

Molecular epidemiology and evolution of A(H1N1)pdm09 and H3N2 viruses in Jordan, 2011–2013

R. El Shesheny¹, N.B. Halasa², J.V. Williams³, A.A. Shehabi⁴, S. Faouri⁵, G. Kayali⁶ and N. Khuri-Bulos⁷

الوبائيات الجزيئية وتطور فيروسات A(H1N1)pdm09 و H3N2 في الأردن

رابح الشيشيني، ناتاشا هلسة، جون ويليامز، عاصم شهابي، سمير فاعوري، غازي كيالي، نجوى خوري-بولس

الخلاصة: لا يوجد سوى القليل من المعلومات عن تطور فيروسات الإنفلونزا الموسمية A(H1N1)pdm09 و H3N2 في الأردن. ولمعالجة هذه القضية ودراسة التغير الجيني والمستضدي لفيروسات A(H1N1)pdm09 و H3N2 أجرينا تحاليل تتعلق بتطور السلالات وبالجينات لجينات HA و NA في فيروسات A(H1N1)pdm09 و H3N2 في الأردن بين عامي 2011 و 2013. وأدرجت في التحليل كامل جينات HA و NA للعينات الـ 16 الإيجابية لـ H1N1 التي تم الحصول عليها في هذه الدراسة، بالإضافة إلى 21 متواليّة HA و 20 متواليّة NA منشورة عن الفيروسات الأردنية التي كانت متاحة في قواعد البيانات الجينية عبر الإنترنت. وبالنسبة لـ H3N2 ولدنا 20 متواليّة HA و 19 متواليّة NA، وأدرجنا في التحليل 19 متواليّة HA و 19 متواليّة NA منشورة. لقد امتلكت فيروسات H1N1 الأردنية الطفرات التي تميز الفئة المستضدية 6، بينما انتمت فيروسات H3N2 إلى الفئة 3. ولم يُكتشف أي صنّاع مقاومة للأوسيلتاميفير. ووصفت الطفرات الفردية بالتفصيل.

ABSTRACT Understanding the genetic evolution of A(H1N1)pdm09 and H3N2 viruses can help better select strains to be included in the annual influenza vaccine. There is little information on their evolution in Jordan so this study investigated the genetic and antigenic variability of A(H1N1)pdm09 and H3N2 viruses in Jordan by performing phylogenetic and genetic analyses of the HA and NA genes of A(H1N1)pdm09 and H3N2 viruses between 2011 and 2013. The full HA and NA genes of 16 H1N1-positive samples obtained in our study and 21 published HA sequences and 20 published NA sequences from Jordanian viruses that were available on online gene databases were analysed. For H3N2, we generated 20 HA and 19 NA sequences and included 19 published HA and NA sequences each in the analysis. Jordanian H1N1 viruses had mutations that are characteristic of antigenic group 6 while H3N2 virus mutations belonged to group 3. No markers of resistance to oseltamivir were detected. The individual mutations are described in detail.

Épidémiologie moléculaire et mutations des virus A(H1N1)pdm09 et H3N2 en Jordanie

RÉSUMÉ La compréhension de l'évolution génétique des virus A(H1N1)pdm09 et H3N2 permet de mieux sélectionner les souches devant être ajoutées au vaccin antigrippal annuel. Peu de renseignements sont disponibles sur les mutations des virus saisonniers de la grippe A(H1N1)pdm09 et H3N2 en Jordanie. Afin de remédier à ce problème et d'étudier les variations génétiques et antigéniques des virus A(H1N1)pdm09 et H3N2, nous avons procédé à des analyses génétiques et phylogénétiques des gènes de l'hémagglutinine (HA) et de la neuraminidase (NA) de ces virus, sur la période 2011-2013 en Jordanie. L'analyse a porté sur les séquences complètes des gènes de l'HA et de la NA de 16 échantillons positifs au virus H1N1 prélevés dans le cadre de cette étude, ainsi que sur 21 séquences publiées de l'HA et 20 séquences publiées de la NA, issues de virus jordaniens disponibles sur les bases de données de gènes en ligne. Pour le virus H3N2, nous avons généré 20 séquences de l'HA et 19 de la NA, et avons également inclus dans l'analyse 19 séquences publiées de l'HA et 19 de la NA. Les virus H1N1 jordaniens présentaient des mutations caractéristiques du groupe antigénique 6, tandis que les virus H3N2 appartenaient au groupe 3. Aucun marqueur de résistance à l'oseltamivir n'a été détecté. Les mutations individuelles sont décrites en détail.

¹Centre of Excellence for Influenza Viruses, National Research Centre, Giza, Egypt. ²Department of Pediatrics, Vanderbilt University, Nashville, Tennessee, United States of America. ³Department of Pediatrics, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America.

⁴Department of Pathology-Microbiology, University of Jordan, Amman, Jordan. ⁵Paediatric Division, Ministry of Health, Amman, Jordan. ⁶Human Link, Baabda, Beirut, Lebanon. ⁷Department of Paediatrics and Division of Paediatric Infectious Diseases. University of Jordan, Amman, Jordan (Correspondence to N. Khuri-Bulos: najwa.khuri@gmail.com).

Received: 28/04/16; accepted: 22/08/16

Introduction

Influenza viruses are considered one of the most common causes of respiratory infection among humans and they are associated with high morbidity and mortality (1). Influenza A viruses are classified based on the antigenic properties of the 2 surface glycoproteins, haemagglutinin (HA) and neuraminidase (NA) (2). Alterations in these proteins are key determinants of the pathogenicity of influenza viruses and their ability to avoid the host immune response.

The influenza pandemic of 2009 was caused by a novel strain of influenza A(H1N1)pdm09, which was first identified in Mexico and then spread globally. It had a massive impact on the health and economy of all affected regions (3, 4). Genetic analysis confirmed that it was a triple reassorted virus, with polymerase basic (PB)1 from viruses of human origin, HA, nucleoprotein (NP) and non-structural (NS) genes from classical swine virus, PB2 and polymerase A (PA) from North American avian viruses and NA and matrix (M) from Eurasian swine avian-like viruses (5,6). In Jordan, the first 2 confirmed case of influenza strain A(H1N1)pdm09 was reported on 17 June 2009 (7).

The A(H1N1)pdm09 virus gradually replaced the seasonal H1N1 virus and has co-circulated with H3N2 and influenza B virus since 2010. H3N2 viruses have been circulating in humans since 1968 (8). Molecular analysis and antigenic characterization of circulating A(H1N1)pdm09 strains and H3N2 viruses has classified them into genetic group and subgroup (9,10).

Understanding the evolution of A(H1N1)pdm09 and H3N2 viruses is essential for studying diversification, determining genetic and antigenic relations and choosing vaccine strains to be included in the annual influenza vaccine. As there is little information on the evolution of seasonal influenza A(H1N1)pdm09 and H3N2 viruses

in Jordan, we performed a phylogenetic analysis and genetic analysis of the HA and NA genes of A(H1N1)pdm09 and H3N2 viruses in Jordan between March 2010 and March 2013.

Methods

Samples

Nasal and throat swabs were collected as part of a large prospective cohort study of 3168 children < 2 years of age who presented with fever and/or respiratory symptoms to a Government hospital in Amman, Jordan, in 2010–2013. Of those, 71 children tested positive for influenza A infection. Details of the cohort study design have been published previously (11, 12).

RNA extraction and PCR

Amplification and sequencing of HA and NA genes

RNA was extracted from original swabs using QIAGEN viral extraction kit (Qiagen, Hilden, Germany). One-step real-time reverse transcriptase polymerase chain reaction (RT-PCR) was performed with Invitrogen Superscript One-Step RT-PCR and a Platinum Taq kit (Invitrogen, Carlsbad, California, USA), with full-length primers, as described previously (13). The PCR products were separated by 1% agarose gel electrophoresis. Amplicons of the appropriate sizes were subsequently excised from the gel and purified with a QIAGEN gel extraction kit (Qiagen). The purified PCR products were used directly for cycle sequencing reactions in a BigDyeR Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) according to the manufacturer's instructions, and were further amplified for 26 cycles at 95 °C, 30 s; 50 °C, 15 s; and 60 °C, 4 min. The reaction product was purified by exclusion chromatography on CentriSep columns (Princeton Separations, Adelphia, NJ, USA). The recovered materials were sequenced in an ABI 3730XL DNA

analyser (Applied Biosystems, Foster City, CA, USA). Sequences were generated with SeqMan DNA Lasergene 7 software (DNASTAR, Madison, Wisconsin, USA). The sequences were submitted to GenBank under accession numbers KU933838–KU933908.

Sequence analysis and phylogenetic tree construction

DNA Lasergene 7 and BioEdit 7.0 (14) were used for multiple sequence alignment and genomic signature analysis with the ClustalW algorithm (15). The full genes of 16 H1N1-positive samples obtained in this study, in addition to 21 published HA sequences and 20 published NA sequences from Jordanian viruses that were available on GenBank and the Global Initiative on Sharing All Influenza Data (GISAID), were included in the analysis. For H3N2, we generated 20 HA and 19 NA sequences and included 19 published HA sequences and 19 published NA sequences in the analysis.

MEGA 6 was used for phylogenetic tree construction by applying the neighbour-joining method with Kimura's 2-parameter distance model and the reliability of phylogenetic inference at each branch node was estimated by the bootstrap method with 1000 replications (16). Sequence group designation was according to WHO nomenclature (17).

Results

H1N1

The results of phylogenetic analysis of the HA and NA genes from H1N1 viruses from Jordan are shown in Figures 1 and 2. A/California/07/09 was used as the root for the tree.

Phylogenetic analysis of the HA gene of pandemic H1N1 viruses revealed that the viruses in Jordan in 2011–2013 viruses belonged to three antigenic groups: 5, 6 and 7 (Figure 1). Antigenic group 6 included isolates from




Figure 1 Phylogenetic tree of the NA gene of pandemic influenza A (H1N1) viruses isolated in Jordan. Vaccine strains are in bold italics and reference strains are in bold. Viruses sequenced for this study are in red font colour.




Figure 2 Phylogenetic tree of the HA gene of pandemic influenza A (H1N1) viruses isolated in Jordan. Vaccine strains are in bold italics and reference strains are in bold. Viruses sequenced for this study are in red font colour.




Figure 3 Phylogenetic tree of the NA gene of influenza A (H3N2) viruses isolated in Jordan. Vaccine strains are in bold italics and reference strains are in bold. Viruses sequenced for this study are in red font colour.




Figure 4 Phylogenetic tree of the HA genes of influenza A (H3N2) viruses isolated in Jordan. Vaccine strains are in bold italics and reference strains are in bold. Viruses sequenced for this study are in red font colour.

2012–2013. Seven isolates belonged to group 6.C, which is characterized by mutations V234I, K283E and E172K. Six viruses from 2012 belonged to antigenic group 6.A, which is characterized by mutations H138R and V249L. One isolate from 2012 (A/Jordan/B1713/2012) belonged to antigenic group 7, which had mutation A197T. One isolate (A/Jordan/A0088/2011) from 2011 belonged to antigenic group 5, which shares two mutations, V234I and K283E, with group 6.C.

The genetic analysis (Table 1 online) showed that all the viruses in this study had one mutation at antigenic site Sb (S185T) and one at antigenic site Ca1 (S203T). Viruses belonging to antigenic group 6.A had one mutation at antigenic site Ca2 (H148R). One virus had mutation L161I in antigenic site Sa. No changes in the glycosylation sites were detected.

Phylogenetic analysis of the NA gene from pandemic H1N1 viruses revealed that the 2011–2013 viruses belonged to 2 antigenic groups, groups 5 and 6 (Figure 2). Analysis of the antiviral susceptibility of Jordanian strains was based on the presence of mutations encoding amino acid substitutions at key residues that confer resistance to NA inhibitors. None of these viruses displayed oseltamivir-resistant clinical markers at positions D198N, I222R, H274Y, N294S.

H3N2

The results of phylogenetic analysis of the HA and NA genes from H3N2 viruses from Jordan during 2011–2012 are shown in Figures 3 and 4. A/Brisbane/10/2007 was used as the root for the tree. Phylogenetic analysis of H3 showed that the viruses belonged to the A/Victoria/361/2011 genetic clade and fell into 2 antigenic groups: 3 and 6. Viruses in group 3 had mutations V223I and D158N and were further divided into 2 groups, 3B and 3C. One isolate belonged to antigenic group 6, which contains mutations D53N,

Y94H, I230V and E280A. Two viruses from 2011 had 3 common mutations, G49A, K92R and V347M, and were grouped together.

The H3N2 viruses in our study accumulated 16 mutations as compared with the reference vaccine strain A/Perth/16/2009 (Table 2 online). The mutations included 4 antigenic site A mutations (I140M, R142G, K144N and N145S), 1 antigenic site B mutation (A198S/P), 4 antigenic site C mutations (D53N, N278K, E280A/P/S and N312S), 4 antigenic site D mutations (T212A, S214I/V, V223I and I230V) and 3 antigenic site E mutations (T48A/I, K62E/I and Y94H).

At the potential N-linked glycosylation sites of H3N2 viruses, amino acid 45 (SSS) was mutated to (NSS) in 4 strains, which created an additional glycosylation site. Two isolates lacked the glycosylation site at amino acid 63. Most isolates lost a glycosylation site at amino acid 402 in the NA gene (Table 3 online).

Discussion

All the influenza H1N1 viruses detected in the world after pandemic 2009 have been related to A(H1N1)pdm09. Genetic variations have been observed in circulating A(H1N1)pdm09 viruses, but they remain antigenically similar to the A/California/07/2009 vaccine strain (18). Most isolates of A(H1N1)pdm09 in our study belonged to antigenic subgroups 6A and 6C. In the case of A(H3N2), all viruses from the 2011–2012 season sequenced in the study belonged to the A/Victoria/208/2011-like lineage, and most fell into antigenic subgroup 3B. Similar results were found in isolates in neighbouring Lebanon (19). The Jordanian isolates closely resembled the WHO-designated reference strains A/Stockholm/18/2011 and A/England/259/2011, with amino acid substitutions in HA1/HA2

(N145S and V223I A198S, N312S and D158N) (20).

Patterns of antigenic site variation were observed by amino acid alignment of the HA genes of A(H1N1)pdm09 and H3N2 viruses. In A(H1N1)pdm09, amino acid mutations were detected at four antigenic sites (Sa, Sb, Ca2 and Ca1), while A(H3N2) showed mutation at antigenic sites (A, B, C, D, and E). N-Glycosylation affects not only the antigenic and functional properties of surface proteins but can also provide an opportunity for immune escape (21,22). The 3 glycosylation sites (residues 146, 367 and 402) on the head of the NA monomer encoded by H3N2 strains are located around the enzymatic active site (23). We observed that most isolates lost a glycosylation site at 402 in the NA gene of A(H3N2).

In the 2014–2015 season, antigenic drift of subtype H3N2 decreased vaccine effectiveness. The H3N2 strain of the 2014–2015 influenza vaccine was A/Texas/50/2012, which belongs to antigenic subgroup 3C.1 (24). In September 2014, the WHO consultation and information meeting on the composition of influenza virus vaccines indicated the emergence of 2 new genetic subgroups, 3C.2a and 3C.3a, of A(H3N2) viruses containing antigenic drift viruses of previously circulating strains (25). WHO recommended that the H3N2 component be updated with an A/Switzerland/9715293/2013-like (antigenic subgroup 3C.3a) virus for the 2015–2016 vaccine for the Northern Hemisphere (26).

Antigenic and genetic analyses are invaluable in guiding effective influenza vaccine selection and control measures. The results of this study contribute to understanding the genetic and antigenic evolution of A(H1N1)pdm09 and H3N2 viruses in Jordan. Further investigation is required to better understand how genetic and antigenic diversity and how to utilize it for vaccine strain selection.

Acknowledgements

We wish to acknowledge our research staff who were responsible for subject enrolment into the cohort: Hanan Amin, Amani Altaber, Hana'a Khalaf, Isra'a Kharbat, Darin Yasin, and Shireen Issa.

Funding: This work was funded by the National Institute of Allergy and

Infectious Diseases, National Institutes of Health, US Department of Health and Human Services (under contract HH-SN272201400006C); the Egyptian Science and Technology Development Fund (under contract no. 5175); and by support from the American Lebanese Syrian Associated Charities. All phases of this project were supported by Union Bank of Switzerland (UBS) Optimus Foundation. The project publication

described was supported by CTSA award No. UL1TR000445 from the National Center for Advancing Translational Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

Competing interests: None declared.

All tables available online at: <http://www.emro.who.int/emh-journal/eastern-mediterranean-health-journal/home.html>

References

- Cox NJ, Subbarao K. Global epidemiology of influenza: past and present. *Annu Rev Med.* 2000;51:407-21.
- Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiol Rev.* 1992 Mar;56(1):152-79.
- Girard MP, Tam JS, Assossou OM, Kieny MP. The 2009 A (H1N1) influenza virus pandemic: a review. *Vaccine.* 2010 Jul 12;28(31):4895-902.
- Keogh-Brown MR, Smith RD, Edmunds JW, Beutels P. The macroeconomic impact of pandemic influenza: estimates from models of the United Kingdom, France, Belgium and The Netherlands. *Eur J Health Econ.* 2010 Dec;11(6):543-54.
- Garten RJ, Davis CT, Russell CA, Shu B, Lindstrom S, Balish A, et al. Antigenic and genetic characteristics of swine-origin 2009 A(H1N1) influenza viruses circulating in humans. *Science.* 2009 Jul 10;325(5937):197-201.
- Peiris JS, Poon LL, Guan Y. Emergence of a novel swine-origin influenza A virus (S-OIV) H1N1 virus in humans. *J Clin Virol.* 2009 Jul;45(3):169-73.
- Rihani R, Hayajneh W, Sultan I, Ghatasheh L, Abdel-Rahman F, Hussein N, et al. Infections with the 2009 H1N1 influenza virus among hematopoietic SCT recipients: a single center experience. *Bone Marrow Transplant.* 2011 Nov;46(11):1430-6.
- Smith DJ, Lapedes AS, de Jong JC, Bestebroer TM, Rimmelzwaan GF, Osterhaus AD, et al. Mapping the antigenic and genetic evolution of influenza virus. *Science.* 2004 Jul 16;305(5682):371-6.
- Nelson M, Spiro D, Wentworth D, Beck E, Fan J, Ghedin E, et al. The early diversification of influenza A/H1N1pdm. *PLoS Curr.* 2009 11 03;1:RRN1126. PMID:20029664
- Holmes EC, Ghedin E, Miller N, Taylor J, Bao Y, St George K, et al. Whole-genome analysis of human influenza A virus reveals multiple persistent lineages and reassortment among recent H3N2 viruses. *PLoS Biol.* 2005 Sep;3(9):e300.
- Halasa N, Williams J, Faouri S, Shehabi A, Vermund SH, Wang L, et al. Natural history and epidemiology of respiratory syncytial virus infection in the Middle East: Hospital surveillance for children under age two in Jordan. *Vaccine.* 2015 Nov 25;33(47):6479-87.
- Schuster JE, Khuri-Bulos N, Faouri S, Shehabi A, Johnson M, Wang L, et al. Human Metapneumovirus Infection in Jordanian Children: Epidemiology and Risk Factors for Severe Disease. *Pediatr Infect Dis J.* 2015 Dec;34(12):1335-41.
- Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. Universal primer set for the full-length amplification of all influenza A viruses. *Arch Virol.* 2001 Dec;146(12):2275-89.
- Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis for Windows 95/98/NT. *Nucleic Acids Symp.* 1999; 41:95-8.
- Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994 Nov 11;22(22):4673-80.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* 2013 Dec;30(12):2725-9.
- ECDC. Influenza Virus Characterization. Surveillance Report 2014 (<http://ecdc.europa.eu/en/publications/Publications/Influenza-ERLI-Net-report-Sept-2014.pdf>, accessed 8 August 2016).
- Klimov AI, Garten R, Russell C, Barr IG, Besselaar TG, Daniels R, et al. Writing Committee of the World Health Organization Consultation on Southern Hemisphere Influenza Vaccine Composition for 2012. WHO recommendations for the viruses to be used in the 2012 Southern Hemisphere Influenza Vaccine: epidemiology, antigenic and genetic characteristics of influenza A(H1N1)pdm09, A(H3N2) and B influenza viruses collected from February to September 2011. *Vaccine.* 2012 Oct 5;30(45):6461-71.
- Zaraket H, Dapat C, Ghanem S, Ali Z, Lteif M, Kondo H, et al. Characterization of human Influenza Viruses in Lebanon during 2010-2011 and 2011-2012 post-pandemic seasons. *Intervirology.* 2014;57(6):344-52.
- ECDC. European Centre for Disease Prevention and Control. Influenza virus characterisation, summary Europe, February 2016. Stockholm: ECDC; 2016. 2016 (<http://ecdc.europa.eu/en/publications/Publications/influenza-virus-characterisation-february-2016.pdf>, accessed 8 August 2016).
- Chen W, Zhong Y, Qin Y, Sun S, Li Z. The evolutionary pattern of glycosylation sites in influenza virus (H5N1) hemagglutinin and neuraminidase. *PLoS One.* 2012;7(11):e49224.
- Cherry JL, Lipman DJ, Nikolskaya A, Wolf YI. Evolutionary dynamics of N-glycosylation sites of influenza virus hemagglutinin. *PLoS Curr.* 2009 08 18;1:RRN1001. PMID:20025194
- Fang Q, Gao Y, Chen M, Guo X, Yang X, Yang X, et al. Molecular epidemiology and evolution of A(H1N1)pdm09 and H3N2

- virus during winter 2012-2013 in Beijing, China. *Infect Genet Evol.* 2014 Aug;26:228-40.
24. Broberg E, Snacken R, Adlhoch C, Beauté J, Galinska M, Pereyaslov D, et al.; WHO European Region and the European Influenza Surveillance Network. Start of the 2014/15 influenza season in Europe: drifted influenza A(H3N2) viruses circulate as dominant subtype. *Euro Surveill.* 2015 01 29;20(4):pii 21023. PMID:25655052
 25. WHO Consultation and Information Meeting on the Composition of Influenza Virus Vaccines for the Northern Hemisphere 2014-2015. Geneva: WHO. 17-19 Feb 2014. 2014 (<http://www.who.int/influenza/vaccines/virus/recommendations/consultation201402/en/>, accessed 8 August 2016).
 26. Anonymous. Recommended composition of influenza virus vaccines for use in the 2016 southern hemisphere influenza season. *Wkly Epidemiol Rec.* 2015 Oct 9;90(41):545-59. PMID:26454888 <tabletit>Table 1 Amino acid substitutions observed in antigenic sites of the haemagglutinin protein of (H1N1) influenza viruses isolated in Jordan