


WHO AFGHANISTAN POLIO SNAPSHOT AS OF 30 APRIL 2017



POLIO TRANSMISSION

No new wild poliovirus (WPV1) cases were reported by the regional reference laboratory in April 2017.
One new WPV1 positive environmental sample was reported in April, collected in March, from Nangarhar province.

AFP and ENVIRONMENTAL SURVEILLANCE

• 217 acute flaccid paralysis (AFP) cases were reported in April. Overall in 2017, 960 AFP cases have been reported, of which 718 have been discarded as non-polio AFP and 239 AFP cases are pending classification.

• Non-polio AFP rate is 14, adequate stool rate is 93%, and non-polio enterovirus rate is 14% at the national level.

• In all regions non-polio AFP rate is above 8, stool adequacy above 86%, and non-polio enterovirus rate ranges from 8 to 22%.

SUPPLEMENTARY IMMUNIZATION ACTIVITIES (SIAs)

• Sub-National Immunization Days (SNIDs), with expanded scope to include case response, targeted over 6.7 million children under the age of 5 with OPV in 26 provinces on 17-21 April.

COMPLEMENTARY VACCINATION ACTIVITIES

• A total of 18 cross-border vaccination points, 44 cross-border teams, 369 permanent transit teams (PTTs) and 22 permanent polio teams (PPTs) were operational across Afghanistan in April.

• Permanent transit teams vaccinated 1,116,420 children against polio and cross-border teams vaccinated 66,351 children in April.

• A total of 11,063 children were vaccinated with OPV and 3,822 with IPV by teams in UNHCR repatriation centers and IOM sites receiving returnees from Pakistan and Iran in April.

• From January 2017 to date, 17,114 returnee children have received OPV and 4,798 children have been vaccinated with IPV.

• In compliance with International Health Regulations (IHR), 12,079 international travellers were vaccinated against polio in April.

Data source: WHO

World Health Organization

POLIO: WHO CORE PRIORITIES IN 2016-17

WHO supports polio eradication activities through the planning, implementation, monitoring and evaluation of vaccination campaigns, training of frontline health workers, and through providing technical assistance to the polio programme at all levels. WHO maintains an active acute flaccid paralysis (AFP) and environmental surveillance system to detect polio transmission, and facilitates cross-border coordination in common reservoirs with Pakistan. WHO supports the Ministry of Public Health and works in close partnership with UNICEF, Bill and Melinda Gates Foundation (BMGF), the US Centers for Disease Control and Prevention (CDC) and Rotary International as part of the Global Polio

Eradication Initiative (GPEI).



WHO supports acute flaccid paralysis (AFP) surveillance through a countrywide network of AFP focal points linked with health facilities and community-based reporting volunteers. There are around 730 AFP focal points, over 1,400 active surveillance sites, 2,400 zero reporting sites and more than 28,000 community-based reporting volunteers across the country. An external AFP surveillance review conducted in June 2016 concluded that the surveillance system is strong and the circulation of wild or vaccine-derived poliovirus is unlikely to be missed in Afghanistan.

Active AFP case search is conducted frequently in all health facilities part of the AFP reporting network. Vaccinators are also carrying out active case search during SIAs. In 2016/17, the AFP reporting network

will continue to be reviewed and expanded based on needs and emphasis will be on including healthcare providers serving high-risk population groups as well as insecure areas, high-risk areas and districts with low non-polio AFP rates.

The environmental surveillance system includes 17 sampling sites in Nangarhar, Kunar, Kandahar, Helmand, Kabul and Khost provinces.

SUPPLEMENTARY IMMUNIZATION ACTIVITIES (SIA)



WHO is responsible for technical assistance, trainings, micro-planning, post-campaign assessment surveys, data analysis and reporting for SIAs. New initiatives put in place in 2016 to improve SIA quality include the roll-out of a new FLW training curriculum, a modified re-visit strategy to reach missed children, the development of district profiles and district-specific plans, the in-depth investigation of reasons for 'lot failure' in Lot Quality Assurance Sampling (LQAS) surveys, the strategic use of inactivated polio vaccine (IPV) and the field validation and revision of microplans.

The programme's current focus is on 50 very high-risk districts (VHRDs). In 2016/17, the oral polio vaccine (OPV) SIA schedule includes two

national immunization days (NIDs) and two sub-national immunization days (SNIDs) in the second half of 2016 and four NIDs and six SNIDs in 2017. The SIA dates will be synchronized with Pakistan. For every new case detected, three case response campaigns will be carried out.

HUMAN RESOURCES AND CAPACITY DEVELOPMENT



WHO supports the Ministry of Public Health in polio eradication through a country-wide network of international and national staff. Each regional team consists of international medical officers (except Central and Northeastern Regions), regional polio officers, assistant regional polio officers, provincial polio officers and district polio officers.

More than 5,500 cluster supervisors and over 58,000 volunteers conduct house-to-house oral polio vaccinations for children below five years of age during national immunization campaigns. There are around 1,000 intra-campaign monitors and 1,100 post-campaign

monitors to monitor the activities during campaigns and assess coverage after the campaign.

WHO supports the training of frontline workers (FLWs) before each vaccination campaign with a newly-revised curriculum. In 2016/17, the programme will focus on improving team performance by ensuring FLWs are carefully selected using a transparent approach, and that they receive high-quality training to keep them motivated.

COMPLEMENTARY VACCINATION ACTIVITIES



Over 320 Permanent Transit Teams (PTTs) vaccinate children who travel in and out of security-compromised areas and children traveling to other destinations. Eighteen cross-border vaccination points in border areas with Pakistan and Iran ensure that children crossing the border are immunized. In selected locations, permanent polio teams (PPTs) provide OPV to children on a continuous basis in their assigned catchment area, in addition to the planned house-to-house vaccination during SIAs.

In 2016/17 the programme will continue to assess and modify the number and location of PTTs according to needs and the evolving

accessibility situation. Full synchronization with vaccination operations on the Pakistan side of the border will continue for cross-border teams (CBTs). Special campaigns are carried out in the South-Eastern, Southern and Western Regions targeting nomads who move widely in the country and across borders.

Vaccination teams are also stationed at airports to administer OPV to travellers as per recommendations under International Health Regulations. Teams are also stationed in UNHCR and IOM sites for returnee refugees from Pakistan and Iran to provide OPV, IPV and measles vaccination to children.

WHO Afghanistan UNOCA Compound, Jalalabad Road Pul-e-Charkhi Kabul, Afghanistan T: +93700045276 E: emacoafgwr@who.int





