


OVERALL SITUATION

Poor recovery from the drought that affected Somalia in 2016/2017 as well as the negative impact of low *Gu* rains (April–June 2019) and low *Deyr* rains (October–December 2018) have led to the 2.2 million in need of urgent humanitarian assistance.

Adverse climatic conditions combined with other drivers of humanitarian crisis, such as armed conflict, have led to internal displacement and weakening of the fragile health system which is contributing to the increased number of cases of measles and diarrhoea.



CHOLERA IN DROUGHT-AFFECTED DISTRICTS

Since December 2017, cholera cases have continued to be reported in Somalia. Among the drought-affected districts, active transmission of cholera has only been reported in Banadir region. A total of 1 508 cholera cases were reported from eight districts of Banadir affected by drought during epidemiological weeks 1 to 42, 2019 (Table 1). Of the 820 stool samples tested since December 2017, 162 samples tested positive for *Vibrio cholerae* serotype Ogawa. Culture and sensitivity studies at the National Public Health Laboratory show that the *V. cholerae* serotype Ogawa isolated is sensitive to chloramphenicol and tetracycline but resistant to ampicillin and nalidixic acid.

ACUTE DIARRHOEAL DISEASES

Cases of acute diarrhoea have increased in 2019 compared with previous years; this is linked to the shortage of safe water, and poor hygiene and sanitation (Fig. 1). Since epidemiological week 1, a total of 107 696 cases of acute diarrhoea have been reported from drought-affected districts through EWARN. The most affected districts are Baidoa, Lasanood, Marka, Beletweyne and Burco. (See Table 1).

MEASLES

As a result of the mass measles vaccination campaign conducted in Somalia in 2018, the number of suspected cases of measles has decreased in 2019 compared with previous years (Fig. 2). Since epidemiological week 1, 2019, a total of 1 927 suspected cases of measles have been reported in drought-affected districts. Madina, Jowhar, Marka, Beletweyne and Baidoa are the most affected districts. (See Table 1). Since week 23, the number of suspected cases of measles has increased and is higher than the number of cases reported during the same period in 2017 and 2018. This may be because of the gradual reduction in the monthly coverage of children vaccinated against measles and the presence of risk factors such as malnutrition in drought-affected districts (Fig. 3).

A total of 153 565 (79.6%) children under 1 year of age out of the targeted 192 825 received Measles 1 vaccine (MCV1) in drought-affected districts from January to September 2019 (Fig. 3).

POLIO UPDATES

No new cases of circulating vaccine-derived polio virus type 2 (cVDPV2) were confirmed this week. Between epidemiological weeks 1 and 41, three new cVDPV2 cases were confirmed in Somalia (Map). The most recent case of cVDPV2 was confirmed on 8 May 2019.

No new cases of circulating vaccine-derived polio virus type 3 (cVDPV3) reported from acute flaccid paralysis cases in 2019. The last case of cVDPV3 in Somalia was confirmed on 7 September 2018.

All environmental samples were negative for both cVDPV2 and cVDPV3 in 2019.

The first round of the response campaign to the three cases of cVDPV2 was conducted from 24 to






Fig. 2. Trends of measles cases reported in drought-affected districts of Somalia, 2017–2019





Fig. 3. Number of children under 1 year of age vaccinated against measles by month, 2019



27 June 2019 in 21 target districts of Somaliland and Puntland using the monovalent type 2 oral poliovirus vaccine (mOPV2).

A total of 170 072 (98%) under five years children were vaccinated using mOPV2 in the targeted 12 districts of Somaliland and Puntland.

A total of 1.59 million (97%) under 5 children were vaccinated using bivalent oral polio vaccine (bOPV) in 41 high- risk districts from 23 to 26 September 2019 across Somalia.

Published by World Health Organization (WHO), Country Office, Mogadishu, Somalia

For Correspondence: Tel: +252616695096; Email address: emacosomwr@who.int; URL: http://www.emro.who.int/countries/som/index.html

Table 1. Cumulative numbers of diarrhoea, measles and cholera cases in drought-affected districts of Somalia (epidemiological weeks 1–42, 2019)^a

State/region	Districts	acute diarrhoea cases	Suspected measles cases	Suspected cholera cases
Banadir ^b	Daynile	4734	116	375
	Hawal Wadag	1600	39	71
	Hodan	225	9	433
	Kahda	3480	59	73
	Karan	45	0	35
	Madina/Wadajir	2806	536	438
	Waberi	574	30	48
	Yaqshid	1392	39	35
Galmudug	Adado	828	13	0
	Dusamareeb	608	17	0
	Abudwaq	242	2	0
HirShabelle	Balad	2006	68	0
	Jowhar	4576	195	0
	Belet Weyne	8005	138	0
Juba land	Kismayo	4796	97	0
	Garbahare	1221	9	0
Puntland	Garowe	5075	15	0
	Bossaso	4500	36	0
	Qardho	1879	4	0
	Galkayo	5741	38	0
Somaliland	Erigavo	3246	5	0
	Hargeisa	3737	31	0
	Las Anod	13891	31	0
	Burao	8659	45	0
South West state	Wajid	395	0	0
	Hudur	910	0	0
	Baidoa	13982	213	0
	Marka	8543	142	0
Total		107 696	1 927	1 508

^a The total number of cases reported on EWARN may change after verification by surveillance teams.

^b Banadir is a region not a state.

WHO and the Federal Ministry of Health continue to monitor trends of epidemic-prone diseases in drought-affected districts using the electronic EWARN. WHO and health cluster partners are implementing preparedness and response activities to prevent the negative effects of drought. WHO is also supporting different states to increase the number of health facilities submitting alerts of epidemic-prone diseases in EWARN. With support from Central Emergency Response Fund (CERF), WHO in collaboration with state level health authorities are implementing activities to avert the negative consequence of drought in selected districts of Jubbaland, Southwest state and Hirshabelle.