




For epidemic prone diseases in Somalia for week 49, 02-08 December, 2019

OVERALL SITUATION

Poor recovery from the drought that affected Somalia in 2016/2017 as well as the negative impact of low *Gu* rains (April–June 2019) and low *Deyr* rains (October–December 2018) have led to the 2.2 million in need of urgent humanitarian assistance.

Adverse climatic conditions combined with other drivers of humanitarian crisis, such as armed conflict, have led to internal displacement and weakening of the fragile health system which is contributing to the increased number of cases of measles and diarrhoea.



diarrhoea, circulating vaccine-derived polio virus (cVDPV) type 2 and measles DJIBOUTI **ETHIOPIA** Drought Drought affected districts Other districts Acute diarrhoea cases 1 - 1000 1001 - 2000 2001 - 3000 **KENYA** Over 3000 cVDPV cases **Banadir Region** cVDPV2 Suspected measles cases 1-30 31-50 51-100 101 - 481 The designation employed and presentation of material on the map do not imply the expression on any opinion whatsoever or United Nations concerning the legal status of any country, territory, city or any any area of authority, or concerning determinal

Map showing drought-affected areas of Somalia, and locations of cases of

¹ This data is from drought affected districts only

CHOLERA IN DROUGHT-AFFECTED DISTRICTS

Since December 2017, cholera cases have continued to be reported in Somalia. Among the drought-affected districts, active transmission of cholera has only been reported in Banadir region. A total of 1959 cholera cases and one death were reported from eight districts of Banadir affected by drought during epidemiological weeks 1 to 49, 2019 (Table 1). Of the 895 stool samples tested since December 2017, 197 samples tested positive for *Vibrio cholerae* serotype Ogawa. Culture and sensitivity studies performed at the National Public Health Laboratory show that the *V. cholerae* serotype Ogawa isolated is sensitive to chloramphenicol and tetracycline but resistant to ampicillin and nalidixic acid.

ACUTE DIARRHOEAL DISEASES

Cases of acute diarrhoea have increased in 2019 compared with previous years; this is linked to the shortage of safe water, and poor hygiene and sanitation (Fig. 1). Since epidemiological week 1, a total of 126 618 cases of acute diarrhoea have been reported from drought-affected districts through EWARN. The most affected districts are Baidoa, Lasanood, Marka, Beletweyne and Burco. (See Table 1).

MEASLES

As a result of the mass measles vaccination campaign conducted in Somalia in 2018, the number of suspected cases of measles have decreased in 2019 compared with previous years (Fig. 2). Another measles campaign conducted in November 2019 is expected to further contribute to reduction of measles cases. epidemiological week 1, 2019, a total of 2337 suspected cases of measles have been reported in drought-affected districts. Madina, Jowhar, Marka, Beletweyne and Baidoa are the most affected districts. (See Table 1). Since week 23, the number of suspected cases of measles has increased and is higher than the number of cases reported during the same period in 2018. This may be because of the gradual reduction in the monthly coverage of children vaccinated against measles and the presence of risk factors such as malnutrition in drought-affected districts (Fig. 3).

A total of 126 653 (74%) children under 1 year of age out of the targeted 171 400 received measles 1 vaccine (MCV1) in drought-affected districts from March to October 2019 (Fig. 3). During the drought monitoring period, March to October 2019, the vaccination coverage has been ranging between 61% and 83% per month against a monthly target of 21 425 children under 1 year of age.

POLIO UPDATES

No new cases of circulating vaccine-derived polio virus type 2 (cVDPV2) were confirmed this week. Between epidemiological weeks 1 and 48, three new cVDPV2 cases were confirmed in Somalia (Map). The most recent case of cVDPV2 was confirmed on 8 May 2019.

No new cases of circulating vaccine-derived polio virus type 3 (cVDPV3) reported from acute flaccid paralysis cases in 2019. The last case of cVDPV3 in Somalia was confirmed on 7 September 2018.

All environmental samples were negative for both cVDPV2 and cVDPV3 in 2019.

Fig. 1. Trends of acute diarrhoea cases reported in drought-affected districts of Somalia, 2017–2019



Jan 2019 - December 2019

1000

Fig. 2. Trends of measles cases reported in drought-affected districts of Somalia, 2017–2019

11 13 15 17 19 21 23 25 27 29 31 33 35 37 39 41 43 45 47 **Epidemiological week**




Fig. 3. Number of children under 1 year of age vaccinated against measles by month, 2019



Integrated measles and polio vaccination campaign was conducted in Southwest, Jubbaland, Hirshabelle and Galmudug states from 24 to 28 November. WHO polio and emergency team jointly coordinated the implementation of the campaign. A total of 918 338 under five years children received polio vaccine, 825 654 received measles vaccine, 54 8917 received deworming and 820 052 received Vitamin A supplementation. Post campaign evaluation results are in the process of compilation, once completed then the final results will be shared.

Table 1. Cumulative numbers of diarrhoea, measles and cholera cases in drought-affected districts of Somalia (epidemiological weeks 1-49, 2019)^a

State/region	Districts	acute diarrhoea cases	Suspected measles cases	Suspected cholera cases
Banadir ^b	Daynile	5 863	137	506
	Hawal Wadag	1 860	44	88
	Hodan	249	9	533
	Kahda	3 890	64	103
	Karan	45	0	42
	Madina/Wadajir	3 294	719	573
	Waberi	800	30	75
	Yaqshid	1 521	43	39
Galmudug	Adado	1 017	31	0
	Dusamareeb	818	41	0
	Abudwaq	345	2	0
HirShabelle	Balad	2 115	75	0
	Jowhar	5 338	227	0
	Belet Weyne	9 400	138	0
Juba land	Kismayo	5 379	107	0
	Garbahare	1 729	9	0
Puntland	Garowe	6 013	21	0
	Bossaso	5 550	38	0
	Qardho	2 160	7	0
	Galkayo	6 433	40	0
Somaliland	Erigavo	3 822	6	0
	Hargeisa	4 585	33	0
	Las Anod	16 929	34	0
	Burao	9 538	47	0
South West state	Wajid	495	0	0
	Hudur	1 133	0	0
	Baidoa	16 623	226	0
	Marka	9 674	209	0
Total		126 618	2 337	1 959

^a The total number of cases reported on EWARN may change after verification by surveillance teams.

WHO and the Federal Ministry of Health continue to monitor trends of epidemic-prone diseases in drought-affected districts using the electronic EWARN. WHO and health cluster partners are implementing preparedness and response activities to prevent the negative effects of drought. WHO is also supporting different states to increase the number of health facilities submitting alerts of epidemic-prone diseases in EWARN. With support from Central Emergency Response Fund (CERF), WHO in collaboration with state level health authorities are implementing activities to avert the negative consequence of drought in selected districts of Jubbaland, Southwest state and Hirshabelle.

^b Banadir is a region not a state.

FLOOD MONITORING

During week 49, health facilities in flood and drought-affected districts reported three alerts of: acute watery diarrhea (140 cases), malaria (416 cases) and measles (43 cases). The alerts were verified through field investigation by WHO deployed rapid response teams (RRT). A total of 15 malaria cases were verified as true. Rapid diagnostic testing was done to test for malaria in the flood affected districts. All cases were treated and referred to the nearest health center for further management.

Reporting week	Alert description	No. Of alerts reported	No. Of alerts verified	No. Of true alerts
Week 46	Acute watery diarrhoea (AWD)	80	5	1
	bloody diarrhoea	16	1	0
	Malaria	73	15	7
	Measles	27	1	0
Total week 46		196	22	8
Week 47	Acute watery diarrhoea (AWD)	93	5	2
	bloody diarrhoea	22	2	0
	Malaria	93	17	6
	Measles	38	1	0
Total week 47		246	25	8
Week 48	Acute watery diarrhoea (AWD)	104	5	0
	bloody diarrhoea	37	3	0
	Malaria	100	20	15
	Measles	32	2	0
	Total week 48	273	30	15
Week 49	Acute watery diarrhoea (AWD)	140	5	2
	bloody diarrhoea	0	0	0
	Malaria	416	5	2
	Measles	43	5	4
	Total week 49	345	15	8