




## **OVERALL SITUATION**

Poor recovery from the drought that affected Somalia in 2016/17 as well as the negative impact of low *Gu* rains (April–June 2019) and low *Deyr* rains (October–December 2018) have led to the 2.2 million in need of urgent humanitarian assistance.

Adverse climatic conditions combined with other drivers of humanitarian crisis such as armed conflict leading to internal displacement and weakening of the fragile health system are contributing to increase number of measles and diarrhoeal cases.





## **CHOLERA IN DROUGHT AFFECTED DISTRICTS**

Since December 2017, Cholera cases continue to be reported in Somalia. Among the drought affected districts, active transmission of cholera is reported only in Banadir region. A total of 1,276 cholera cases have been reported from 8 districts of Banadir affected by drought during epidemiological week 1 to 33, 2019 (table 1). Of the 790 stool samples tested since December 2017, a total of 153 samples tested positive for *Vibrio cholerae* serotype Ogawa.

### ACUTE DIARRHOEAL DISEASES SITUATION

Linked to shortage of safe water, poor hygiene and sanitation, the cases of acute diarrhoea have increased in 2019 compared to previous years (fig 1). Since epidemiological week 1, a total of 87,655 cases of acute diarrhoea were reported from drought affected districts through the Early Warning Alert and Response Network (EWARN). However, there is a progressive reduction in the number of new cases of acute diarrhoea reported since week 24. This may be attributed to improved implementation of activities aimed at improved access to safe water and sanitation to drought affected communities. The most affected districts include Baidoa, Lasanood, Marka, Beletweyne and Burco. Please see table-1 and map.

### **MEASLES SITUATION**

Linked to the mass measles vaccination campaign that was conducted in Somalia in 2018, the number of suspected cases of measles has reduced in 2019 when compared to previous years (fig 2). Since epidemiological week 1 of 2019, a total of 1,467 suspected cases of measles were reported in drought affected districts with Madina, Jowhar, Marka, Beletweyne and Baidoa being the most affected districts. Please see table-1 and map. Since week 23, the number of suspected cases of measles have increased above the numbers reported during the same period of time in 2017 and 2018. This may be attributed to gradual reduction in the number of children vaccinated against measles and the presence of risk factors such as malnutrition in drought affected districts (fig-3).

A total of 119,912 (82%) out of 146,426 under one year children targeted received Measles 1 Vaccine (MCV1) in drought affected districts from January to July 2019 (fig 3).

# **POLIO UPDATES**

No new case of circulating Vaccine Derived Polio Virus type 2 (cVDPV2) confirmed this week. Between epidemiological week 1-32, a total of 3 new cVDPV2 cases have been confirmed in Somalia (please see map). The most recent case of cVDPV2 was confirmed on 8 May 2019.

No new circulating Vaccine Derived Polio Virus type 3 (cVDPV3) reported from AFP case in 2019. The last case of cVDPV3 in Somalia was confirmed on 7 September 2018.

All Environmental Samples (ES) were negative for both cVDPV2 and cVDPV3 in 2019.

The first round of response campaign to 3 cases of

Fig 1. Trends of acute diarrhoea cases reported in drought affected districts of Somalia during the same corresponding weeks of 2017, 2018 and 2019

Fig 2. Trends of measles cases reported from drought affected districts of Somalia during the same corresponding weeks of 2017, 2018 and 2019

wk 19 21 23 25 27 29 31 33 35 37

**Epidemiological week** 





Fig 3. Number of children under 1 year vaccinated against measles by month, 2019



cVDPV2 was conducted from 24 to 27 June 2019 in 21 target districts of Somaliland and Puntland using monovalent Type 2 OPV (mOPV2) vaccine.

Of the 945,480 children under five years targeted for vaccination, a total of 992,629 (105%) received mOPV2

The second round of response campaign will commence from 19th to 22nd August 2019 in 34 districts of Somaliland and Puntland using monovalent type 2 OPV (mOPV2) vaccine targeting 1.1 million children under five years.

Table 1. Cumulative number of acute diarrhoea, suspected cholera and suspected measles cases reported from 28 drought affected districts of Somalia (Epidemiological week 1-33 of 2019)

State/region	Districts	acute diarrhoea cases	Suspected measles cases	Suspected cholera cases
Banadir <sup>1</sup>	Daynile	3659	62	314
	Hawal Wadag	1133	31	61
	Hodan	183	3	366
	Kahda	3006	51	69
	Karan	33	0	32
	Madina/Wadajir	2142	424	366
	Waberi	357	19	39
	Yaqshid	1090	36	29
Galmudug	Adado	599	2	0
	Dusamareeb	456	0	0
	Abudwaq	203	2	0
HirShabelle	Balad	1750	54	0
	Jowhar	3715	174	0
	Belet Weyne	6366	120	0
Juba land	Kismayo	4057	81	0
	Garbahare	897	8	0
Puntland	Garowe	4567	11	0
	Bossaso	3711	33	0
	Qardho	1660	3	0
	Galkayo	4961	33	0
Somaliland	Erigavo	2601	3	0
	Hargeisa	2981	28	0
	Las anod	10708	28	0
	Burao	7456	32	0
South West state	Wajid	310	0	0
	Hudur	649	0	0
	Baidoa	10991	117	0
	Markat	7414	112	0
Total		87,655	1,467	1,276

<sup>&</sup>lt;sup>1</sup> Banadir is not a state but a region

Note: The number of cases as shown in table 1 are subject to change after alert verification and validation of data by surveillance officers in different states

WHO and Federal Ministry of Health continue to monitor trends of epidemic prone diseases in drought affected districts. WHO and Health cluster partners are implementing preparedness and response activities to avert the negative consequence of drought. In line with the Global Task Force for Cholera Control (GTFCC) strategy to reduce cholera related deaths by 90% by 2030, Ministry of Health is developing a cholera strategy to ensure that this vision in realised in Somalia.