

Association between oral health knowledge and practices of Saudi pregnant women in Dammam, Saudi Arabia

B.O. Gaffar,¹ M. El Tantawi,¹ A. Al-Ansari¹ and A.S. AlAgl¹

الارتباط بين المعلومات والممارسات المتعلقة بصحة الفم لدى النساء السعوديات الحوامل في الدمام بالمملكة العربية السعودية

بلقيس جعفر، مها الطنطاوي، عاصم الأنصاري، عادل العقل

الخلاصة: لقد قيِّمت هذه الدراسة الارتباط بين المعلومات والممارسات المتعلقة بصحة الفم لدى السعوديات الحوامل تم اختيارهن من بين زائرات مستشفى حكومي في الدمام بالمملكة العربية السعودية عام 2014. حيث أجابت النساء على أسئلة حول المعلومات المتعلقة بصحة الفم أثناء الحمل والمعلومات المتعلقة بصحة الفم لدى الرضع. كما جرى تقييم ممارساتهن المتعلقة بالمحافظة على صحة الفم (تنظيف الأسنان بالفرشاة أو بالخيط أو استخدام السواك) ومعدل زيارة أطباء الأسنان. لقد وُجد أن معظم النساء (>70%) يعرفن أن تسوس الأسنان عند الأطفال يمكن الوقاية منه، وأن الحمل يؤثر على صحة الفم، وأن معالجة الأسنان أثناء الحمل يمكن أن تؤثر سلباً على الرضع. ومعظم النساء (>80%) قمنَ باتباع عادات للمحافظة على صحة الفم، ولكن 18% منهن فقط قمنَ بزيارة طبيب الأسنان بانتظام. ولدى إجراء تحليل تحوُّفي تبين أن المعلومات المتعلقة بصحة الفم لم تكن مرتبطة ارتباطاً ملحوظاً بالممارسات المذكورة منهن لحفظ صحة الفم. وكانت النساء اللواتي زرن طبيب الأسنان بانتظام أكثر معرفة بكيفية الوقاية من التسوس لدى الأطفال، وبأن معالجة الأسنان أثناء الحمل كانت مرتبطة بصحة الرضع.

ABSTRACT This study assessed the association between oral health knowledge and practices of pregnant Saudi women selected from visitors to a government hospital in Dammam, Saudi Arabia in 2014. Women answered questions on oral health knowledge during pregnancy and knowledge of infant oral health. Most women (> 70%) knew that dental caries in children can be prevented, that pregnancy affects oral health and that dental treatment during pregnancy can negatively affect infants. Most women (> 80%) performed oral hygiene procedures but only 18% regularly visited the dentist. In a regression analysis, oral health knowledge was not significantly associated with reported oral hygiene practices. Women who visited the dentist regularly were more likely to know how to prevent caries in children, and that dental treatment during pregnancy and infant health were associated.

Association entre les connaissances et les pratiques de santé bucco-dentaire des femmes enceintes saoudiennes à Dammam (Arabie saoudite)

RÉSUMÉ La présente étude avait pour objectif d'étudier l'association entre les connaissances et les pratiques en matière de santé bucco-dentaire de femmes enceintes saoudiennes sélectionnées dans un hôpital public à Dammam, Arabie saoudite, en 2014. Les femmes interrogées ont répondu à des questions concernant leurs connaissances sur la santé bucco-dentaire pendant la grossesse et la santé bucco-dentaire du nourrisson. La plupart des femmes (plus de 70 %) avaient connaissance du fait que les caries dentaires chez l'enfant peuvent être évitées, que la grossesse a une influence sur la santé bucco-dentaire, et que les traitements dentaires pendant la grossesse peuvent avoir un impact négatif sur le nourrisson. La plupart des femmes (plus de 80 %) appliquaient des pratiques d'hygiène buccale, mais seulement 18 % se rendaient régulièrement chez un dentiste. Une analyse de régression a montré que les connaissances en matière de santé bucco-dentaire n'étaient pas associées de façon significative à l'application de pratiques d'hygiène buccale. Les femmes consultant régulièrement un dentiste étaient plus susceptibles de savoir comment prévenir les caries chez l'enfant, et qu'un traitement dentaire pendant la grossesse et la santé du nourrisson étaient corrélées.

¹ Department of Preventive Dental Sciences, College of Dentistry, University of Dammam, Dammam, Saudi Arabia (Correspondence to M. El Tantawi: maha_tantawy@hotmail.com).

Received: 21/06/15; accepted: 31/05/16

Introduction

Pregnancy is a critical period that involves physiological and psychological changes. These alterations may impose a burden on many women, leading them to neglect their general and oral health. There is a positive association between oral health problems such as periodontitis and adverse pregnancy outcomes such as preterm birth and eclampsia, and good oral health can reduce the risk of these outcomes (1–5). In the same context, several factors are known to be associated with children's oral health status, including maternal oral health knowledge and status, education, occupation and age (6–10).

Saudi Arabia is known for its high prevalence of caries (11,12). Children as young as 71 months have high rates of severe caries (13,14). Caries prevention programmes aim to control promoting factors (15) and reinforce protective factors (16). Oral health education programmes are cost-effective methods that can decrease disease risk through modifying social and dietary factors (17). These programmes aim to improve knowledge of oral health issues, and are based on the assumption that proper oral health practices occur when people have adequate knowledge (17). Many studies have investigated the applicability of this model among pregnant women in several countries with different cultures (18–20), and similar studies have been conducted among pregnant women in some regions of Saudi Arabia (21,22). However, there is inadequate information about whether knowledge of oral health issues leads to adoption of proper practices by Saudi pregnant women. The present study was conducted to: (1) assess oral health knowledge and reported oral health practices in pregnant Saudi women; and (2) investigate the association between these reported practices and oral health knowledge.

Methods

Study design and setting

A cross-sectional study was designed to collect data from pregnant women visiting a Ministry of Health hospital in Dammam, Saudi Arabia from February to April 2014. Approval for the study was obtained from the Research Ethics Committee of the College of Dentistry, University of Dammam (#EA2014003). The study was conducted in full accordance with the Helsinki Declaration. Responding to the questionnaire was considered an agreement to participate in the study.

Study sample

The estimated sample size was calculated to be 196 women (sample size calculator: <http://www.dartmouth.edu/~eugened/power-samplesize.php>). The following assumptions were used to estimate sample size: type I error = 5%; type II error = 20%; and odds that proper knowledge (of the relation between dental care during pregnancy and fetal health) is associated with seeking care during pregnancy = 2.5. Correct knowledge of this issue was assumed to be present in 50% of pregnant women, and actual regular seeking of care by visiting the dentist was assumed to be done by 20% of pregnant women.

Consecutive women were selected from those visiting the hospital between October and December 2014 who fulfilled the inclusion criteria that included: Saudi pregnant women; literate or accompanied by a literate companion to read and respond to the questionnaire; and free from systemic diseases as verified from the hospital records. This governmental hospital was selected for recruitment of study participants because of its large and well-equipped obstetrics and gynaecology department, to which pregnant women are referred from several cities in the Eastern Province, with an estimated 100 pregnant women visiting monthly for various reasons. Other hospitals were

not included because of their lower patient flow, which would have added only a small number to our sample. In addition, the diversity of patients from the selected hospital had a better chance of representing Saudi pregnant women in the Eastern Province than samples based on small hospitals with limited pools of patients.

Data collection tool

Data were collected using a self-administered, anonymous questionnaire in Arabic. The questionnaire was assessed for content and face validity and pilot tested on 20 women who were not part of the study. Previously published Arabic questionnaires could not be accessed because we could not reach their authors. Individual questions in the questionnaire were used and reliability was not assessed because there was no calculation of overall scores that expressed concept/status.

The study purpose was explained in an attached covering letter. The questionnaire included 20 close-ended questions that were separated into three sections. The first section was based on demographics and whether the women had children before the current pregnancy and the second section assessed oral health knowledge. In the second section, women answered true, false or do not know to each of 10 statements. They were also asked about their source of oral health information. The third and last section in the questionnaire assessed oral health practices. It included questions about: whether the participant performed any oral hygiene practices; frequency of brushing, flossing and using miswak; and if she had regular dental visits. The questionnaire was distributed to pregnant women attending the prenatal clinic for routine follow-up. Completed questionnaires were collected at the same session. Data were coded and entered into an Excel file.

Statistical analysis

Descriptive statistics were calculated to characterize the sample and assess the level of knowledge and oral health practices of the study sample. Univariate logistic regression models were developed for which the outcomes were: practicing oral hygiene (yes/no); and visiting the dentist regularly before and after pregnancy (yes/no). The independent variables were university education, having children, and knowledge related to each of the two outcomes. For oral hygiene practices, association with knowledge of the following was investigated: oral health relation with pregnancy; oral health relation with pregnancy outcomes; relation between maternal and infant oral health; transmission of cariogenic microorganisms to infant through saliva; and correct methods to prevent caries in children. For regularly visiting the dentist, the following were investigated as independent variables in addition to those listed above: could dental visits be scheduled during pregnancy; effect of dental treatment on infants; and actual practice of oral hygiene. Multivariate regression models were constructed to adjust mutually for all included variables. Odds ratios and confidence intervals were calculated.

Results

Two hundred and seventeen Saudi pregnant women were approached with the questionnaire and 197 (91%) responded. The greatest percentages of participants were in the age groups 21–30 and 31–40 years (47.4% and 42.2% respectively) and had less than university education (69.6%). Two thirds (59.1%) of them already had at least one child (Table 1).

The majority of participants knew that food affects oral health (88.7%), infants should not sleep while breast or bottle feeding (86.5%), oral health is affected by pregnancy (82.8%), caries in

Table 1 Characteristics of the sample of Saudi Arabian pregnant women included in the study (n = 197)

Variable	No. (%)
Age (years)	
≤ 20	12 (6.2)
21–30	91 (47.4)
31–40	81 (42.2)
> 41	8 (4.2)
Education	
Less than university	128 (69.6)
University and above	56 (30.4)
Has children	
Yes	114 (59.1)
No	79 (40.9)

children can be prevented (75.5%) and dental treatment does not negatively affect fetal health (72.9%). Around half the participants knew that dental visits can be scheduled during pregnancy (53.6%) and that pregnancy hormones

can affect oral health (44.7%). A minor proportion of participants knew that cariogenic bacteria can be transmitted from mother to child through saliva (28.9%), the correct methods to prevent caries in children (27.7%), and

Table 2 Responses related to self-reported oral health practices among Saudi pregnant women (n = 197)

Variable	No. (%)
Practicing oral hygiene	
Yes	161 (81.7)
No	36 (18.3)
Brushing	
Less than once daily (every other day or less frequently)	17 (8.8)
Once daily	77 (39.7)
Twice or more daily	100 (51.5)
Flossing	
Always	13 (6.8%)
Sometimes	84 (43.8)
Never	95 (49.5)
Miswak	
Always	17 (8.8)
Sometimes	92 (47.7)
Never	84 (43.5)
Dental visits	
Did not visit the dentist before or after pregnancy	66 (35.1)
Regularly visited the dentist before pregnancy but not after	62 (33)
Did not visit the dentist regularly before pregnancy but started after	26 (13.8)
Regularly visited the dentist for check-up before and after pregnancy	34 (18.1)

Table 3 Association between knowledge and practicing oral hygiene and regularly visiting the dentist (*n* = 197)

Variable	Practicing oral hygiene: OR (95% CI)		Regular visits to dentists: OR (95% CI)	
	Univariate regression	Multivariate regression	Univariate regression	Multivariate regression
University education				
Has children				
Knowledge				
Oral health affected by pregnancy	1.01 (0.44–2.29)	1.00 (0.41–2.34)	2.20 (0.85–5.70)	2.18 (0.83–5.71)
Oral health affects pregnancy	0.41 (0.20–0.88)*	0.32 (0.13–0.80)*	0.69 (0.32–1.46)	0.72 (0.27–1.56)
Oral health affects infants' oral health	0.72 (0.34–1.52)	0.69 (0.23–1.49)	0.67 (0.30–1.52)	0.65 (0.35–1.48)
Oral health affects infants' oral health	0.69 (0.26–1.79)	0.61 (0.24–1.70)	0.41 (0.14–1.26)	0.40 (0.11–1.22)
Oral health affects infants' oral health	0.34 (0.08–1.53)	0.30 (0.10–1.43)	0.84 (0.27–2.65)	0.85 (0.23–2.61)
Cariogenic microorganisms can be transmitted to children by saliva	0.85 (0.37–1.96)	0.89 (0.40–1.87)	0.74 (0.31–1.78)	0.72 (0.33–1.81)
Correct methods to prevent caries in children	2.33 (0.85–6.44)	2.48 (0.80–6.51)	2.97 (1.34–6.62)*	2.76 (1.07–7.12)*
Dentist can be visited during pregnancy	-	-	1.64 (0.76–3.57)	1.62 (0.74–3.58)
Dental treatment does not negatively affect fetal health	-	-	7.11 (1.63–31.05)*	5.14 (1.12–9.17)*
Practicing oral hygiene	-	-	2.60 (0.75–9.08)	2.59 (0.77–9.10)

*Statistically significant where CI does not include the null value
CI = confidence interval; OR = odds ratio

that maternal oral health may affect pregnancy outcomes (22.6%).

Most participants obtained oral health information from dentists or the media (36% and 30.6%, respectively), which together accounted for 66.6% of oral health information sources, followed by family and friends (12.9%) and several different sources (8.1%). Some patients (12.4%) reported no source of health information.

One hundred and sixty-one (81.7%) participants reported performing oral hygiene procedures, with 100 (51.5%) of them brushing at least twice daily, 84 (43.8%) sometimes flossing and 92 (47.7%) sometimes using miswak (Table 2). Only 34 (18.1%) reported regular dental visits before and after pregnancy.

Table 3 shows the association between oral health practices and knowledge in addition to two potential confounders: education and having children. Having children was associated with lower odds of practicing oral hygiene procedures, which remained statistically significant in the multivariate model that included knowledge and education [odds ratio (OR) in the multivariate model = 0.32]. This was the only factor in the model that was significantly associated with performing oral hygiene procedures. Knowledge of the correct methods to prevent caries in children was associated with higher odds of regular visits to the dentist (OR in multivariate regression = 2.76). Correct knowledge of the relation between dental treatment during pregnancy and fetal health was also associated with higher odds of regular dental visits (OR in multivariate model = 5.14). Practicing any form of oral hygiene was not associated with visiting the dentist (OR = 2.60, 95% confidence interval = 0.75–9.08).

Discussion

This study assessed the association between oral health knowledge and practices among Saudi pregnant women. The majority of the participants (>70%) revealed good oral health knowledge related to pregnancy and infancy. This disagrees with some previous studies (19,23) but agrees with many others (9,18,20,22). This reasonable level of knowledge may be related to the fact that one-third of women relied on dentists as a source of oral health information. Some participants had poor knowledge, which may be attributed to their educational level, as only one-third had a university degree. This agrees with Moawad et al., who found that women with lower education level had a lower score for oral health knowledge, particularly about the relation between periodontitis and preterm birth (22). Similarly, Chacko et al. (24) found that women with higher education expressed more awareness about bacterial

transmission from mother to child and about caries preventive methods, although this finding was not statistically significant. The present study also highlighted the role of dentists and mass media in providing oral health information. Chacko et al. (24) found that oral health information was delivered mainly through non-healthcare providers, and those who received information from healthcare workers had better oral health knowledge.

Half the participants reported brushing their teeth twice daily, although another half of them never flossed. These findings were similar to other studies that assessed oral health practices (18–20,22). Having children was negatively associated with performing oral hygiene procedures; most probably because mothers' time was taken up by tending to their children. Some studies have found a significant relation between regular dental visits and oral health knowledge; especially information about complications of bad oral health and bacterial transmission to infants (18,22). This is also similar to our findings. Dentists should therefore allocate time to explain the role of maternal oral health in preventing negative pregnancy outcomes, in addition to the maternal role in controlling transmission of cariogenic bacteria to infants. An emphasis on aspects related to child

health may be important in motivating pregnant women to care more for their own health. Preventive programmes using mass media may also be of value in providing correct oral health information, as well as programmes provided by nurses at antenatal clinics, while expectant mothers wait for check-ups.

The importance of oral health in ensuring positive pregnancy outcomes is acknowledged (1–5). Nevertheless, hormonal alterations affecting oral health during pregnancy cannot be prevented. As such, pregnant women need regular dental check-ups as part of their prenatal care. In the present study, only 18.1% visited the dentist before and after pregnancy and 33% stopped their dental visits during pregnancy. This agrees with previous studies that found that women may decrease or stop attending dental check-ups during pregnancy (18,19,22,25,26).

Health practices are assumed to be related to knowledge and a good level of knowledge is followed by adoption of proper practices (19). This was not the case in our study, which showed only a limited relation between oral health knowledge and practices. Our study also showed that adopting one oral health practice does not necessarily mean that others will be followed. We did not find an association between performing oral

hygiene and regular dental visits, which disagrees with Moawed et al. (22).

The main limitation of our study was relying on self-reports, in which the desire to give the correct response may have overestimated oral health practices. Future studies assessing oral hygiene clinically using plaque indices and assessing actual visits from records may avoid this bias. Our study explored some important aspects such as the extent of oral health knowledge among Saudi pregnant woman in and around Dammam and the relation between their oral health knowledge and practices.

In conclusion, Saudi pregnant women had basic oral health knowledge but this knowledge was not strongly associated with reported self-care through performing oral hygiene procedures. Similarly, pregnant women were more likely to visit the dentist during pregnancy if they had proper knowledge about the effect of oral health and dental treatment on children's oral health. Incorporating messages related to infant oral health in the information package delivered to pregnant women may further encourage Saudi pregnant women to visit the dentist regularly.

Funding: None.

Competing interests: None declared.

References

- Han YW. Oral health and adverse pregnancy outcomes – what's next? *J Dent Res*. 2011 Mar;90(3):289–93. PMID:21041548
- Sanz M, Kornman K; Working group 3 of joint EFP/AAP workshop. Periodontitis and adverse pregnancy outcomes: consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *J Clin Periodontol*. 2013 Apr;40 Suppl 14:S164–9. PMID:23627326
- Corbella S, Taschieri S, Francetti L, De Siena F, Del Fabbro M. Periodontal disease as a risk factor for adverse pregnancy outcomes: a systematic review and meta-analysis of case-control studies. *Odontology*. 2012 Jul;100(2):232–40. PMID:21739194
- Ide M, Papapanou PN. Epidemiology of association between maternal periodontal disease and adverse pregnancy outcomes – systematic review. *J Clin Periodontol*. 2013 Apr;40 Suppl 14:S181–94. PMID:23627328
- Kumar A, Basra M, Begum N, Rani V, Prasad S, Lamba AK, et al. Association of maternal periodontal health with adverse pregnancy outcome. *J Obstet Gynaecol Res*. 2013 Jan;39(1):40–5. PMID:22845916
- Bozorgmehr E, Hajizamani A, Malek Mohammadi T. Oral health behavior of parents as a predictor of oral health status of their children. *ISRN Dent* 2013 May 8; 2013:741783. PMID:23738088
- Dye BA, Vargas CM, Lee JJ, Magder L, Tinanoff N. Assessing the relationship between children's oral health status and that of their mothers. *J Am Dent Assoc*. 2011 Feb;142(2):173–83. PMID:21282684
- Weintraub JA, Prakash P, Shain SG, Laccabue M, Gansky SA. Mothers' caries increases odds of children's caries. *J Dent Res*. 2010 Sep;89(9):954–8. PMID:20505046
- Kishi M, Abe A, Kishi K, Ohara-Nemoto Y, Kimura S, Yonemitsu M. Relationship of quantitative salivary levels of *Streptococcus mutans* and *S. sobrinus* in mothers to caries status and colonization of mutans streptococci in plaque in their

- 2.5-year-old children. *Community Dent Oral Epidemiol.* 2009 Jun;37(3):241–9. PMID:19508271
10. Saied-Moallemi Z, Virtanen JI, Ghofranipour F, Murtomaa H. Influence of mothers' oral health knowledge and attitudes on their children's dental health. *Eur Arch Paediatr Dent.* 2008 Jun;9(2):79–83. PMID:18534175
 11. Al Agili DE. A systematic review of population-based dental caries studies among children in Saudi Arabia. *Saudi Dent J.* 2013 Jan;25(1):3–11. PMID:23960549
 12. Al-Ansari AA. Prevalence, severity, and secular trends of dental caries among various Saudi populations: A literature review. *Saudi J Med Med Sci.* 2014;2(3):142–50.
 13. Al-Malik MI, Holt RD, Bedi R. Prevalence and patterns of caries, rampant caries, and oral health in two- to five-year-old children in Saudi Arabia. *J Dent Child (Chic).* 2003 Sep-Dec;70(3):235–42. PMID:14998208
 14. Baghdadi ZD. Managing dental caries in children in Saudi Arabia. *Int Dent J.* 2011 Apr;61(2):101–8. PMID:21554279
 15. Daly B, Watt R, Batchelor P, Treasure E. *Essentials of dental public health.* Oxford: Oxford University Press; 2002.
 16. Pine C, Harris R, editors. *Community oral health*, 2nd ed. London: Quintessence; 2007.
 17. Watt RG. Strategies and approaches in oral disease prevention and health promotion. *Bull World Health Organ.* 2005 Sep;83(9):711–8. PMID:16211164
 18. George A, Johnson M, Blinkhorn A, Ajwani S, Bhole S, Yeo AE, et al. The oral health status, practices and knowledge of pregnant women in south-western Sydney. *Aust Dent J.* 2013 Mar;58(1):26–33. PMID:23441789
 19. Bamanikar S, Kee LK. Knowledge, attitude and practice of oral and dental healthcare in pregnant women. *Oman Med J.* 2013 Jul;28(4):288–91. PMID:23904926
 20. Abiola A, Olayinka A, Mathilda B, Ogunbiyi O, Modupe S, Olubunmi O. A survey of the oral health knowledge and practices of pregnant women in a Nigerian teaching hospital. *Afr J Reprod Health.* 2011 Dec;15(4):14–9. PMID:22571100
 21. Al-Zahrani AM, Al-Mushayt AS, Otaibi MF, Wyne AH. Knowledge and attitude of Saudi mothers towards their preschool children's oral health. *Pak J Med Sci.* 2014 Jul;30(4):720–4. PMID:25097504
 22. Moawed S, Hawsawi A, Al Ahmed SS, Al-Atawi N, Awadien AAZ. Knowledge and oral health care practices among Saudi pregnant women. *Life Sci J.* 2014;11(5):32–41 (http://www.lifesciencesite.com/lj/life1105/005_23165blife110514_32_41.pdf, accessed 2 June 2016)
 23. Jain R, Oswal KC, Chitguppi R. Knowledge, attitude and practices of mothers toward their children's oral health: A questionnaire survey among sub-population in Mumbai (India). *J Dent Res Sci Develop.* 2014;1(2):40–5.
 24. Chacko V, Shenoy R, Prasy HE, Agarwal S. Self-reported awareness of oral health and infant oral health among pregnant women in Mangalore, India – a prenatal survey. *Int J Health Rehab Sci.* 2013;2(2):109–15.
 25. Boggess KA, Urlaub DM, Massey KE, Moos MK, Matheson MB, Lorenz C. Oral hygiene practices and dental service utilization among pregnant women. *J Am Dent Assoc.* 2010 May;141(5):553–61. PMID:20436103
 26. Al-Swuailem AS, Al-Jamal FS, Helmi MF. Treatment perception and utilization of dental services during pregnancy among sampled women in Riyadh, Saudi Arabia. *Saudi J Dent Res.* 2014;5(2):123–9