

Drop-out from a drug treatment clinic and associated reasons

Leila Hoseinie¹, Zhaleh Gholami¹, Behrang Shadloo¹, Azarakhsh Mokri¹, Masoumeh Amin-Esmaeili¹ and Afarin Rahimi-Movaghar¹

معدلات التسرب من عيادات العلاج من الإدمان والأسباب المرتبطة بها

ليلي حسيني، جاله غلامي، بهرنج شادلو، آذرخش مكري، معصومة أمين-إسماعيلي، آفرين رحيمي-موقر

الخلاصة: تمثّل الهدف من هذه الدراسة في تقييم معدلات التسرب من العلاج والأسباب المرتبطة بها في عيادة المركز الوطني الإيراني لدراسات الإدمان. وفي فترة سنة واحدة (أبريل/ نيسان 2014 حتى مارس/ آذار 2015)، أدرج في هذه الدراسة جميع المرضى الذين يعانون من إدمان المخدرات ممن أحيلوا للعلاج وخضعوا لتقييم أولي (حجم العينة = 242). وتمت متابعة الأشخاص الذين تلقوا العلاج حتى مارس/ آذار 2016. وأظهر تحليل القدرة على البقاء على قيد الحياة تسرب 70.2٪ من العلاج. وأظهر اختبار رتبة اللوغريتمية أن معدلات التسرب من العلاج تختلف باختلاف الأساليب المستخدمة ($P < 0.001$)، وهو أدنى ميل في العلاج المستمر بالإنبيورينورفين وأعلى ميل في برنامج إزالة السُمّية. وبلغت معدلات التسرب خلال الأشهر الثلاثة الأولى 62٪ (الانحراف المعياري = 0.05) و 82.4٪ (الانحراف المعياري = 0.03) لإدمان الأفيونيات والمنبهات، على التوالي. وأجريت تحليلات باستخدام برامج SPSS (الإصدار 21.0)، و STATA الإحصائية (الإصدار 13.0). ومن منظور المرضى، اعتُبرت أوجه التناقض في الحوافز السبب الرئيسي لعدم البدء في العلاج أو تركه. ويمكن أن توفر نتائج هذه الدراسة فهماً أفضل لمقدمي الخدمات بشأن معدلات التسرب وما يرتبط بها من أسباب.

ABSTRACT The aim of this study was to assess drop-out rates and associated reasons among patients at the Iranian National Center for Addiction Studies (INCAS) clinic. In a one-year period (April 2014 to March 2015), all patients with drug dependence who had been referred for treatment and attended for a first assessment were included in this study (N=242). Those who received treatment were followed until March 2016. Survival analysis showed that 70.2% had dropped out from treatment. Log rank test showed that treatment drop-out rates differed between the different approaches used ($P < 0.001$), with the lowest slope in buprenorphine maintenance treatment and the highest in the detoxification programme. Drop-out rates within the first three months was 62% (SE= 0.05) and 82.4% (SE=0.03) for opioids and stimulants dependence, respectively. Analyses were performed using SPSS (Version 21.0) and STATA software, (version 13.0). From the patients' perspective, motivational inconsistencies were considered as the main reason for not starting or leaving treatment. The findings of this study could give service providers a better grasp of drop-out rates and the associated reasons.

Taux d'abandon dans un centre de traitement de la dépendance aux drogues et raisons associées

RÉSUMÉ La présente étude avait pour objectif d'évaluer les taux d'abandon et les raisons associées parmi les patients de la clinique du Centre national iranien d'études sur les addictions (INCAS). Sur une période d'un an (avril 2014 à mars 2015), tous les patients souffrant de dépendance aux drogues ayant été transférés en vue d'un traitement et consultant pour la première fois dans cette structure ont été inclus dans l'étude (n=242). Les patients sous traitement ont été suivis jusqu'en mars 2016. L'analyse de survie a montré que 70,2 % des patients avaient abandonné le traitement. Le test du log-rank a révélé que les taux d'abandon de traitement différaient selon les approches utilisées ($p < 0,001$), la baisse la moins importante concernant le traitement d'entretien à la buprénorphine et la baisse la plus forte les programmes de désintoxication. Les taux d'abandon durant les trois premiers mois étaient de 62 % (erreur-type = 0,05) et de 82,4 % (erreur-type = 0,03) pour la dépendance aux opioïdes et aux stimulants respectivement. Les analyses ont été réalisées à l'aide des logiciels SPSS (version 21.0) et STATA (version 13.0). Selon les patients, le manque de motivation constituait l'une des raisons principales pour ne pas commencer ou abandonner un traitement. Les résultats de cette étude pourraient offrir aux prestataires de service une meilleure compréhension des taux d'abandon et des raisons associées.

¹Iranian National Center for Addiction Studies (INCAS), Iranian Institute for Reduction of High-Risk Behaviour, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran. (Correspondence to: Masoumeh Amin-Esmaeili: dr.m.a.esmaeili@gmail.com)

Received: 31/05/16; accepted: 01/03/17

Introduction

Substance use disorder is a chronic relapsing condition which requires long-term treatment. In the Islamic Republic of Iran, it is estimated that > 1 million people are suffering from addiction to an illicit drug (1). During the past 2 decades, drug treatment services have expanded dramatically, both in service coverage and diversity of services. There are now > 5000 outpatient drug treatment clinics in the country. However, only 21% of people with drug use disorder have received services from these clinics in the past 12 months (1).

Addiction treatment is a complex process and achieving desirable outcomes relies on several factors. Retention in the treatment and completion of its course are some of the most prominent factors. Completing the course of treatment is associated with higher abstinence, lower crime rate, lower relapse and higher employment compared to those who have dropped out (2). At the same time, there are studies indicating that drop-out might result in wasting resources of the individual, the healthcare system and society. There have been several studies on treatment retention, drop-out and associated factors; however, due to differences in the types of treatment, treatment setting and patients' characteristics, different rates of drop-out, from 0.4% to 90%, have been reported (3).

Different reasons have been given for drop-out. Lack of motivation, fear of treatment failure and interpersonal conflicts have been reported as the reasons for treatment drop-out (4–6). Therefore, with proper detection and removal of such obstacles, one may be able better to provide the required services and fulfil treatment protocols and decrease the cost to the healthcare system.

Our knowledge regarding drop-out usually comes from the results of controlled trials in experimental settings; however, the results might not

be applicable to the real world. The clinic of the Iranian National Center for Addiction Studies (INCAS) could be considered as a prototype of a specialized drug treatment clinic and observations of treatment course and outcomes from this centre can be applied to other similar facilities. The INCAS clinic, with more than a decade of experience in the field of delivering drug treatment services, is an academic centre affiliated to the Tehran University of Medical Sciences. The clinic is located at the heart of the city of Tehran and receives a variety of clients from all over Tehran.

For opioid dependence, the INCAS clinic offers an assortment of services including methadone and buprenorphine maintenance treatment, short-term detox programmes with counselling, and other psychosocial rehabilitation approaches. The provided treatment modality for each patient is based on patient characteristics, severity of dependence, previous treatment attempts and patient preference. Maintenance treatments could continue for several years but usually buprenorphine maintenance treatment (BMT) is provided for 1 year and detox programmes are followed by 6-months relapse-prevention counselling. For amphetamine-type stimulants dependence, the centre offers a treatment package, called the Matrix model, which is a structured 24-session individual psychotherapy with a cognitive-behavioural approach along with family education. The therapy sessions are provided 2 or 3 times a week. These sessions cover areas such as: trigger and craving management techniques; basic required skills for remission (e.g., managing boredom and other negative emotions, building trust, stress and anger management); and relapse-prevention counselling. For cannabis dependence, similar core cognitive-behavioural techniques, as in the Matrix model, are provided for the patients. Group family education sessions are also provided. Patients are encouraged to attend group therapy,

focusing on relapse prevention, after the completion of the structured therapy sessions. After 1 session of initial assessment and development of a treatment plan, the second appointment for all modalities is made within the next 7 days.

This study was carried out in a group of treatment-seeking clients of the INCAS clinic. The aims of this study were: (1) to assess the rates of drop-out in the total sample and in different treatment categories; and (2) to assess the reasons for drop-out in those who had left the treatment and provide pragmatic recommendations in order to achieve higher treatment retention rates.

Methods

We included 242 clients from April 2014 to end of March 2015 who had been referred to the INCAS clinic for treatment of opioids, amphetamine-type stimulants, prescription drugs and cannabis dependence. The INCAS clinic benefits from an electronic registry system, called Addiction Treatment Centers Automation Software (ATCAS). ATCAS was used to obtain the clients' sociodemographics, contact information, and data regarding the pattern of substance abuse and the treatment approach at each entry point. The study was conducted in 2 phases. First, drop-out was assessed using the ATCAS registry up to the end of March 2016 and survival analysis was carried out. Second, the reasons for drop-out were gathered via telephone interviews using a questionnaire.

Drop-out was defined either as not attending at all after the first assessment or otherwise according to the specific treatment modality started. For opioid substitution treatment, a ≥ 7 -day leave of treatment, and for psychotherapy and detox and relapse-prevention programmes, failure to show up for the next therapy session, even after setting

another appointment were considered as drop-out.

The interviewer contacted all individuals who dropped out (either those who did not start treatment or those who had left after treatment initiation) from April to September 2015. The interviewer was a trained nurse making telephone calls with the use of ATCAS registry. For ethical and professional considerations, interviews were conducted after confirmation of the identity of the respondent. This was achieved by asking for national identity and birth certificate numbers and other identifying data. After confirming the identity of the client, the interviewers introduced themselves and gave the reasons for calling. After obtaining oral consent and stating the participant's right of refusal, the interview began. First, an open-ended question regarding the client's overall opinion toward treatment obstacles was asked. Then the interview continued by using a structured questionnaire.

Instruments

A modified version of Reason for Leaving Treatment Questionnaire (RLTQ) was implemented (6). RLTQ consists of 28 true/false questions and covers 7 domains: motivational inconsistencies (4 questions), staff conflicts (3 questions), boundary concerns (5 questions), outside influences (4 questions), programme expectations (4 questions), problem severity (4 questions) and logistic problems (4 questions). Respondents were asked to determine if the statement related to their decision not to start or to leave the treatment. A domain was considered positive if one of its related questions was true. In order to implement RLTQ, the questionnaire was translated and modified according to the cultural context of the addiction treatment services in the country. Two other questions for boundary concerns and 2 questions to assess pharmacological adverse effects for those receiving maintenance treatments were added to the questionnaire. Therefore, the total

number of questions for those on maintenance programmes and nonpharmacological approaches were 32 and 30, respectively. Content validity of the questionnaire was assessed by an expert panel of drug treatment specialists.

Ethical considerations

The research protocol was approved by the Ethics Committee of Tehran University of Medical Sciences in Iran (No. IR.TUMS.REC.1394.2136). Participation in the study was on a voluntary basis, and oral consent was obtained before all interviews. All questionnaires and the database of the study were kept confidential by the research team.

Data analysis

SPSS Statistics for Windows version 21.0 (IBM Corp., Armonk, NY, USA) and Stata Statistical Software version 13 (Stata Corp., College Station, TX, USA) were used for data analysis. For quantitative variables (such as age), *t* test and analysis of variance were used. For qualitative variables (such as gender, marital status, type of treatment and substance of abuse) χ^2 test was applied. Survival analysis (Kaplan–Meier: log rank test and Cox regression) was used to assess days staying in treatment. For those who started treatment, the period of survival of each case was calculated between the date of admission to the clinic and the date of leaving treatment or completion of treatment. For the case of methadone maintenance treatment (MMT), survival was calculated until the end of the study period in March 2016. The survival data were considered to be subject to right censoring. *P* values < 0.05 were considered statistically significant in all tests.

Results

Between April 1, 2014 and March 31, 2015, 242 individuals with substance use disorders (226 men and 16 women) were referred to the INCAS clinic

for treatment and attended for initial assessment. A total 16 689 person–days of services were delivered. Following the initial assessment, treatment need was identified as solely for stimulants (56.2%), solely for opioids (32.6%), both for stimulants and opioids (7.4%) or solely for cannabis or prescription drugs (3.8%). Among the clients, 61 (25.2%) visited only once and did not return again, although the initial assessment was done and the treatment approach was determined and recommended. Treatment initiation failure was 26.4%, 26.9%, 21.0% and 8.3% for psychotherapy, MMT, BMT and detox programme, respectively. Moreover, treatment initiation failure was 28.7%, 25.3% and 11.1% for stimulant, opioid and combination of opioid and stimulant dependence, respectively and zero for cannabis and prescription drugs.

In addition to the 61 clients who, following the first assessment, did not take up further treatment offered at the centre, from the remaining 181 clients who did engage further, 127 (70.2%) subsequently dropped out. Drop-out rate after treatment initiation was 74.3% for psychotherapy and 63.0–66.7% for the other 3 medically assisted treatment modalities. The overall incidence rate of drop-out from the treatment was 7.6 per 1000 person–days of service [95% confidence interval (CI): 6.4–9.1]. Twenty-five percent had left the treatment in the first 12 days and 50% in the first 42 days. Drop-out during the first 3 months was seen in 62.0% [standard error (SE) 0.05] of opioid-dependents and 82.4% (0.03) of stimulant-dependents ($P < 0.001$).

Comparison of those who had dropped out with those remaining on treatment revealed that age, gender, marital status, employment, substance of abuse and treatment modality were not significantly correlated with drop-out (Table 1).

Rates of drop-out from different treatment approaches (detox programmes, psychotherapy, BMT and

Table 1 Characteristics of participants who either remained in treatment or left/not started treatment

Patient characteristics	All patients (n = 242)	Patients who stayed in treatment (n = 54)	Patients who did not start or left treatment (n = 188)	P
Age, yr, mean (SD)	36.1 (9.4)	36.9 (10.6)	35.9 (9.1)	0.49
Min–Max	15–74	18–63	15–74	
Sex, male (%)	226 (93.4)	49 (90.7)	177 (94.2)	0.37
Marital status (%)				0.86
Married	137 (57.3)	29 (54.7)	108 (58.1)	
Never married	79 (33.1)	18 (34.0)	61 (32.8)	
Previously married	23 (9.6)	6 (11.3)	17 (9.1)	
Educational status (%)				0.02*
Illiterate or elementary school	21 (9.2)	6 (11.3)	15 (8.6)	
Middle school	47 (20.6)	6 (11.3)	41 (23.4)	
High school	104 (45.6)	33 (62.3)	71 (40.6)	
Higher education	56 (24.6)	8 (15.1)	48 (27.4)	
Occupation (%)				0.65
Employed	147 (61.5)	34 (64.2)	113 (60.8)	
Unemployed or not in labour force	92 (38.5)	19 (35.8)	73 (39.2)	
Reason for referral (%)				0.39
Stimulants	136 (56.2)	24 (44.4)	112 (59.6)	
Opioids	79 (32.6)	23 (42.6)	56 (29.8)	
Opioids+ stimulants	18 (7.4)	5 (9.2)	13 (6.9)	
Prescription drugs	5 (2.1)	1 (1.9)	4 (2.1)	
Cannabis	4 (1.7)	1 (1.9)	3 (1.6)	
Treatment (%)				0.43
Methadone maintenance treatment	63 (26.0)	17 (31.5)	46 (24.5)	
Buprenorphine maintenance treatment	19 (7.8)	5 (9.3)	14 (7.4)	
Detoxification + relapse prevention program	12 (5.1)	4 (7.4)	8 (4.3)	
Psychotherapy	148 (61.2)	28 (51.8)	120 (63.8)	

*Level of significance < 0.05.

MMT) are demonstrated in a Kaplan–Meier curve (Figure 1). Log rank test showed that treatment drop-out differed significantly for different approaches ($P < 0.001$), with the lowest slope for BMT and highest for detox. Cox regression analysis showed that detoxification resulted in a significant decrease (hazard ratio: 3.27, 95% CI: 1.10–9.78) in remaining on treatment, after adjustment for education and type of substance.

To assess the reasons for not starting or leaving treatment, a minimum

of 3 and a maximum of 7 telephone calls were made to the 188 individuals who either did not start treatment or had dropped out after initiation. Seventy people (37.2%) could not be reached due to several possible reasons: wrong number in the registry, change of phone number, disconnected line or non-response to the call. Contact was made with 118 individuals, from which 40 (33.9%) refused to participate in the study. In addition, 2 were incarcerated and 2 were in a residential drug treatment facility. Another 2 had passed away, both opioid dependent, 1 due

to accidental electrocution and 1 had committed suicide. The questionnaire was filled out for 76 individuals (Figure 2). Those who could not be reached or refused to participate did not significantly differ in age ($P = 0.66$), gender ($P = 0.29$), type of substance ($P = 0.29$) and treatment modality ($P = 0.42$) from those who completed the telephone interview.

While answering the initial open-ended question regarding the reason of not starting or leaving treatment, the following answers were found to be




Figure 1 Rates of drop-out demonstrated in a Kaplan–Meier curve for staying in treatment (n = 181). BMT = buprenorphine maintenance treatment; MMT = methadone maintenance treatment

more common: commuting difficulties, dissatisfaction with the services, cost of treatment and finally lack of motivation. Some indicated adverse effects of methadone; another group had difficulties matching themselves with the working hours of the centre, had conflicts in the family or were concerned about treatment failure. Few indicated medical conditions or stigma regarding attending a drug clinic.

Results of the interview with RLTQ (Table 2) showed that outside influences (86.3%) followed by motivational inconsistencies (65.8%) were the most common reasons for not starting or leaving treatment. Staff conflicts were reported by 27.6% of respondents. In the outside influences domain, 61.6% believed that they could get better on their own or through self-help groups. In the motivational inconsistencies domain, 50.0% mentioned that they had

changed their minds about being in treatment at that moment.

Apart from the outside influences domain ($P = 0.01$), others were not significantly correlated with treatment modality. Three clients from detox programmes, 43 from psychotherapy services, 13 in MMT and 4 in BMT indicated outside influences for not starting or leaving treatment. Motivational inconsistency did not have a significant association with gender (P




Figure 2 Flow diagram indicating participants' responses.

Table 2 Participants' reasons for not starting or leaving treatment

Item	<i>n</i>	%
Motivational inconsistencies (<i>n</i> = 76)	50*	65.8*
I changed my mind about being in the treatment at this point	38	50.0
I had no good reason to stop using drugs	21	27.6
I did not feel motivated enough to keep coming	25	32.9
I lost hope in my ability to change right now	27	35.5
Staff conflicts (<i>n</i> = 76)	21*	27.6*
I had a negative interaction with another staff member	11	14.5
I did not like or trust some of the staff	15	19.7
I felt that staff did not like, respect or want to help me	15	20.0
Boundary concerns (<i>n</i> = 75)	34*	45.3*
I had a personality conflict with people at the programme	19	25.3
I was worried that the therapist would talk about my situation to my family	11	14.7
I was worried that the therapist would give my personal information to the police	6	8.0
I felt my privacy or confidentiality might not be respected	12	16.0
Somebody I know is a client or staff member in the programme	7	9.3
I said or did some things that would make it hard for me to go back	9	12.0
I was worried I would get involved in the wrong things like drugs, sex or crime because of people around the programme	8	10.7
Outside influences (<i>n</i> = 73)	63*	86.3*
Problems with family or acquaintances kept me from coming in	16	21.9
I felt that I could get better on my own or with self-help meetings	45	61.6
I did not have enough support from people in my life to stay in the programme	21	28.8
I decided to go to another programme for help	32	43.8
Program expectations (<i>n</i> = 73)	42*	57.5*
I did not like the rules the programme had	24	32.9
I was confused about what the programme wanted me to do	13	17.8
I did not like the kind of services offered at the programme	20	27.4
The wait to start the programme was too long	25	34.3
Problem severity (<i>n</i> = 73)	34*	46.6*
My medical problems kept me from coming	7	9.6
My drug use was so heavy I could not come in	17	23.3
My alcohol use was so heavy I could not come in	2	2.7
My mental health or psychological problems kept me from coming	27	37.0
Logistical problems (<i>n</i> = 73)	42*	57.5*
I had transportation problems that kept me from coming	26	35.6
I had childcare problems that kept me from coming in	1	1.4
The hours of the programme were not good for me	24	32.9
I did not have money or insurance to pay for the programme	22	30.1
Drug treatment problem (<i>n</i> = 12)	7*	58.3*
I thought that I would become dependent to methadone or buprenorphine	6	50.0
I thought that I would experience side effects of methadone or buprenorphine	3	25.0

*Any item endorsed.

= 0.20), marital status ($P = 0.29$), level of education ($P = 0.56$), employment ($P = 0.83$) and the type of substance ($P = 0.28$). Clients aged ≤ 30 years mentioned motivational inconsistencies

significantly more than those aged > 30 years ($P = 0.03$). Among those who completed the questionnaire, 3 were women. All 3 mentioned motivational inconsistency as the main reason and

none reported conflicts with staff or severity of the problem.

Service-related factors constituted a lower, but still considerable proportion of the reasons given for not starting

or leaving treatment. Long waiting list before the initiation of the treatment package (34.3%), service delivery hours (32.9%), having difficulties with following the regulations of the centre (32.9%), cost of services and lack of insurance coverage (30.1%), disinterest in the proposed treatment modality (27.4%) and ambiguities regarding expectations from treatment outcomes (17.8%) were the mentioned reasons in this regard.

A higher number of those who had not started treatment following their initial assessment reported motivational factors, staff conflicts and logistic problems as their main reasons for drop-out. A significant number of these clients mentioned that they decided to go to another programme after their first visit to this centre.

Discussion

This study showed that more than two thirds of the clients had dropped out of treatment, even after excluding those for whom the treatment had not been initiated. Other studies have also indicated high drop-out rates. In a systematic review, Brorson et al. indicated that in more than one-third of the 122 included studies, attrition rates of 50–90% were reported (3). Drop-out differed according to the definition and the treatment setting and the time span of evaluation. A systematic review of randomized clinical trials showed that 47.9% on BMT and 37.7% on MMT had dropped out within 2–52 weeks (7). The studies included in the mentioned systematic review benefitted from a structured and highly supervised treatment plan. However, the effectiveness is not usually the same in the real world. For instance, a study reported 67.6% drop-out within 18 months in MMT clinics (8). In another study half of the patients dropped out within 12 months (9). Another study showed that the rate of 12-week opioid abstinence was 16% in BMT

and 8% in those on oral naltrexone (10). The result of a national household study in the Islamic Republic of Iran indicated that nearly 40% of those receiving an outpatient health service for addiction have dropped out of treatment (1).

Several factors influence treatment retention. Treatment modality has been proposed as one of the most important factors. In the current study, treatment drop-out happened earlier in the detox programme and later in BMT. Opioid substitution treatment programmes are associated with lower mortality, crime, injecting drug use, HIV acquisition, and also benefit from higher retention rates (11). Although, maintenance treatments are associated with more desirable outcomes, the cost of the treatment and the associated financial burden could influence the individual's choice (12,13). When comparing MMT with BMT, MMT was associated with lower cost and higher effectiveness and higher retention rates in randomized trials (7,14). Better retention rates of BMT in our study might have been due to the selection of patients. Those with less severe opioid dependence are preferably offered BMT. Although all 3 approaches of detox, MMT and BMT are provided in this centre and most of the clients were married, literate and employed, MMT was preferred. Client preference has an important role in treatment choice and could be the reason for higher rates of methadone implementation at the centre.

The type of substance can also influence treatment retention. In the current study, drop-out in the first 3 months was significantly higher in those with stimulant compared to opioid dependence. The review by Brorson et al. indicated that in most of the studies, type of substance was not correlated with treatment retention (3). There have been a few studies on retention rates for stimulant dependence (15–17). A study on the trend of drop-out in this group indicated 53–64% rates of drop-out (15). Another study indicated that

treatment attrition was not significantly different in heroin versus amphetamine users. For decades, opium has been the main substance of abuse in the Islamic Republic of Iran, while methamphetamine is a new phenomenon. The results from 3 repeated national studies on drug use show that methamphetamine appeared in the mid-2000s (18–20). Since amphetamine-type stimulant treatment has just recently started in the country, it is necessary to enhance staff training and supervision to improve the quality of such services and to try to optimize retention rates.

In the current study, 25.2% of the clients attended only a single visit at the clinic, and although a thorough assessment was done and the treatment plan was agreed, the client did not return for initiation of the programme. Other studies have also reported significant attrition after the first visit. For example, drop-out rates of 15% (21), 27% (22) and 25.9–41% (23) have also been reported after the initial assessment. Therefore, the first contact is crucial. Building a proper rapport, providing useful information and raising awareness regarding treatment outcomes could help in retaining the individual in the treatment process. It is necessary to assess the motivational aspects of the client and utilize proper techniques to enhance them during the first contact.

Regarding the reasons for drop-out, factors associated with service delivery although reported by a minority of respondents, are also of importance. Other studies have indicated that these factors constitute a minority of the reasons for leaving treatment (6,24,25). However, adding reasons related to boundary concerns, programme expectations and logistical problems results in a significant contribution of clinic-related factors. Removing such obstacles could promote the quality of the provided services. The clinics should closely monitor organizational factors that might negatively affect patient adherence. Developing quality assurance

and improvement measures, providing social support packages, flexible timing, insurance and commuting facilitation could have desirable effects on the treatment process and thus lower drop-out.

Another finding of this study was that motivational inconsistencies were one of the main reasons for not starting or leaving treatment. Most of the respondents stated that they had changed their mind about being on treatment and some felt they were not ready for a change at the moment. Lack of motivation has been reported as a reason for drop-out in several different studies (3,4,6,25,26). Addiction treatment is a long-lasting process and requires several changes in the individual's lifestyle and abstaining from pleasure-seeking behaviour. It is obvious that the individual's motivation will change during this process. Therefore, regular monitoring and use of motivational enhancement techniques and teaching adaptive coping skills could increase treatment retention.

Outside influences were also frequently mentioned as reasons for not starting or leaving treatment. Most of the respondents believed that they could use other methods for recovery, either on their own or through self-help groups. It is possible that particularly for some of those initially engaging with and requiring psychotherapy, alternative available supports may be associated with a capacity for self-recovery outside of treatment and so a person may reasonably feel that he or she does not need to attend specific treatment

services. This points to a potential positive reflection on the possible value of practitioners and services providing good information and visibility of the additional or alternative external supports available, including from the initial assessment onwards.

Another group mentioned conflicts with the family members as an obstacle for treatment. External factors are a part of the patient's real world, but could serve for denial or as an excuse to procrastinate the treatment process. Assessing the clients' beliefs in this regard and their support from significant others could be used to tackle this obstacle.

Demographics did not correlate with drop-out in our study. The only exception was that younger individuals had lower levels of motivation than older ones. In other studies, low motivation has been found to be more prevalent in clients who are single than those who are married (6). Considering the findings of the current study, it seems that use of motivational techniques and obtaining the support of the family, especially for those who are younger, could help in achieving better results.

The study had some limitations. The sample size was small in some subgroups: detoxification as a treatment modality, young age group and female gender. Moreover, building the trust of the interviewees through telephone calls was challenging. Although several measures were adopted to facilitate engagement in the process and to obtain consent, a significant number did not either answer or refused to participate in

the study. Recall bias was another limitation of the study, and in some cases it had been a while since the patients had dropped out of treatment and had difficulty to recall the reason why they left.

Conclusions

The results of the current study indicated that the motivational inconsistencies had the highest impact on not starting or leaving treatment, while several other influential, and potentially remediable, service-related factors were also identified. The findings of this study could be of use for service providers and policy-makers to adopt proper measures to increase treatment retention. It is also recommended that for enhancing the motivation of the clients, novel and more client-centred approaches with proper involvement of family members be adopted.

Acknowledgements

The authors would like to express their gratitude to Ms. Mehri Noori, Ms. Rahimeh Azimi and Ms. Mahboubeh Abbasi for their contribution in this study. We wish to thank all participants in this study.

Funding: This study was financially supported by Tehran University of Medical Sciences through the contract number 132-1114.

Competing interests: None declared

References

1. Amin-Esmaeili M, Rahimi-Movaghar A, Sharifi V, Hajebi A, Radgoodarzi R, Mojtabai R, et al. Epidemiology of illicit drug use disorders in Iran: prevalence, correlates, comorbidity and service utilization results from the Iranian Mental Health Survey. *Addiction*. 2016 Oct;111(10):1836-47. PMID:27177849
2. Kleber HD, Weiss RD, Anton RF Jr, George TP, Greenfield SF, Kosten TR, et al. Treatment of patients with substance use disorders, second edition. American Psychiatric Association. *Am J Psychiatry*, 2007 Apr;164(4 Suppl):5-123. PMID:17569411
3. Brorson HH, Ajo Arnevik E, Rand-Hendriksen K, Duckert F. Drop-out from addiction treatment: a systematic review of risk factors. *Clin Psychol Rev*. 2013 Dec;33(8):1010-24. PMID:24029221
4. Palmer RS, Murphy MK, Piselli A, Ball SA. Substance user treatment dropout from client and clinician perspectives: a pilot study. *Subst Use Misuse*. 2009;44(7):1021-38. PMID:19938942
5. Stark MJ. Dropping out of substance abuse treatment: a clinically oriented review. *Clin Psychol Rev*. 1992;12(1):93-116.

6. Ball SA, Carroll KM, Canning-Ball M, Rounsaville BJ. Reasons for dropout from drug abuse treatment: Symptoms, personality, and motivation. *Addict Behav.* 2006 Feb;31(2):320-30. PMID:15964152
7. Mattick RP, Breen C, Kimber J, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Database Syst Rev.* 2004;(3):CD002207. PMID:15266465
8. Lin HC, Chen KY, Wang PW, Yen CF, Wu HC, Yen CN, et al. Predictors for dropping-out from methadone maintenance therapy programs among heroin users in southern Taiwan. *Subst Use Misuse.* 2013 Jan;48(1-2):181-91. PMID:23368704
9. Smyth BP, Fagan J, Kernan K. Outcome of heroin-dependent adolescents presenting for opiate substitution treatment. *J Subst Abuse Treat.* 2012 Jan;42(1):35-44. PMID:21940134
10. Mokri A, Chawarski MC, Taherinakhost H, Schottenfeld RS. Medical treatments for opioid use disorder in Iran: a randomized, double-blind placebo-controlled comparison of buprenorphine/naloxone and naltrexone maintenance treatment. *Addiction.* 2016 May;111(5):874-82. PMID:26639678
11. Connock M, Juarez-Garcia A, Jowett S, Frew E, Liu Z, Taylor RJ, et al. Methadone and buprenorphine for the management of opioid dependence: a systematic review and economic evaluation. *Health Technol Assess.* 2007 Mar;11(9):1-171. PMID:17313907
12. Stotts AL, Dodrill CL, Kosten TR. Opioid dependence treatment: options in pharmacotherapy. *Expert Opin Pharmacother.* 2009 Aug;10(11):1727-40. PMID:19538000
13. Kleber HD. Pharmacologic treatments for opioid dependence: detoxification and maintenance options. *Dialogues Clin Neurosci.* 2007;9(4):455-70. PMID:18286804
14. King JB, Sainski-Nguyen AM, Bellows BK. Office-based buprenorphine versus clinic-based methadone: a cost-effectiveness analysis. *J Pain Palliat Care Pharmacother.* 2016;30(1):55-65. PMID:27007583
15. Brecht ML, Greenwell L, Anglin MD. Methamphetamine treatment: trends and predictors of retention and completion in a large state treatment system (1992-2002). *J Subst Abuse Treat.* 2005;29(4):295-306. PMID:16311182
16. Gonzales R, Ang A, McCann MJ, Rawson RA. An emerging problem: methamphetamine abuse among treatment seeking youth. *Subst Abuse.* 2008;29(2):71-80. PMID:19042326
17. Maglione M, Chao B, Anglin MD. Correlates of outpatient drug treatment drop-out among methamphetamine users. *J Psychoactive Drugs.* 2000 Apr-Jun;32(2):221-8.
18. Razaghi EM, Rahimi A, Hosseini M, Madani S, Chatterjee A. Rapid situation assessment of drug abuse in Iran (<http://www.unodc.org/pdf/iran/publications/RSA2000SUMMARY.pdf>, accessed 14 March 2017).
19. Narenjiha H, Rafiei H, Baghestani AR, Nouri R, Shirinbayan P, Farhadi MH, et al. Rapid situation assessment of drug abuse in Iran (year 2004): research report. Tehran: Dariush Institute; 2005 (in Persian).
20. Narenjiha H, Rafiei H, Baghestani AR, Nouri R, Ghafouri B, Soleimaninia L, et al., editors. Rapid situation assessment of drug abuse in Iran (year 2007) Tehran: Dariush Institute; 2009 (in Persian).
21. Claus RE, Kindleberger LR. Engaging substance abusers after centralized assessment: predictors of treatment entry and dropout. *J Psychoactive Drugs.* 2002 Jan-Mar;34(1):25-31. PMID:12003110
22. Rohrer JE, Vaughan MS, Cadoret RJ, Carswell C, Patterson A, Zwick J. Effect of centralized intake on outcomes of substance abuse treatment. *Psychiatr Serv.* 1996 Nov;47(11):1233-8. PMID:8916242
23. Li X, Sun H, Marsh DC, Anis AH. Factors associated with pre-treatment and treatment dropouts: comparisons between Aboriginal and non-Aboriginal clients admitted to medical withdrawal management. *Harm Reduct J.* 2013 Dec;10:38. PMID:24325629
24. Laudet AB, Stanick V, Sands B. What could the program have done differently? A qualitative examination of reasons for leaving outpatient treatment. *J Subst Abuse Treat.* 2009 Sep;37(2):182-90. PMID:19339133
25. Evans E, Li L, Hser Y-I. Client and program factors associated with dropout from court mandated drug treatment. *Eval Program Plann.* 2009 Aug;32(3):204-12. PMID:19150133
26. Longshore D, Teruya C. Treatment motivation in drug users: a theory-based analysis. *Drug Alcohol Depend.* 2006 Feb;81(2):179-88. PMID:16051447