

Nutritional risk screening of hospitalized children aged < 3 years

Sanaa Shaaban,¹ May Nassar,¹ Yasmin El-Gendy¹ and Bassam El-Shaer¹

¹Faculty of Medicine, Ain Shams University, Cairo, Egypt (Correspondence to: M. Nassar: maie_nassar@yahoo.co.uk).

Abstract

Background: Identification of children at risk of malnutrition is not easily achieved in hospital settings.

Aims: To assess the merits of using the Screening Tool for Risk on Nutritional status and Growth (STRONG_{kids}) as a nutrition screening tool in hospitalized children aged < 3 years and correlate it with the severity of their nutritional derangements.

Methods: This cross-sectional study was conducted on 500 children aged < 3 years admitted to the Children's Hospital, Ain Shams University, Cairo, Egypt. STRONG_{kids} score was used to assess the risk for nutritional derangements and World Health Organization growth charts were used to define underweight, wasted and stunted patients upon admission and discharge.

Results: According to STRONG_{kids} score, 19.6% of patients were low risk, 42.6% were moderate risk and 37.8% were high risk. Out of the enrolled patients, 62.4% were underweight, 58.4% were stunted and 57.8% were wasted. Among the 66 patients with severe wasting, nutritional status improved in 6.06% while deterioration was observed in 13.0% of the moderately wasted patients. STRONG_{kids} score was worse among those who deteriorated, which together with its significant positive correlation with the duration of hospital stay, emphasized that STRONG_{kids} score can be a predictive tool.

Conclusions: The use of STRONG_{kids} screening tool can ensure early identification of children vulnerable to malnutrition, ensuring prompt interventions that may contribute to overall improvements in patient care, as well as shortening hospitalization period.

Keywords: malnutrition, screening, underweight, paediatrics, hospital.

Citation: Shaaban S; Nassar M; El-Gendy Y; El-Shaer B. Nutritional risk screening of hospitalized children aged < 3 years. East Mediterr Health J. 2019;25(1):18–23. <https://doi.org/10.26719/emhj.18.019>

Received: 13/04/17; accepted: 26/09/17

Copyright © World Health Organization (WHO) 2019. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license (<https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Introduction

Malnutrition in hospitalized children is an important pathological condition and a risk factor for unfavourable outcomes, prolonged hospital stay, delayed recovery and increased care costs. Reduction of dietary intake and increased energy requirements are the main causes of hospital undernutrition (1). The reported prevalence of acute malnutrition in infants and children admitted to hospitals from different countries ranges from 6.1 to 40.9% (2). In children with an underlying disease, higher prevalence of chronic malnutrition (44–64%) was reported in several studies (3).

To prevent hospital-acquired malnutrition, the risk of nutritional depletion needs to be identified as soon as possible, ideally at admission, so that appropriate nutritional intervention can be initiated at an early stage (4). Routine nutritional screening is rarely carried out in paediatric patients because of the lack of a simple and properly validated nutritional screening tool. The current practice of identifying children at risk of malnutrition is reliant on interpretation of anthropometric data and clinical judgement; the reliability of which is dependent on nutritional knowledge of paediatricians (5). Severe cases of malnutrition are easily recognized; however, the identification of children with lesser degrees

of malnutrition or at risk of malnutrition, which is also important, is not as easily achieved. Reports of malnutrition prevalence among hospitalized Egyptian infants and children are lacking.

This study was thus designed to assess the merits of using the Screening Tool for Risk on Nutritional status and Growth (STRONG_{kids}) as a nutrition screening tool in hospitalized Egyptian children aged < 3 years and correlate it with the severity of their nutritional derangements.

Methods

This cross-sectional study was conducted on 500 newly hospitalized children recruited from the Children's Hospital, Ain Shams University, Cairo, Egypt, between 1 January and 31 July 2015. There were 297 boys (59.4%) and 203 girls (40.6%). Their mean age was 13.73 [standard deviation (SD) 10.68] months with a range of 1–36 months; 315 patients (63%) were ≤ 12 months old and 185 (37%) were > 12 months old. The mean hospital stay was 6.62 (3.85) days with a range of 2–14 days. They were classified, as surgical or nonsurgical patients and underlying diseases were explored clinically and using laboratory and imaging assessment methods.

For all enrolled children aged < 3 years, we recorded age, sex, diagnosis and length of hospital stay. Nutritional

status was assessed using $STRONG_{kids}$ and complete anthropometric evaluation of body weight, body length/height, weight for length/height, skinfold thickness and mid arm circumference was done upon admission and discharge. $STRONG_{kids}$ is an easy to apply nutritional risk screening tool developed according to the latest European Society for Parenteral and Enteral Nutrition (ESPEN) guidelines (6). It consists of 4 elements: subjective clinical assessment, high-risk disease, nutritional intake and weight loss or poor weight gain. It is a comprehensive summary of commonly asked questions concerning nutritional issues, combined with a clinical view of the child's status. Each of the 4 elements of the questionnaire was allocated a score of 1 or 2 points with a maximum total score of 5 points. Patients obtaining 0 points were considered low risk; 1–3 points, moderate risk; and 4 or 5 points, high risk.

Anthropometric measurements were estimated by 2 trained investigators (Y. El-Gendy and B. El-Shaer). Height was measured to the nearest 0.1 cm with a portable stadiometer (Marsden, Rotherham, UK) with children standing bare foot, and recumbent length was measured by an infantometer (Model 416; Seca, Hamburg, Germany). Body weight was recorded to the nearest 0.1 kg using a calibrated baby scale (Model 834; Seca, Germany), with the patients' wearing only underpants or a clean diaper. Triceps skinfold thickness was measured vertically over the left triceps muscle midway between the acromion and olecranon process using a triceps skinfold caliper (Beta Technology Inc., Houston, TX, USA). Mid arm circumference was measured to the nearest centimetre using a nonstretchable tape (Butterfly, China), with the left arm hanging and relaxed in a sitting or lying position, midway between the tip of the acromion and the olecranon process.

Children with malnutrition were divided according to the World Health Organization (WHO) Global Database on Child Growth and Malnutrition, which uses a Z-score cutoff point of < -2 SD to classify low weight-for-age, low height-for-age and low weight-for-height as moderate undernutrition, and < -3 SD to define severe

undernutrition (7).

IBM SPSS version 20 was used for data analysis. Descriptive statistics were generated and numbers and percentages were used. Multivariate logistic regression analysis was performed for predictors of higher $STRONG_{kids}$ score. Correlation studies were demonstrated in figures and r values provided ($P < 0.05$ was considered significant).

Results

According to disease type 86 (17.2%) patients had chronic illnesses and 414 (82.8%) had acute conditions; the most common causes of which were chest infection in 190 (38%) and gastroenteritis in 176 (35.2%). According to $STRONG_{kids}$ score, 98 (19.6%) patients were classified as low risk, 213 (42.6%) as moderate risk and 189 (37.8%) as high risk. Table 1 shows the details of the points given to the screened patients.

Two hundred and eighty-nine (57.8%) patients were underweight (weight for age < -2 Z score), 292 (58.4%) were stunted (height for age < -2 Z score) and 312 (62.4%) were wasted (weight for height < -2 Z score). Among the wasted cases, 66 had severe wasting and the rest moderate wasting. Table 2 shows that among the 66 patients with severe wasting, nutritional state was not altered in 62 (93.93%) while it improved in 4 (6.06%) who became moderately wasted. Nutritional deterioration was observed in 32 (13.00%) children, who had been moderately wasted at admission and progressed to severe wasting, while 214 (86.99%) remained moderately wasted. Also nutritional deterioration was observed in 6 (3.19%) children who had been normal at admission and progressed to moderate wasting while 182 (96.8%) remained normal.

Five of the 6 patients in the normal weight for height group and 28 of the 32 patients in the moderate wasting group who deteriorated were high risk according to $STRONG_{kids}$ score. Three of the 4 severely wasted patients who improved were moderate risk according to $STRONG_{kids}$ score, and the other one was high risk. Among the 62 severely wasted patients who showed no

Table 1 Nutritional risk screening tool $STRONG_{kids}$ results among the studied series

	Yes	No
(1) Subjective clinical assessment (1 point) Is the patient in a poor nutritional status judged by subjective clinical assessment (diminished subcutaneous fat and/or muscle mass and/or hollow face)?	215 (43%)	285 (57%)
(2) High-risk disease (2 points) Is there an underlying illness with a risk of malnutrition or expected major surgery?	270 (54%)	230 (46%)
(3) Nutritional intake and losses (1 point) Is one of the following items present? Excessive diarrhoea (> 5 times/day) and/or vomiting (> 3 times/day) in the last few days? Reduced food intake during the last few days before admission (not including fasting for an elective procedure or surgery)? Pre-existing dietetically advised nutritional intervention? Inability to consume adequate intake because of pain?	378 (75.6%)	122 (24.4%)
(4) Weight loss or poor weight gain? (1 point) Is there weight loss or no weight gain (infants aged < 1 year) during the last few weeks/months?	440 (88%)	60 (12%)

Table 2 Follow-up of nutritional status in children aged < 3 years during hospitalization according to weight for height Z score

Classification	Patient progress		
	Severe wasting	Moderate wasting	Normal
Severe wasting, 66 (100%)	62 (93.93%)	4 (6.06%)	0 (0.00%)
Moderate wasting, 246 (100%)	32 (13.00%)	214 (86.99%)	0 (0.00%)
Normal weight for height, 188 (100%)	0 (0.00%)	6 (3.19%)	182 (96.8%)
Total, 500 (100%)	94 (18.8%)	224 (44.8%)	182 (36.4%)

improvement, 51 were high risk and 11 moderate risk.

Figure 1 demonstrates a significant positive correlation between $STRONG_{kids}$ score and duration of hospital stay ($r = 0.114, P = 0.01$). However, there was a significant negative correlation between $STRONG_{kids}$ score and maternal education ($r = -0.633, P = 0.005$). The logistic regression showed that after elimination of all other factors, there was significant association between higher $STRONG_{kids}$ score and each of the following: low maternal education, high duration of hospital stay and low admission weight for age (Table 3).

Discussion

We showed that 17.2% of the patients had chronic illnesses and 82.8% had acute ones. The most common acute illnesses were chest infection in 38% and gastroenteritis in 35.2%. This patient profile is similar to that of Silveira et al. (8) and Saccardo Sarni et al. (9) who reported that respiratory diseases were the main reason for hospitalization. Additionally, Rocha et al. (5) found that the most frequent disease responsible for hospital admission was pneumonia (33%) followed by diarrhoea (6.4%). The noticeable difference in the current study figures is the percentage of hospitalization from gastroenteritis compared to pneumonia, which still has a high disease burden, despite the various preventive efforts of Egyptian governmental and nongovernmental agencies.

According to WHO cutoff values, 62.4% of our patients were underweight, 58.4% were stunted and

57.8% were wasted, which are higher than those for children aged < 5 years (6%, 21% and 8%, respectively) reported in the 2014 Egyptian Demographic and Health Survey (10). Although Rocha et al. (5) reported lower figures for underweight, stunting and wasting in Brazil (18.7, 18.2 and 6.9%, respectively), they mentioned that hospital malnutrition in Latin America can reach up to 70–80%, which agrees with our results. Ozturk et al. (11) found that 31.8% of hospitalized children in Turkey were malnourished and added that well-nourished children do not carry nutritional risk due to hospitalization for other medical reasons. Another Turkish study by Dogan et al. (12) reported that 27% of the hospitalized patients were stunted, 52.4% were underweight and 40.9% were wasted, which is closer to the results in the current study. Malnutrition rates of 32% among hospitalized children in Turkey (13) and 60% among hospitalized children in Thailand (14) further demonstrate the diversity of the published results.

Nutritional deterioration was observed in 13% of the moderately wasted children and 3.19% of patients who were normal at admission. Ferreira and França (15) observed that 20% of children who were well nourished upon admission became malnourished. Rocha et al. (5) reported that 51.6% of 186 hospitalized children lost weight and 9.17% of well-nourished children developed mild malnutrition during hospitalization. Pacheco-Acosta et al. (16), also reported nutritional deterioration in their series of hospitalized children with nonserious disease and advised early




Figure 1 Correlation between $STRONG_{kids}$ score and duration of hospital stay.

Table 3 Multivariate logistic regression analysis for predictors of higher STRONG_{kids} score

	Significance	OR	95% CI
Age	0.221 (NS)	2.009	0.988–3.029
Maternal education	0.002 (S)	8.007	1.979–10.036
Admission weight	0.01 (S)	6.989	1.952–8.027
Duration of hospital stay	0.001 (S)	8.022	1.995–10.049
Cause of admission	0.128 (NS)	1.071	0.896–1.280

detection of children at risk to enable early interventions. Special attention should be paid to children who are already malnourished upon admission, as they are at risk of further nutritional deterioration during their hospital stay (17).

In the current study 42.6% of the patients were at moderate risk and 37.8% at high risk of developing malnutrition according to STRONG_{kids}. These percentages are higher than the risk score recorded in a multicentre Dutch study by Hulst et al. (18). The latter authors reported that 62% of hospitalized children were classified at moderate and high risk by the STRONG_{kids} tool. In Romania 58% of children were found to be at risk of malnutrition (24% high risk) by the STRONG_{kids} tool (19). The higher figures in the present study can be attributed to the initial increased incidence of underweight, stunting and wasting, as well as persistence of the high rate of hospitalization for gastroenteritis which, if prolonged, can affect weight greatly (20).

The current study showed significant associations between STRONG_{kids} score and both prolonged hospital stay and low admission weight for age. These findings are consistent with the above-mentioned Dutch study (18), which predicted a significant relationship between high risk score, a negative SD score for weight for height, and prolonged hospital stay. Several other studies have also documented that malnourished patients stay longer in hospital than well-nourished patients (21,22), confirming the need for early detection of such vulnerable patients.

The results of the current study showed that patients whose nutritional status deteriorated had initial high risk score by STRONG_{kids} compared to moderate risk score for those who improved. In retrospect, this emphasizes that STRONG_{kids} can be a predictive prognostic tool. Similarly,

Sermet-Gaudelus et al. (23) advised implementation of their simple pediatric nutrition risk score to prevent hospital-acquired malnutrition. In contrast, Huysentruyt and associates (24) did not find a significant correlation between STRONG_{kids} risk categories and weight loss during hospitalization. However, they mentioned that these categories correlated with the length of hospital stay and establishment of nutritional intervention during hospitalization.

To our knowledge, this is the first study to explore the STRONG_{kids} as a screening and prognostic tool in an Egyptian hospital setting. Nevertheless, this study has its limitations; mainly in the small sample size and lack of long-term follow-up. Additionally, there should be a larger multicentre study including other age groups from all over Egypt to allow us to draw conclusions on a nationwide basis.

Conclusion

Use of the STRONG_{kids} for screening hospitalized children aged < 3 years revealed that nearly 80% were at risk of nutritional derangements, and its scores correlated positively with the length of hospital stay and negatively with body weight at admission. Moreover, the overall malnutrition among these children is a significant problem and patients whose nutritional status deteriorated had higher STRONG_{kids} scores. We thus recommend implementation of the STRONG_{kids} nutritional risk assessment tool for early screening of hospitalized Egyptian children to avoid prolonged hospitalization and further compromise in their nutritional status.

Funding: None.

Competing interests: None declared.

Évaluation du risque nutritionnel des enfants de plus de trois ans hospitalisés

Résumé

Contexte : L'identification des enfants à risque de malnutrition n'est pas une entreprise facile en milieu hospitalier.

Objectifs : Évaluer les mérites du recours à l'outil d'évaluation du risque pour l'état nutritionnel et la croissance (STRONG_{kids}) en tant qu'outil de dépistage nutritionnel pour les enfants de plus de trois ans hospitalisés et le corrélér à la sévérité des troubles nutritionnels.

Méthodes : La présente étude transversale a été réalisée auprès de 500 enfants de plus de trois ans admis à l'hôpital des enfants de l'Université d'Ain Shams, au Caire (Égypte). Le score STRONG_{kids} a été utilisé pour évaluer le risque de trouble nutritionnel et les diagrammes de croissance de l'Organisation mondiale de la Santé ont été employés pour définir les enfants présentant une insuffisance pondérale, une émaciation et un retard de croissance à l'admission et à la sortie d'hôpital.

Résultats : Selon le score $STRONG_{kids}$, 19,6 % des patients présentaient un faible risque, 42,6 % un risque modéré et 37,8 % un risque élevé. Sur les patients participant à l'étude, 62,4 % avaient une insuffisance pondérale, 58,4 % une émaciation et 57,8 % un retard de croissance. Sur les 66 patients présentant une émaciation sévère, l'état nutritionnel s'est amélioré pour 6,06 % et on a observé une détérioration de cet état chez 13,0 % des patients ayant une émaciation modérée. Le score $STRONG_{kids}$ était pire chez ceux ayant connu une détérioration de leur état, ce qui, en association avec sa corrélation positive significative avec la durée du séjour hospitalier, soulignait le fait que le score $STRONG_{kids}$ peut constituer un outil prédictif.

Conclusions : L'utilisation de l'outil de dépistage $STRONG_{kids}$ peut permettre l'identification précoce des enfants vulnérables à la malnutrition, ainsi que la mise en place d'interventions rapides qui peuvent contribuer à des améliorations générales des soins aux patients, ainsi qu'à une réduction de la période d'hospitalisation.

تحري المخاطر التغذوية لدى الأطفال الذين تقل أعمارهم عن ٣ سنوات داخل المستشفيات

سناء شعبان، مي نصار، ياسمين الجندي، بسام الشاعر

الخلاصة

الخلفية: لا يجري تحديد الأطفال المعرضين لخطر سوء التغذية بسهولة داخل المستشفيات.

الأهداف: هدفت الدراسة إلى تقييم مزايا استخدام أداة الكشف عن المخاطر التي تهدد الحالة التغذوية والنمو ($STRONG_{kids}$) كأداة للكشف عن سوء التغذية لدى الأطفال الذين تقل أعمارهم عن ٣ سنوات داخل المستشفيات، وربطها بمدى شدة الاختلالات التغذوية لديهم.

طرق البحث: أُجريت هذه الدراسة المقطعية على ٥٠٠ طفل تقل أعمارهم عن ٣ سنوات والذين أُدخلوا إلى مستشفى الأطفال، جامعة عين شمس، القاهرة، مصر. واستُخدمت أداة $STRONG_{kids}$ في تقييم خطر الاختلالات التغذوية، كما استُخدمت مخططات النمو لمنظمة الصحة العالمية في تحديد نقص الوزن والهزال ونقص النمو لدى المرضى عند إدخالهم إلى المستشفى وإخراجهم منها.

النتائج: وفقاً لأداة $STRONG_{kids}$ ، كان ١٩,٦ % من المرضى معرضين لخطر منخفض، و ٤٢,٦ % معرضين لخطر متوسط، و ٣٧,٨ % معرضين لخطر مرتفع. ومن بين المرضى المسجلين، كان ٦٢,٤ % يعانون من نقص الوزن، و ٥٨,٤ % يعانون من نقص النمو و ٥٧,٨ % يعانون من الهزال. ومن بين ٦٦ مريضاً يعانون من شدة الهزال، تحسنت الحالة التغذوية لدى ٦,٠٦ %، بينما لوحظ تدهور لدى ١٣,٠ % من المرضى الذين يعانون من هزال متوسط. وقد كانت نقاط $STRONG_{kids}$ أسوأ لدى المرضى الذين تدهورت حالتهم هذا إلى جانب ارتباطها الإيجابي الكبير مع مدة الإقامة في المستشفى مما يثبت انه يمكن استخدام أداة $STRONG_{kids}$ بمثابة أداة تنبؤية.

الاستنتاجات: يمكن أن يضمن استخدام أداة التحري $STRONG_{kids}$ للتشخيص المبكر للأطفال المعرضين لخطر سوء التغذية، مما يساعد على التدخل الفوري الذي يساهم في تحسين رعاية المرضى بشكل عام ويقلل من مدة الإقامة في المستشفى.

References

1. Bejon P, Mohammed S, Mwangi I, Atkinson SH, Osier F, Peshu N, et al. Fraction of all hospital admissions and deaths attributable to malnutrition among children in rural Kenya. *Am J Clin Nutr.* 2008 Dec;88(6):1626–31. <http://dx.doi.org/10.3945/ajcn.2008.26510> PMID:19064524
2. Pawellek I, Dokoupil K, Koletzko B. Prevalence of malnutrition in paediatric hospital patients. *Clin Nutr.* 2008 Feb;27(1):72–6. <http://dx.doi.org/10.1016/j.clnu.2007.11.001> PMID:18086508
3. Joosten KF, Zwart H, Hop WC, Hulst JM. National malnutrition screening days in hospitalized children in the Netherlands. *Arch Dis Child.* 2010 Feb;95(2):141–5. <http://dx.doi.org/10.1136/adc.2008.157255> PMID:19414435
4. Mahdavi AM, Ostadrahimi A, Safaiyan A. Nutritional status of children hospitalized in Tabriz Paediatric Hospital, Islamic Republic of Iran, 2008. *East Mediterr Health J.* 2011 Jan;17(1):36–40. <http://dx.doi.org/10.26719/2011.17.1.36> PMID:21735799
5. Rocha GA, Rocha EJ, Martins CV. The effects of hospitalization on the nutritional status of children. *J Pediatr (Rio J).* 2006 Jan–Feb;82(1):70–4. <http://dx.doi.org/10.2223/JPED.1440> PMID:16532151
6. Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr.* 2003 Aug;22(4):415–21. [http://dx.doi.org/10.1016/S0261-5614\(03\)00098-0](http://dx.doi.org/10.1016/S0261-5614(03)00098-0) PMID:12880610
7. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. WHO Technical Report Series No. 854. Geneva: World Health Organization; 1995 (https://www.who.int/childgrowth/publications/physical_status/en/, accessed 9 January 2019).
8. Silveira KB, Alves JF, Ferreira HS, Sawaya AL, Florêncio TM. Association between malnutrition in children living in slums, maternal nutritional status, and environmental factors. *J Pediatr (Rio J).* 2010 May–Jun;86(3):215–20. <http://dx.doi.org/10.1590/S0021-75572010000300009> PMID:20440445

9. Saccardo Sarni RO, Suano de Souza FI, Catherino P, Kochi C, Ceragioli Oliveira FL, de Nóbrega FJ. [Treatment of severe malnourished children with WHO protocol: experience of a referral center in São Paulo, Brazil]. *Arch Latinoam Nutr.* 2005 Dec;55(4):336–44 (in Portuguese). PMID:16640196
10. Ministry of Health and Population (Egypt), El-Zanaty and Associates (Egypt), and ICF International. *Egypt Demographic and Health Survey 2014*. Cairo: Ministry of Health and Population; Rockville, MD: ICF International; 2015 (<https://dhsprogram.com/pubs/pdf/fr302/fr302.pdf>, accessed 20 December 2018).
11. Öztürk Y, Büyükgöbüz B, Arslan N, Ellidokuz H. Effects of hospital stay on nutritional anthropometric data in Turkish children. *J Trop Pediatr.* 2003 Jun;49(3):189–90. <http://dx.doi.org/10.1093/tropej/49.3.189> PMID:12848214
12. Doğan Y, Erkan T, Yalvaç S, Altay S, Cokuğraş FC, Aydın A, et al. Nutritional status of patients hospitalized in pediatric clinic. *Turk J Gastroenterol.* 2005 Dec;16(4):212–6. PMID:16547850
13. Joosten KF, Hulst JM. Prevalence of malnutrition in pediatric hospital patients. *Curr Opin Pediatr.* 2008 Oct;20(5):590–6. <http://dx.doi.org/10.1097/MOP.0b013e32830c6ede> PMID:18781124
14. Tienboon P. Nutrition problems of hospitalised children in a developing country: Thailand. *Asia Pac J Clin Nutr.* 2002;11(4):258–62. <http://dx.doi.org/10.1046/j.1440-6047.2002.00307.x> PMID:12495256
15. Ferreira HS, França OSA. [Evolution of nutritional status in hospitalized children]. *J Pediatr (Rio J).* 2002 Nov-Dec;78(6):491–6 (in Portuguese). PMID:14647730
16. Pacheco-Acosta JC, Gomez-Correa AC, Florez ID, Cortés JE, Velez D, Gomez J, et al. Incidence of nutrition deterioration in nonseriously ill hospitalized children younger than 5 years. *Nutr Clin Pract.* 2014 Oct;29(5):692–7. <http://dx.doi.org/10.1177/0884533614533122> PMID:24888729
17. Campanozzi A, Russo M, Catucci A, Rutigliano I, Canestrino G, Giardino I, et al. Hospital-acquired malnutrition in children with mild clinical conditions. *Nutrition.* 2009 May;25(5):540–7. <http://dx.doi.org/10.1016/j.nut.2008.11.026> PMID:19230617
18. Hulst JM, Zwart H, Hop WC, Joosten KF. Dutch national survey to test the STRONG(kids) nutritional risk screening tool in hospitalized children. *Clin Nutr.* 2010 Feb;29(1):106–11. <http://dx.doi.org/10.1016/j.clnu.2009.07.006> PMID:19682776
19. Mărginean O, Pitea AM, Voidăzan S, Mărginean C. Prevalence and assessment of malnutrition risk among hospitalized children in Romania. *J Health Popul Nutr.* 2014 Mar;32(1):97–102. PMID:24847598
20. Giannattasio A, Guarino A, Lo Vecchio A. Management of children with prolonged diarrhea. *F1000Res.* 2016 Feb 23. <http://dx.doi.org/10.12688/f1000research.7469.1> PMID:26962439
21. Aurangzeb B, Whitten KE, Harrison B, Mitchell M, Kepreotes H, Sidler M, et al. Prevalence of malnutrition and risk of under-nutrition in hospitalized children. *Clin Nutr.* 2012 Feb;31(1):35–40. <http://dx.doi.org/10.1016/j.clnu.2011.08.011> PMID:21945311
22. Joosten KFM, Hulst JM. Malnutrition in pediatric hospital patients: current issues. *Nutrition.* 2011 Feb;27(2):133–7. <http://dx.doi.org/10.1016/j.nut.2010.06.001> PMID:20708380
23. Sermet-Gaudelus I, Poisson-Salomon AS, Colomb V, Brusset MC, Mosser F, Berrier F, et al. Simple pediatric nutritional risk score to identify children at risk of malnutrition. *Am J Clin Nutr.* 2000 Jul;72(1):64–70. <http://dx.doi.org/10.1093/ajcn/72.1.64> PMID:10871562
24. Huysentruyt K, Alliet P, Muyschont L, Rossignol R, Devreker T, Bontems P, et al. The STRONGkids nutritional screening tool in hospitalized children: A validation study. *Nutrition.* 2013 Nov-Dec;29(11-12):1356–61. <http://dx.doi.org/10.1016/j.nut.2013.05.008> PMID:24103513