



Review Article

Targeted therapies in the management of breast cancer

G.O. Abdulrahman Jnr.

Institute of Medical Education, Cardiff University School of Medicine, Cardiff, UK

Abstract

The incidence of cancer is increasing in the United Kingdom. Breast cancer is the most commonly diagnosed cancer among women in the UK with an age standardised rate of 124 per 100000 women. It accounts for almost one-third of all female cases of cancer in the UK and it is the most common form of cancer among women in both developing and developed countries. There are many management options for cancer such as surgery, chemotherapy and radiotherapy. Surgery has been the main treatment option for many solid tumours for several decades. However, research has shown that attacking specific targets within tumours such as receptors, intracellular proteins and genes could result in better

clinical outcome. Promising therapeutic targets that have been identified include Raf kinase, Src, HER2, epidermal growth factor receptor and vascular endothelial growth factor receptor. After several decades of intensive research, it appears that we are finally hitting cancer where it hurts. It is very likely that in the near future, more targets for potential therapies would be identified and ultimately, there would be significant reduction in morbidity and mortality from cancers, including breast cancer.

Key words

Breast cancer, HER2, BRCA genes, Raf kinase, anti-microtubule, targeted therapies

Introduction

The incidence of cancer is increasing in the United Kingdom. In 2011, approximately 331000 people were diagnosed with cancer in the UK, more than half of which are breast, lung, colorectal and prostate. ⁽¹⁾ Cancer continues to be a major killer despite the rigorous research and rapid developments in recent years, and the incidence and mortality are likely to increase in the near future. It is expected that the world population would increase to 7.5 billion by 2020 and of this number, approximately 16 million new cancer cases will be diagnosed and 12 million cancer patients will die. ⁽²⁾ Approximately 90–95% of cases can be attributed to environmental factors and lifestyle such as smoking, alcohol, diet and infections. ⁽³⁾ Only 5–10% of all cancer cases can be attributed to genetic defects. ⁽³⁾ There are many management options for cancer such as surgery, chemotherapy and radiotherapy. Surgery has been the main treatment option for many solid tumours for several decades; however research has shown that attacking specific targets within tumours

such as receptors, intracellular proteins and genes could result in better clinical outcome. Promising therapeutic targets that have been identified include Raf kinase ⁽⁴⁾, Src ⁽⁵⁾ and HER2 ⁽⁶⁾. The aim of this review is to highlight some of the therapeutic targets that have been identified in the treatment of solid tumours, with particular emphasis on breast cancer – the most common cancer in women in the UK.

Breast cancer

Breast cancer is the most commonly diagnosed cancer among women in the UK with almost 50000 women diagnosed with the malignancy in 2011. ⁽¹⁾ It accounts for almost one-third of all female cases of cancer in the UK ⁽¹⁾ and it is the most common

Corresponding author: **Ganiy Opeyemi Abdulrahman, Jnr**, Currently an Academic Clinical Fellow / Specialty Trainee, Department of Obstetrics & Gynaecology, Women's Services, Level 5, Leazes Wing, Royal Victoria Infirmary (RVI), Queen Victoria Road, Newcastle upon Tyne, Tyne and Wear NE1 4LP, United Kingdom
Email: ussheadie@hotmail.co.uk

form of cancer among women in both developing and developed countries. ⁽⁷⁾ The age-standardised incidence rate of breast cancer in Wales, United Kingdom is 113.4 per 100,000. ⁽⁸⁾ Certain risk factors have been associated with breast cancer. These include age, sex, obesity, family history of breast cancer, late menopause and long-term use of hormone replacement therapy. Like many other cancers, breast cancer is caused by an interaction between the environment and genetic defect. During cell division, DNA is usually copied with several mistakes, which is normally repaired by error-correcting proteins or undergo apoptosis. When there is mutation in these error-correcting mechanisms, the cells begin to divide uncontrollably resulting in tumourigenesis. A number of mutations have been implicated in the pathophysiology of breast cancer. These include mutations in the RAS-RAF-MEK-ERK pathway, ⁽⁹⁾ PI3K/AKT pathway, ⁽¹⁰⁾ breast cancer susceptibility genes 1 and 2 (BRCA1 and BRCA2) ⁽¹⁰⁾ and HER2. ⁽¹⁰⁾ As a result of increasing knowledge about the cellular mechanisms that underlie the genesis of cancer, specific targeted therapies are being developed that block the growth and spread of cancer by interfering with specific molecules that are involved in the growth and progression of tumours. It is thought that this therapeutic strategy may be more effective as it is more specific than treatments such as chemotherapy and radiotherapy which are non-specific and usually have many side effects.

HER2 inhibitors

HER2 belongs to the human epidermal growth factor receptor (HER) family of tyrosine kinases which consists of EGFR (HER1, erbB1), HER2 (erbB2, HER2/neu), HER3 (erbB3) and HER4 (erbB4). ⁽¹¹⁾ The HER2 gene is located on the long arm of human chromosome 17 and encodes a transmembrane receptor protein with tyrosine kinase activity. ^(12,13) Structurally, the receptor is made up of an extracellular ligand-binding domain, transmembrane domain and an intracellular tyrosine kinase catalytic domain. ⁽¹⁴⁾ On activation by a ligand, the receptors dimerize and undergo transphosphorylation in order to activate the various intracellular signaling pathways which mediate the proliferation and differentiation of cells. ^(15,16) HER proteins control cell growth, survival, adhesion, migration and differentiation. The cellular

mechanism by which HER2 is activated is not entirely understood and it is believed that it is an orphan receptor as there is no known stimulatory ligand for HER2 receptor homodimers. ⁽¹⁴⁾

The amplification of HER2 has been implicated in some cancers. Slamon and colleagues reported that approximately 25–30% of breast cancers have amplification and overexpression of HER2, with such cancers having worse biologic behaviour and prognosis. ⁽¹⁷⁾ When alterations of the gene in 189 primary human breast cancers were investigated, HER2/neu oncogene was found to be amplified from 2- to greater than 20-fold in 30% of the tumours. ⁽¹⁸⁾ It soon became clear that there is a correlation between the amplification of HER2 and the genesis of human cancers. HER2 amplification or protein overexpression is now used to predict clinical outcome and prognosis in clinical practice. HER2 amplification or protein overexpression has been shown to be a poor predictor of clinical outcome in node-positive patients. ⁽¹⁶⁾ It is also associated with increased tumour recurrence and decreased survival.

As a result of the correlation between HER2 amplification and tumourigenesis, it was proposed that inhibiting oncogenic HER2 could be an effective treatment for HER2-driven cancers. This has resulted in the development and use of anti-HER2 antibodies such as trastuzumab (Herceptin) in clinical management. Herceptin is an anti-HER2 monoclonal antibody with an anti-proliferative effect on cells transformed by HER2 overexpression and is an effective treatment for breast cancer. ⁽¹⁹⁾ It is thought that herceptin stimulates HER2 endocytosis and removal of HER2 from the cell surface. Sliwkowski and colleagues claim that trastuzumab induces p27KIP1 – a tumour suppressor protein, and Rb-related protein, p130, which in turn significantly reduces the number of cells undergoing S-phase. ⁽²⁰⁾ Herceptin downmodulates HER2 receptor, inhibits tumour cell growth, reverses cytokine resistance, restores E-cadherin expression levels and reduces vascular endothelial growth factor production. ⁽²⁰⁾ The use of herceptin in clinical practice has resulted in significant reduction in mortality from HER2-overexpressing disease. In an American trial to compare adjuvant chemotherapy with or without concurrent trastuzumab in women with surgically removed HER2-positive breast cancer, trastuzumab

therapy was associated with a 33% reduction in the risk of death ($P=0.015$).⁽²¹⁾

Lapatinib is another drug that has been developed which interrupts the HER2 growth receptor pathway. Lapatinib inhibits the tyrosine kinase activity associated with two oncogenes – EGFR and HER2/neu.⁽²²⁾ Lapatinib works intracellularly by reversibly binding to the cytoplasmic ATP-binding site of the kinase, therefore blocking receptor phosphorylation and activation.⁽²³⁾ This results in the suppression of proliferation pathways of solid tumours, most notably advanced or metastatic breast cancer.⁽²⁴⁾ In a study by Cameron and associates, the use of lapatinib in combination with capecitabine in women with HER2-positive, locally advanced and metastatic breast cancer resulted in improved overall survival and prolonged time to progression.⁽²⁵⁾

In view of the success of HER2 targeting agents, it is very likely that in the coming decade, a second generation of HER2-targeting agents would be introduced into clinical testing in an attempt to treat HER2-driven cancers via inactivation of HER2. With a better understanding of the cellular mechanisms by which HER2 is activated, more novel therapies are likely to be developed with different mechanisms of action.

BRCA genes

Breast cancer susceptibility genes 1 and 2 (BRCA1 and BRCA2) are tumour suppressor genes. In normal cells, BRCA 1 and BRCA 2 help to ensure the stability of the cell's genetic material and prevent uncontrolled cell growth.⁽²⁶⁾ Mutation of BRCA1 and BRCA2 genes have been shown to be associated with an increased risk of developing breast cancer. BRCA1 and BRCA2 both interact with RAD51 – a protein – in the repair of DNA double-strand breaks.⁽²⁷⁾ When BRCA1 or BRCA2 gene becomes defective, damaged DNAs begin to accumulate as the repair mechanism is lost, resulting in uncontrolled division and tumorigenesis. In a study by Risch and colleagues, they claim that the estimated cumulative incidence of breast cancer to age 80 years among women carrying the BRCA1 and BRCA2 mutations was 90% and 41% respectively.⁽²⁸⁾ It is therefore evident that mutation in BRCA1 or BRCA2 gene increases the risk of developing breast cancer. Hence it has been suggested that specifically targeting the mechanism for single-strand DNA

repair on which the tumour cells depend could offer a therapeutic strategy in inhibiting tumour growth and progression since the double-strand DNA repair mechanism is already damaged. This resulted in the development of PARP inhibitors. Poly ADP ribose polymerase (PARP) is a protein that plays a crucial role in the repairing of single-strand breaks, particularly PARP1 which acts as a sensor of DNA damage and initiate base excision repair pathway.⁽²⁹⁾ In a phase 1 trial by Fong and colleagues, they administered olaparib (a PARP inhibitor) to patients who carry the BRCA1 or BRCA2 mutation and reported anti-tumour activity in all the mutation carriers.⁽³⁰⁾ Although no lethal adverse effects were reported, it is important to study the potential long-term adverse effects of PARP inhibitors. This is because PARP1 has several other important roles including restarting stalled replication, inhibiting non-homologous end-joining repair, regulating transcription, initiating a unique cell death pathway and modulating cellular biogenesis.⁽³¹⁾ Hence inhibiting PARP may significantly affect these processes, which would be detrimental to cells.

Although there may be potential drawbacks to the use of PARP inhibitors in the treatment of breast cancers, it would be interesting to see the results of the Phases II and III trials as there is a potential that clinical benefits would be produced in BRCA-deficient tumours.

Rapidly Accelerated Fibrosarcoma (Raf) kinase inhibitors

Raf kinases are a family of three serine/threonine-specific protein kinases that are related to retroviral oncogenes and were discovered in 1983.⁽³²⁾ Members of the RAF family include A-RAF, B-RAF and C-RAF (formerly known as RAF-1). RAF kinases participate in the RAS-RAF-MEK-ERK signal transduction cascade, which is sometimes denoted as the mitogen-activated protein kinase (MAPK) cascade.⁽³²⁾ Activation of RAF kinase activity is a multistage process requiring interaction with RAS-GTP. The RAF kinases have restricted substrate specificity and catalyze phosphorylation and activation of MEK1 and MEK2. [33] MEK1/2, which are dual specificity protein kinases, mediate the phosphorylation of tyrosine before threonine in ERK1 or ERK2 (Extracellular-signal Regulated protein Kinase) – their only substrates.⁽³³⁾ The phosphorylation activates ERK1/2,

which are protein–serine/threonine kinases.⁽³³⁾ Unlike RAF and MEK1/2 that have narrow substrate specificity, ERK1 and ERK2 have many substrates. This cascade is important in the regulation of several processes including apoptosis, cell cycle progression, differentiation, proliferation and transformation to the cancerous state.⁽³³⁾

The physiological regulation of RAF kinases is complex and involves several steps such as protein–protein interactions, phosphorylation, dephosphorylation and conformational changes.⁽³⁴⁾ The activation of oncogenic mutation within the upstream RAS gene implicated in tumourigenesis can result in constitutive activation of downstream RAF–1⁽³⁵⁾. This then induces DNA synthesis resulting in malignant transformation, loss of growth factor dependence and contact inhibition, increased cell survival and invasion/metastasis, all of which are pathognomonic features of cancerous cells. It has been claimed that oncogenic ras mutations occur in as many as 90% of pancreatic and half of thyroid and colorectal carcinomas.⁽³⁵⁾ RAS mutations have also been implicated in acute myeloid leukemia, melanoma and kidney tumours.^(35, 36) The activation of b–raf mutations also occur at a high frequency in certain human cancers, with the most common variant being b–raf V600E present in 63% of melanomas,⁽³⁷⁾ and 40% of sporadic colorectal tumours.⁽³⁸⁾ Mutations in k–ras and b–raf have also been implicated in human breast cancer cell lines. Although initially thought to be rare, k–ras mutation was noted in 13% of breast cancer cell lines studied by Hollestelle and colleagues while 25% of the breast cancer lines were found to have mutational activation of RAS signaling pathways.⁽³⁹⁾

There has been development of therapies targeting the RAS–RAF–MEK–ERK pathway with the aim of downregulating the cellular activities, therefore inhibiting tumour cell growth. Sorafenib has been developed to inhibit RAF isoforms. It is a synthetic, orally available bi–aryl urea and is the most promising of the RAF kinase inhibitors in clinical development.⁽⁴⁰⁾ In cellular assays, sorafenib inhibits the phosphorylation of ERK1/2 (pERK) and MEK 1/2 (pMEK), thus supporting its RAF inhibitory effects in human cells.^(41,42) It has also been demonstrated that sorafenib reduces pERK levels in tumour cell lines with oncogenic k–ras or b–raf genes.⁽⁴¹⁾ The inhibitory

effects of Sorafenib on RAF isoforms in vivo is also supported by preclinical studies demonstrating its dose–dependent growth inhibitory effect on human colon and breast tumour xenografts containing k–ras or b–raf mutations.⁽⁴¹⁾ Sorafenib has been shown to have activity against ovarian, pancreatic, breast and melanoma tumour xenograft models.⁽⁴¹⁾ In a study by Gradishar and associates, it was observed that sorafenib, in combination with paclitaxel, resulted in significant improvement in survival in patients with HER2–negative, locally recurrent or metastatic breast cancer.⁽⁴³⁾

It would be interesting to study the effectiveness of each of the targeting agents discussed above in relation to the genotype of patients. It may well be demonstrated, for example, that patients who are HER2–positive have better prognosis when treated with Trastuzumab or Lapatinib while patients who are HER2–negative have improved clinical outcome when treated with Sorafenib, rather than Trastuzumab.

Conclusion

Specific targeted therapies have been discussed with particular emphasis on breast cancer. Some of these therapies are still undergoing clinical trials and it would be interesting to see the results of these trials. It is paramount that researchers ascertain the optimal dosing of these targeting agents in order to achieve maximal anti–tumour effect with minimal or tolerable toxicity. After several decades of intensive research, it appears that we are finally hitting cancer where it hurts. It is very likely that in the near future, more targets for potential therapies would be identified and ultimately, there would be significant reduction in morbidity and mortality from cancers.

References

1. Cancer Research UK [Internet]. London: Cancer Research UK; c2014 [updated 2011 January; cited 2015 Jun 17]. Available from: http://publications.cancerresearchuk.org/downloads/Product/CS_REPORT_INCIDENCE.pdf
2. Bray F, Møller B. Predicting the future burden of cancer. *Nat Rev Cancer* 2006; 6(1): 63–74
3. Anand P, Kunnumakkara AB, Sundaram C, et al. Cancer is a preventable disease that requires major lifestyle changes. *Pharm Res* 2008; 25(9): 2097–116

4. Lyons JF, Wilhelm S, Hibner B, et al. Discovery of a novel Raf kinase inhibitor. *Endocr Relat Cancer* 2001; 8(3): 219–25
5. Nagathihalli NS, Merchant NB. Src-mediated regulation of E-cadherin and EMT in pancreatic cancer. *Front Biosci (Landmark Ed)* 2012; 17: 2059–69
6. Ko BK, Lee SY, Lee YH, et al. Combination of novel HER2-targeting antibody 1E11 with trastuzumab shows synergistic antitumor activity in HER2-positive gastric cancer. *Mol Oncol* 2015; 9(2): 398–408
7. Torre LA, Bray F, Siegel RL, et al. Global cancer statistics, 2012. *CA Cancer J Clin* 2015; 65(2): 87–108
8. Abdulrahman GO Jnr. Breast cancer in Wales: time trends and geographical distribution. *Gland Surg* 2014; 3(4): 237–42
9. McGlynn LM, Kirkegaard T, Edwards J, et al. Ras/Raf-1/ MAPK pathway mediates response to tamoxifen but not chemotherapy in breast cancer patients. *Clin Cancer Res* 2009; 15(4): 1487–95
10. McCubrey JA, Franklin RA, Bertrand FE, et al. Targeting survival cascades induced by activation of Ras/Raf/MEK/ ERK and PI3K/Akt pathways to sensitize cancer cells to therapy. In: Bonavida B, ed, *Sensitization of cancer cells for chemo/immuno/radio-therapy*. 1st ed. Totowa, NJ: Humana Press, 2008; 81–114
11. Moasser MM. Targeting the function of the HER2 oncogene in human cancer therapeutics. *Oncogene* 2007; 26(46): 6577–92
12. Fukushige S, Matsubara K, Yoshida M, et al. Localization of a novel v-erbB-related gene, c-erbB-2, on human chromosome 17 and its amplification in a gastric cancer cell line. *Mol Cell Biol* 1986; 6(3): 955–8
13. Akiyama T, Sudo C, Ogawara H, et al. The product of the human c-erbB-2 gene: a 185-kilodalton glycoprotein with tyrosine kinase activity. *Science* 1986; 232(4758): 1644–6
14. Shah S, Chen B. Testing for HER2 in Breast Cancer: A Continuing Evolution. *Patholog Res Int* 2010;2011: 903202
15. Yarden Y, Sliwkowski MX. Untangling the ErbB signalling network. *Nat Rev Mol Cell Biol* 2001; 2(2): 127–37
16. Moasser MM. The oncogene HER2: its signaling and transforming functions and its role in human cancer pathogenesis. *Oncogene* 2007; 26(45): 6469–87
17. Slamon DJ, Godolphin W, Jones LA, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science* 1989; 244(4905): 707–12
18. Slamon DJ, Clark GM, Wong SG, et al. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science*. 1987; 235(4785): 177–82
19. Slamon DJ, Leyland-Jones B, Shak S, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 2001; 344(11): 783–92
20. Sliwkowski MX, Lofgren JA, Lewis GD, et al. Nonclinical studies addressing the mechanism of action of trastuzumab (Herceptin). *Semin Oncol* 1999; 26(4 Suppl 12): 60–70
21. Romond EH, Perez EA, Bryant J, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005; 353(16): 1673–84
22. Wood ER, Truesdale AT, McDonald OB, et al. A unique structure for epidermal growth factor receptor bound to GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells. *Cancer Res* 2004; 64(18): 6652–9
23. Untch M, Lück HJ. Lapatinib – Member of a New Generation of ErbB-Targeting Drugs. *Breast Care (Basel)* 2010; 5(s1): 8–12
24. Nelson MH, Dolder CR. Lapatinib: a novel dual tyrosine kinase inhibitor with activity in solid tumors. *Ann Pharmacother* 2006; 40(2): 261–9
25. Cameron D, Casey M, Press M, et al. A phase III randomized comparison of lapatinib plus capecitabine versus capecitabine alone in women with advanced breast cancer that has progressed on trastuzumab: updated efficacy and biomarker analyses. *Breast Cancer Res Treat* 2008; 112(3): 533–43
26. National Cancer Institute [Internet]. Bethesda, MD: National Institutes of Health [updated 2014 January 22; cited 2014 December 21]. Available from: <http://www.cancer.gov/cancertopics/factsheet/Risk/BRCA>
27. Lo T, Pellegrini L, Venkitaraman AR, et al. Sequence fingerprints in BRCA2 and RAD51: implications for DNA repair and cancer. *DNA Repair (Amst)* 2003; 2(9): 1015–28
28. Risch, McLaughlin JR, Cole DE, et al. Population BRCA1 and BRCA2 mutation frequencies and cancer penetrances: a kin-cohort study in Ontario, Canada. *J Natl Cancer Inst* 2006; 98(23): 1694–706
29. El-Khamisy SF, Masutani M, Suzuki H, et al. A requirement for PARP-1 for the assembly or stability of XRCC1 nuclear foci at sites of oxidative DNA damage. *Nucleic Acids Res* 2003; 31(19): 5526–33
30. Fong PC, Boss DS, Yap TA, et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med* 2009; 361(2): 123–34
31. Rouleau M, Patel A, Hendzel MJ, et al. PARP inhibition: PARP1 and beyond. *Nat Rev Cancer* 2010; 10(4): 293–301
32. Zebisch A, Troppmair J. Back to the roots: the remarkable RAF oncogene story. *Cell Mol Life Sci* 2006; 63(11): 1314–30

33. Roskoski R Jr. RAF protein–serine/threonine kinases: structure and regulation. *Biochem Biophys Res Commun* 2010; 399(3): 313–7
34. Wellbrock C, Karasarides M, Marais R. The RAF proteins take centre stage. *Nat Rev Mol Cell Biol* 2004; 5(11): 875–85
35. Downward J. Targeting RAS signalling pathways in cancer therapy. *Nat Rev Cancer* 2003; 3(1): 11–22
36. Scharovsky OG, Rozados VR, Gervasoni SI, et al. Inhibition of ras oncogene: a novel approach to antineoplastic therapy. *J Biomed Sci* 2000; 7(4): 292–8
37. Brose MS, Volpe P, Feldman M, et al. BRAF and RAS mutations in human lung cancer and melanoma. *Cancer Res* 2002; 62(23): 6997–7000
38. Domingo E, Laiho P, Ollikainen M, et al. BRAF screening as a low–cost effective strategy for simplifying HNPCC genetic testing. *J Med Genet* 2004; 41(9): 664–8
39. Hollestelle A1, Elstrodt F, Nagel JH, et al. Schutte. Phosphatidylinositol–3–OH Kinase or RAS Pathway Mutations in Human Breast Cancer Cell Lines. *Mol Cancer Res* 2007; 5(2): 195–201
40. Gollob JA, Wilhelm S, Carter C, et al. Role of Raf kinase in cancer: therapeutic potential of targeting the Raf/MEK/ERK signal transduction pathway. *Semin Oncol* 2006; 33(4): 392–406
41. Wilhelm SM, Carter C, Tang L, et al. BAY 43–9006 exhibits broad spectrum oral antitumor activity and targets the RAF/MEK/ERK pathway and receptor tyrosine kinases involved in tumor progression and angiogenesis. *Cancer Res* 2004; 64(19): 7099–109
42. Tannapfel A, Sommerer F, Benicke M, et al. Mutations of the BRAF gene in cholangiocarcinoma but not in hepatocellular carcinoma. *Gut* 2003; 52(5): 706–12
43. Gradishar W, Kaklamani V, Prasad Sahoo T, et al. A Double–Blind, Randomized, Placebo–Controlled, Phase 2b Study Evaluating the Efficacy and Safety of Sorafenib (SOR) in Combination with Paclitaxel (PAC) as a First–Line Therapy in Patients (pts) with Locally Recurrent or Metastatic Breast Cancer (BC). *Cancer Res* 2009; 69(24 Supplement): 44