

Biological Activity and Phytochemical Study of *Scutellaria platystegia*

Seyedeh Neda Madani mousavi^{a, b}, Abbas Delazar^{a, b}, Hossein Nazemiyeh^b and Laleh Khodaie^{a*}

^aDrug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. ^bSchool of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Abstract

This study aimed to determine biological activity and phytochemical study of *Scutellaria platystegia* (family Labiatae). Methanolic (MeOH) extract of aerial parts of *S. platystegia* and SPE fractions of methanolic extract (specially 20% and 40% methanolic fractions), growing in East-Azarbaijan province of Iran were found to have radical scavenging activity by DPPH (2, 2-diphenyl -1- pycryl hydrazyl) assay. Dichloromethane (DCM) extract of this plant exhibited antimalarial activity by cell free method providing IC₅₀ at 1.1876 mg/mL. Crude extracts did not exhibit any toxicity assessed by brine shrimp lethality assay.

Phytochemical study of methanolic extract by using reverse phase HPLC method and NMR instrument for isolation and identification of pure compounds respectively, yielded 2-(4- hydroxy phenyl) ethyl-O-β-D- glucopyranoside from 10% and apigenin 7-O-glucoside, verbascoside and martynoside from 40% SPE fraction. Occurance of verbascoside and martynoside as biochemical markers appeared to be widespread in this genus.

Antioxidant and antimalarial activity of MeOH and DCM extracts, respectively, as well as no general toxicity of them could provide a basis for further *in-vitro* and *in-vivo* studies and clinical trials to develop new therapeutical alternatives.

Keywords: *Scutellaria platystegia*; Antimalaria; Antioxidant; DPPH; Labiatae.

Introduction

Scutellaria which belongs to Labiatae family, spread throughout the world, with approximately 350 currently recognized species (1-2). Plants of this genus have been traditionally used in China, Korea, and Japan as an agent for activating blood circulation, inducing diuresis and reducing oedema. Some other applications of these plants in folk medicine are due to their anti-inflammatory, antiviral, sedative and antioxidant effects (3). Modern pharmacologic researches on crude extracts and isolated compounds of the plants of this genus, confirmed multiple biological

activities, including anticonvulsant (2, 4), prolyl oligopeptidase inhibitory, hepatoprotective (3, 5), memory improvement (6-7) effects. In some *in-vitro* methods, phytochemicals of some species of *Scutellaria*, exhibited potent cytotoxic effects on some of the human tumour cell lines (8-10). Flavonoids (3, 11-12) and neoclerodan diterpenoids (1, 13-14) as well as iridoids (3, 15), phenyl alcohol glycosides and alkaloids (3, 16) have been isolated from several species of *Scutellaria*. Due to Remarkable and diverse biological activities of other species of *scutellaria* genus, this study aimed to evaluate biological activity and identify chemical composition of this plant. To our knowledge there have been no reports on biological activities and chemical composition of *S. platystegia*.

* Corresponding author:

E-mail: khodaiel89@gmail.com

Experimental

Plant material

Aerial parts of *Scutellaria platystegia* (family Labiatae) were gathered from Yam region of East Azarbaijan province of Iran in June 2009. A voucher specimen (Tbz-Fph-724) for this collection has been deposited in the Herbarium of pharmacy faculty, Tabriz University of Medical Sciences, Tabriz, Iran. 100 g of the air dried and grounded sample was extracted by the aid of Soxhlet apparatus using hexane, dichloromethane (DCM) and methanol (MeOH) respectively (1 L for each solvent). Obtained extracts were individually concentrated under vacuum in a rotary evaporator (Heidolph, Germany), yielding 1.67 g, 1.50 g and 24.50 g respectively.

Fractionation of methanolic extract

Fractionation of dried methanolic extract (2 g) was carried out using SPE (solid phase extraction) cartridge (Sep- pak ODS C18), eluting with methanol and water mixture (10:90, 20:80, 40:60, 60:40, 80:20 and 100:0). All SPE fractions were dried by rotary evaporator at a temperature not exceeding 50 °C. In order to increase the available SPE fractions, procedure was repeated 3 times yielding 1.421, 0.279, 0.655, 0.220, 0.162 and 0.190 g of each fraction respectively.

DPPH assay

Antioxidant activity of extracts and different SPE fractions was performed using DPPH assay. The basis of this experiment was Bleaching of purple coloured methanolic solution of 2, 2-diphenyl -1- pycryl hydrazyl (DPPH) (sigma). In order to obtain antioxidant activity, different sample solution series were prepared. 5 mL of each concentration of methanolic extract and SPE fractions were added to 5 mL of 0.004% methanolic solution of DPPH. After 30 min incubation of solutions at room temperature and bleaching of DPPH, absorption of samples was monitored at 517 nm against a blank. Inhibition of DPPH was calculated as RC_{50} that was extrapolated from dose-response curve. Tests were carried out in duplicate (17-18).

Antimalaria assay

The antimalaria potential of extracts was determined using cell free method which was described by Fitch *et al.* (19) with some modifications (20). Different concentrations of the n-hexan, DCM and methanolic extracts ranging from 0-2 mg/mL in 10% DMSO, were incubated in 300 µL of haematin which was freshly dissolved in 0.1 M NaOH, 10 mM oleic acid and 10 µM HCl. Afterwards 500 mM sodium acetate buffer, pH 5, was added to test tubes for adjusting reaction volume to 1000 µL. Positive control was chloroquine diphosphate in this test. The samples were incubated overnight at 37 °C with regular shaking. After incubation, samples were centrifuged at 14,000 × g, for 10 min, at 21 °C and the hemozoin pellet repeatedly washed with sonication (30 min, at 21 °C; FS100 bath sonicator; Decon Ultrasonics Ltd.) in 2.5% (w/v) SDS in phosphate buffered saline followed by a final wash in 0.1 M sodium bicarbonate, pH 9.0, until the supernatant became clear which usually happens after 3-5 washes. After the final wash, the supernatant was removed and the pellets were re-suspended in 1 mL of 0.1 M NaOH before determining the hemozoin content by measuring the absorbance at 400 nm (Beckmann DU640 spectrophotometer). Results were recorded as IC%, which is inhibition of heme crystallization compared to chloroquine as positive control, using the following formula: $IC\% = [(AB-AA)/AB] \times 100$, where AB and AA are absorbance of blank and test samples respectively. Final results which mean inhibition of hemozoin polymerization have been shown as IC50.

Brine shrimp lethality assay

This test is proposed as a preliminary and simple assay to study general toxicity of plant extracts. The eggs of *Artemia salina* purchased from Water Life, Middlesex, UK, were hatched in a flask containing 300 mL of artificial sea water aerated by the aid of an air pump. The flasks were kept in a 29-30 °C water bath and a bright light was left on. Afterwards, nauplii were hatched after 48 h. 1 mg/mL of n-hexan, DCM, MeOH were prepared by dissolving them in 5% DMSO. These solutions were serially diluted to obtain 7 concentrations, by the aid of aerated sea

water this experiment was carried out for twice. About ten nauplii were transferred in to each test tube. The number of alive nauplii were counted after 24 h. The control test tubes contained 5% DMSO, saline and podophyllotoxin (21-23).

Isolation of compounds

Preparative reversed phase HPLC with photodiode array detector was used for isolation of phytochemicals from 20 and 40% SPE fractions. Each fraction was analysed repeatedly by preparative reverse phase HPLC (Knauer, preparative pump 1800), equipped with a Reprosil 100 C18 (250 mm length, 20 mm *i.d.*, particle size 10 μm , Dr. Maisch, Germany) column. The mobile phase consisted of (A) methanol and (B) water. The following mobile phase program was used over 60 min to isolate glycosylated phenylethanoid (Figure 1) from the 10% SPE fraction: A initially changed to 10% in 15 min. then it changed to 20% in 50 min, maintained there for 10 min. A program over a run time of 50 min was applied for separation of 7-glucoapigenin (Figure 2), verbascoside (Figure 3) and martynoside (Figure 4) from the 40% SPE fraction: 17% A initially changed to 28% in 40 min. Then, it stayed there for 10 min. Photodiode Array Detector (PDA) was used to monitor the chromatogram, and the HPLC separation was carried out at room temperature. The flow rate was 8 mL/min and the injection volume was 1 mL. Structures of compounds were determined by H and C NMR (Bruker-spectrospin at 200MHz) as well as comparison with the literature data of respective compounds.

2-(4-hydroxy phenyl) ethyl-O- β -D-glucopyranoside (Figure 1); pale green solid; ^1H NMR (200 MHz, D_2O): Aglycone moiety: δ 7.04 (2 H, d, $J=7.9$ Hz, H2,6), 6.68 (2 H, d, $J=7.9$ Hz, H3,5), 3.8-4.1 (2 H, overlapped peak, H8), 2.7 (2 H, t, $J=6.9$ Hz, H7), Glucose moiety: δ 4.29 (1 H, d, $J=7.9$ Hz, H1'), 3.2-3.8 (signal patterns unclear due to overlapping, H2'-3'-4'-5'-6'), ^{13}C NMR (200 MHz, CD3OD): Aglycone moiety: δ 153.88(C4), 130.42(C1), 130.19 (C2,6), 115.28(C3,5), 73.02(C8), 34.22(C7).

Apigenin 7-O-glucoside (Figure 2); brown solid; ^1H NMR (200 MHz, CD3OD). Aglycone moiety: δ 7.90 (2H, d, $J=8.7$, H2',6'), 6.93 (2H, d, $J=8.7$, H3',5'), 6.85 (1H,d, $J=1.8$ Hz, H8), 6.63

(1H, S, H3), 6.46 (1H,d, $J=1.8$ Hz, H6). Glucose moiety: δ 4.65 (1 H, d, $J=7.8$ Hz, H1'), 3.2-3.8 (signal patterns unclear due to overlapping, H2'', H 3'', H 4'', H5''and H6'').

Verbascoside (Figure 3): brown solid; ^1H NMR (200 MHz, CD3OD): aglycone moiety: δ 7.6 (1H, d, $J=15.8$ Hz, H7'), 7.06(1H, d, $J=1.82$ Hz, H2'), 6.98(1H, dd, $J=8.1,1.8$ Hz, H6'), 6.78(1H, d, $J=8.1$ Hz, H5'), 6.69(1H, d, $J=1.8$ Hz, H2), 6.64(1H, d, $J=8.1$ Hz, H5), 6.58(1H, dd, $J=8.1,1.8$ Hz, H6), 6.25 (1H, d, $J=15.8$ Hz, H8'), 3.8-4.1 (2H, overlapped peak, H8), 2.79 (2H, t, $J=8$ Hz, H7), glucose moiety: δ 4.37 (1H, d, $J=7.8$ Hz, H1''), 3.2-3.8 (signal patterns unclear due to overlapping, H2''-3''-4''-5''-6''), rhamnose moiety: δ 5.19 (1H, d, $J=1.5$ Hz, H1'''), 3.2-3.8 (signal patterns unclear due to overlapping, H2'''-3''' -4''' -5'''), 1.085 (3H, d, $J=6.0$ Hz, H6'''). ^{13}C NMR (200 MHz, CD3OD) Aglycone moiety: δ 166.8(C9'), 148.4(C4'), 146.6(C7'), 145.4(C3'),), 143.2(C3), 130(C1), 126.2(C1'), 121.8(C6'), 119.8(C6), 115.7(C2), 115.7(C5'), 114.86(C5), 114.7(C4), 113.7 (C2'), 113.2 (C8'), 72.3(C8), 35.1(C7), Glucose moiety: δ 101.6(C1''), 80.2(C3''), 74.8(C2''), 74.6(C5''), 69(C4''), 60.9(C6''), Rhamnose moiety: 102.77(C1'''), 72.3(C4'''), 70.9(C2'''), 70.6(C3'''), 69.1(C5'''), 17(C6'''). Data were in agreement with the published data (24).

Martynoside (Figure 4); dark solid; ^1H NMR (200 MHz, CD3OD): aglycone moiety: δ 7.63 (1H, d, $J=15.8$ Hz, H7'), 7.16(1H, d, $J=1.8$ Hz, H2'), 7.03(1H, dd, $J=8.0,1.8$ Hz, H6'), 6.79(1H, d, $J=8.0$ Hz, H5), 6.79(1H, d, $J=8$ Hz, H5'), 6.69(1H, d, $J=1.8$ Hz, H2), 6.67(1H, dd, $J=8.0,1.8$ Hz, H6), 6.40 (1H, d, $J=15.8$ Hz, H8'), 3.8-4.1 (2H, overlapped peak, H8), 3.87 (3H, S, 3'-OMe), 3.75 (3H, S, 3-OMe), 2.8 (2H, t, $J=7.3$ Hz, H7), glucose moiety: δ 4.34 (1H, d, $J=7.8$ Hz, H1''), 3.2-3.8 (signal patterns unclear due to overlapping, H-2'', H-3'', H-4'', H-5'', H-6''), Rhamnose moiety: δ 5.18 (1H, d, $J=1.7$ Hz, H1'''), 3.2-3.8 (signal patterns unclear due to overlapping, H-2''', H-3''', H-4''', H-5'''), 1.085 (3H, d, $J=6.0$ Hz, H6'''). ^{13}C NMR (200 MHz, CD3OD); Aglycone moiety: δ 166.85 (C9'), 150.81 (C4'), 149.41 (C3'), 148.03 (C7'), 147.98 (C3), 147.37(C4), 132.93 (C1), 127.71 (C1'), 124.46 (C6'), 122.94 (C6), 117.19 (C5), 116.66 (C5'), 115.19 (C8'), 112.95 (C2), 111.91 (C2'),




Figure 1. 2-(4-hydroxy phenyl) ethyl-O- β -D- glucopyranoside.




Figure 2. Apigenin 7-O-glucoside.




Figure 3. Verbascoside.




Figure 4. Martynoside.

Table 1. Free radical scavenging activity of the extracts of *S. platystegia* by DPPH assay.

Extract	<i>n</i> -hexan	DCM	MeOH	Quercetin
RC50 (mg/m)	1.4283	0.7482	0.0342	0.0039

Table 2. Free radical scavenging activity of SPE methanolic fractions of *S. platystegia* by DPPH assay.

Fraction	10%	20%	40%	60%	80%	100%	Quercetin
RC50 (mg/mL)	0.0677	0.0085	0.0144	0.0314	0.2479	0.7321	0.0039

Table 3. Antimalarial activity of the extracts of *S. platystegia* by inhibition of Heme biocrystallisation assay

Extract	<i>n</i> -hexan	DCM	MeOH	Chloroquin
IC50 (mg/mL)	-	1.1876	16.9356	0.043

102.79 (C1'''), 101.58 (C1''), 81.65 (C3''), 76.24 (C2''), 76.04 (C5''), 73.87 (C4'''), 72.42 (C2'''), 72.35 (C8), 72.20 (C3'''), 70.74 (C4''), 70.31 (C5'''), 62.49 (C6''), 53.49 (O-CH3), 53.39 (O-CH3), 35.12 (C7), 17.03 (C6'''). Data were in agreement with the published data (25).

Results

DPPH assay was performed to determine Radical scavenging activity of extracts and SPE fractions of this plant. As it can be seen in Table 1, methanolic extract showed better antioxidant activity than other crude extracts. Table 2 demonstrated that among SPE fractions, 20% and 40% hydroalcoholic fractions were more potent antioxidants. Results of antimalarial activity of this plant, which was determined by

cell free method, exhibited in Table 3. DCM crude extract, providing IC50 at 1.1876 mg/mL showed potential of antimalarial activity.

Observation of radical scavenging activity from methanolic extract and SPE fractions encouraged us to study this plant phytochemically. Reverse-phase prep-HPLC analysis of SPE fractions of methanolic extract of aerial parts of *S. platystegia* (Labiatae) yielded 2-(4- hydroxy phenyl) ethyl-O-β-D- glucopyranoside from 10% ,one flavonoid (Apigenin 7-O-glucoside) and 2 phenyl ethanoid glycosides (verbascoside and martynoside) from 40% SPE fractions. In fact compounds 1-4 have been identified previously from other species of this genus (26-34), whereas this is the first report of these phytochemicals from *S. platystegia*. Distribution of compounds 1-4 has been demonstrated in Table 4.

Table 4. Distribution of compounds 1-4 within genus *Scutellaria*.

Scutellaria spp.	compounds			
	1	2	3	4
<i>S. baicalensis</i>	+	N.R.*	+	+
<i>S. baicalensis</i> Georgi	+	N.R.	N.R.	N.R.
<i>S. barbata</i>	N.R.	+	N.R.	N.R.
<i>S. immaculate</i>	N.R.	+	N.R.	N.R.
<i>S. pontifica</i>	N.R.	+	+	+
<i>S. ramosissima</i>	N.R.	+	+	+
<i>S. salviifolia</i>	N.R.	N.R.	+	+
<i>S. orientalis</i> sub. <i>pinnatifida</i>	N.R.	N.R.	+	+
<i>S. lateriflora</i>	N.R.	N.R.	+	+
<i>S. albida</i>	N.R.	N.R.	+	+
<i>S. salviifolia</i>	N.R.	N.R.	+	+
<i>S. galericulata</i>	N.R.	N.R.	+	+

* N.R.: not reported

Discussion

Recently, phytochemicals as bioactive components of plant extracts has received considerable attention. Antioxidants, the agents against oxidative stress-mediated disorders, with free radical scavenging activity, can prevent damages caused by various disorders (35-38). Antioxidant capacity of phenolic compounds isolated from plants (39-42) and Correlation of phenol content and antioxidant activity has been shown in different studies as well (43-44). According to some evidences, flavonoids and phenylethanoids as plant derived polyphenolic compounds act as free radical acceptors, and potent antioxidants (45-49). Plants of *Scutellaria* genus, used in traditional medicine for thousands years, with variety of confirmed pharmacological effects in modern researches, has been shown free radical scavenging and antioxidant activities due to existence of different phenolic compounds such as flavonoids and phenylethanoids (50-57). Results of this study demonstrated that verbascoside, martynoside and apigenin as antioxidant compounds (45, 58-62), which existed in methanolic extract and isolated from 40% fraction were responsible for good radical scavenging activity of this methanolic fraction (0.0342 mg/mL). Further investigations will reveal other phytochemicals responsible for radical scavenging activity of 20% and 60% fractions. Identification of apigenin, martynoside and verbascoside as anti-inflammatory and antioxidant constituents from methanolic extract, is in agreement with traditional usage of this plant as an anti-inflammatory agent (63-66) and confirm its use in folk medicine.

Malaria, a malignancy with worldwide spread, results in loss of lives each year. Resistance of deadly forms of malaria parasites to anti-malarial drugs highlights needs for new antimalarial drugs (67). Anti-fever herbal plants used in traditional medicine might contain some antimalarial phytochemicals which could lead to development of new drugs for treatment of this mortal malignancy (68). Diverse nature of Iran possesses medicinal plants which could be alternative choice for treatment of malaria. In previous works, traditionally used febrifuge

plants to treat fever as a symptom of malaria, have been selected for antimalarial studies (69-71). Since several species of *Scutellaria* have been used as febrifuge in folk medicine, extracts of *S. platystegia* were subjected to *in-vitro* antimalarial test (57, 72-73) and interestingly DCM extract of this plant demonstrated antimalarial activity. Further investigations are needed to identify and purify compounds responsible for antimalarial activity of this plant.

Brine shrimp lethality assay, a suitable, simple, rapid procedure with low cost (74-75), was chosen to determine general toxicity of plant extracts. None of the plant extracts exhibited any toxicity at highest test concentrations (1 mg/mL).

Biological activity of methanolic extract conducted us to phytochemical study of this plant. 2-(4-hydroxy phenyl) ethyl-O- β -D-glucopyranoside (from 10% SPE fraction), Apigenin-7-O-glucoside, verbascoside and martynoside (from 40% SPE fraction) were isolated and identified by UV and NMR analysis. All spectroscopic data were in agreement with respective published data. Identified components have been reported from other species of *Scutellaria* genus, while distribution of verbascoside and martynoside appears to be widespread (Table 4). So, it is concluded that these two phytochemicals could be determined as chemical biomarkers in this genus. To our knowledge, this is the first report on the antioxidant and antimalarial activity as well as occurrence of compounds 1-4 within this species.

Conclusion

It can be concluded that antioxidant and antimalarial activity of MeOH and DCM extracts, respectively, as well as no general toxicity of them, could provide a basis for further *in-vitro* and *in-vivo* studies and clinical trials to develop new therapeutical alternatives. More over the results of present study show that it is worth to do further phytochemical studies on Iranian *Scutellaria* species and to isolate compounds responsible for antioxidant and antimalarial activities.

Acknowledgments

We are grateful to Dr. Asnaashari and Mrs Bamdad for their valuable contributions to this study.

References

- (1) Dai SJ, Qu GW, Yu QY, Zhang DW and Li GS. New neo-clerodane diterpenoids from *Scutellaria barbata* with cytotoxic activities. *Fitoterapia* (2010) 81: 737-741.
- (2) Zhang Z, Lian XY, Li S and Stringer JL. Characterization of chemical ingredients and anticonvulsant activity of American skullcap (*Scutellaria lateriflora*). *Phytomed.* (2009) 16: 485-93.
- (3) Shang X, He X, Li M, Zhang R, Fan P and Zhang Q. The genus *Scutellaria* an ethnopharmacological and phytochemical review. *J. Ethnopharmacol.* (2010) 128: 279-313.
- (4) Park HG, Yoon SY, Choi JY, Lee GS, Choi JH and Shin CY. Anticonvulsant effect of wogonin isolated from *Scutellaria baicalensis*. *Eur. J. Pharmacol.* (2007) 574: 112-119.
- (5) De BJB, Quiney B, Walter PB, Thomas C, Hodgson K and Murch SJ. Protection against aflatoxin-B1-induced liver mutagenesis by *Scutellaria baicalensis*. *Mutat. Res.* (2005) 578: 15-22.
- (6) Shang Y, Cheng J, Qi J and Miao H. *Scutellaria* flavonoid reduced memory dysfunction and neuronal injury caused by permanent global ischemia in rats. *Pharmacol. Biochem. Behav.* (2005) 82: 67-73.
- (7) Heo H, Shin Y, Cho W, Choi Y, Kim H and Kwon YK. Memory improvement in ibotenic acid induced model rats by extracts of *Scutellaria baicalensis*. *J. Ethnopharmacol.* (2009) 122: 20-27.
- (8) Cha YY, Lee EO, Lee HJ, Park YD, Ko SG and Kim DH. Methylene chloride fraction of *Scutellaria barbata* induces apoptosis in human U937 leukemia cells via the mitochondrial signaling pathway. *Clin. Chim. Acta.* (2004) 348: 41-48.
- (9) Ozmen A, Madlener S, Bauer S, Krasteva S, Vonach C and Giessrigl B. *In-vitro* anti-leukemic activity of the ethno-pharmacological plant *Scutellaria orientalis* ssp. *carica* endemic to western Turkey. *Phytomed.* (2010) 17: 55-62.
- (10) Kumagai T, Muller CI, Desmond JC, Imai Y, Heber D and Koeffler HP. *Scutellaria baicalensis*, a herbal medicine: anti-proliferative and apoptotic activity against acute lymphocytic leukemia, lymphoma and myeloma cell lines. *Leuk. Res.* (2007) 31: 523-530.
- (11) Liu G, Ma J, Chen Y, Tian Q, Shen Y and Wang X. Investigation of flavonoid profile of *Scutellaria baicalensis* Georgi by high performance liquid chromatography with diode array detection and electrospray ion trap mass spectrometry. *J. Chromatogr. A* (2009) 1216: 4809-4814.
- (12) Sato Y, Suzaki S, Nishikawa T, Kihara M, Shibata H and Higuti T. Phytochemical flavones isolated from *Scutellaria barbata* and antibacterial activity against methicillin-resistant *Staphylococcus aureus*. *J. Ethnopharmacol.* (2000) 72: 483-488.
- (13) Miyaichi Y, Kizu H, Yamaguchi Y and Tomimori T. Studies on the constituents of *Scutellaria* species. XV. On the diterpenoid constituents of the leaves of *Scutellaria alpina* L. *Yakugaku Zasshi.* (1994) 114: 264-271.
- (14) Ezer N, Akcos Y and Rodriguez B. Neo-clerodane diterpenoids from *Scutellaria orientalis* subsp. *sintenisii*. *Phytochem.* (1998) 49: 1825-1827.
- (15) Gousiadou C, Karioti A, Heilmann J and Skaltsa H. Iridoids from *Scutellaria albida* ssp. *albida*. *Phytochem.* (2007) 68: 1799-1804.
- (16) Dai SJ, Liang DD, Ren Y, Liu K and Shen L. New neo-clerodane diterpenoid alkaloids from *Scutellaria barbata* with cytotoxic activities. *Chem. Pharm. Bull.* (2008) 56: 207-209.
- (17) Moein S, Moein M, Khoshnoud MJ and Kalanteri T. *In-vitro* antioxidant properties evaluation of 10 Iranian medicinal plants by different methods. *Iran. Red. Crescent Med. J.* (2012) 14: 771-775.
- (18) Kicel A and Wolbis M. Phenolic content and DPPH radical scavenging activity of the flowers and leaves of *Trifolium repens*. *Nat. Prod. Commun.* (2013) 8: 99-102.
- (19) Fitch CD. Involvement of heme in the antimalarial action of chloroquine. *Trans. Am. Clin. Climatol. Assoc.* (1998) 109: 97-105.
- (20) Tripathi AK, Gupta A, Garg SK and Tekwani BL. *In-vitro* beta-hematin formation assays with plasma of mice infected with *Plasmodium yoelii* and other parasite preparations: comparative inhibition with quinoline and endoperoxide antimalarials. *Life Sci.* (2001) 69: 2725-2733.
- (21) Meyer B, Ferrigni N, Putnam J, Jacobsen L, Nichols D and McLaughlin J. Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medica.* (2007) 45: 31-34.
- (22) Carballo JL, Hernández-Inda ZL, Pérez P and García-Grávalos MD. A comparison between two brine shrimp assays to detect *in-vitro* cytotoxicity in marine natural products. *BMC Biotechnol.* (2002) 2: 17.
- (23) Hasan MS, Ahmed MI, Mondal S, Uddin SJ, Masud M and Sadhu S. Antioxidant, antinociceptive activity and general toxicity study of *Dendrophthoe falcata* and isolation of quercitrin as the major component. *Int. J. Orient. Pharm. Exp. Med.* (2006) 6: 355-360.
- (24) Olivier D, Shikanga E, Combrinck S, Krause R, Regnier T and Dlamini T. Phenylethanoid glycosides from *Lippia javanica*. *S. Afr. J. Bot.* (2010) 76: 58-63.
- (25) Khodaie L, Delazar A, Lotfipour F, Nazemiyeh H, Asnaashari S and Moghadam SB. Phytochemistry and bioactivity of *Pedicularis sibthorpii* growing in Iran. *Rev. Bras. Farmacogn.* (2012) 22: 1268-1275.
- (26) Liu YX, Liu ZG, Su L, Yang RP, Hao DF and Pei YH. Chemical constituents from *Scutellaria baicalensis*

- Georgi. *Chinese J. Med. Chem.* (2009) 1: 14.
- (27) Sato Y, Suzaki S, Nishikawa T, Kihara M, Shibata H and Higuti T. Phytochemical flavones isolated from *Scutellaria barbata* and antibacterial activity against methicillin-resistant *Staphylococcus aureus*. *J. Ethnopharmacol.* (2000) 72: 483-488.
- (28) Çaliş I, Ersöz T, Saracoglu I and Sticher O. Scalbidoside and albidoside, two iridoid glycosides from *Scutellaria albida* subsp. *Colchica*. *Phytochem.* (1993) 32: 1213-127.
- (29) Yuldashev M. Flavonoids of the aerial part of *Scutellaria immaculate*. *Chem. Nat. Compd.* (2001) 37: 428-430.
- (30) Simmonds MS. The search for plant-derived compounds with antifeedant activity. *Adv. Phytomed.* (2006) 3: 291-324.
- (31) Mamadalieva NZ, Herrmann F, El-Readi MZ, Tahrani A, Hamoud R and Egamberdieva DR. Flavonoids in *Scutellaria immaculata* and *S. ramosissima* (Lamiaceae) and their biological activity. *J. Pharm. Pharmacol.* (2011) 63: 1346-1357.
- (32) Saracoglu I, Inoue M, Calis I and Ogihara Y. Studies on constituents with cytotoxic and cytostatic activity of two Turkish medicinal plants *Phlomis armeniaca* and *Scutellaria salviifolia*. *Biol. Pharm. Bull.* (1995) 18: 1396.
- (33) Çaliş İ, Saracoglu İ, Başaran AA and Sticher O. Two phenethyl alcohol glycosides from *Scutellaria orientalis* subsp. *Pinnatifida*. *Phytochem.* (1993) 32: 1621-1623.
- (34) Zhang Z, Lian XY, Li S and Stringer JL. Characterization of chemical ingredients and anticonvulsant activity of American skullcap (*Scutellaria lateriflora*). *Phytomed.* (2009) 16: 485-493.
- (35) Ratnam DV, Ankola D, Bhardwaj V, Sahana DK and Kumar M. Role of antioxidants in prophylaxis and therapy: A pharmaceutical perspective. *J. Control Release* (2006) 113: 189-207.
- (36) Diplock AT. Antioxidant nutrients and disease prevention: an overview. *Am. J. Clin. Nutr.* (1991) 53: 189.
- (37) Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O and Lee JH. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J. Am. Coll. Nutr.* (2003) 22: 18-35.
- (38) Zafra-Stone S, Yasmin T, Bagchi M, Chatterjee A, Vinson JA and Bagchi D. Berry anthocyanins as novel antioxidants in human health and disease prevention. *Mol. Nutr. food Res.* (2007) 51: 675-683.
- (39) Soobrattee MA, Neergheen VS, Luximon-Ramma A, Aruoma OI and Bahorun T. Phenolics as potential antioxidant therapeutic agents: mechanism and actions. *Mutat. Res.* (2005) 579: 200.
- (40) Rice-Evans C, Miller N and Paganga G. Antioxidant properties of phenolic compounds. *Trends Plant Sci.* (1997) 2: 152-159.
- (41) Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K and Kujala TS. Antioxidant activity of plant extracts containing phenolic compounds. *J. Agr. Food Chem.* (1999) 47: 3954-3962.
- (42) Shahidi F, Janitha P and Wanasundara P. Phenolic antioxidants. *Crit. Rev. Food Sci. Nutr.* (1992) 32: 67-103.
- (43) Lopez-Velez M, Martinez-Martinez F and Valle-Ribes CD. The study of phenolic compounds as natural antioxidants in wine. *Crit. Rev. Food Sci.* (2003) 43: 233-244.
- (44) Turkmen N, Sari F and Velioglu YS. Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods. *Food Chem.* (2006) 99: 835-841.
- (45) Rice-evans CA, Miller NJ, Bolwell PG, Bramley PM and Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Res.* (1995) 22: 375-383.
- (46) Ordonez A, Gomez J and Vattuone M. Antioxidant activities of *Sechium edule* (Jacq.) Swartz extracts. *Food Chem.* (2006) 97: 452-458.
- (47) Lee JY, Yoon JW, Kim CT and Lim ST. Antioxidant activity of phenylpropanoid esters isolated and identified from *Platycodon grandiflorum* A. DC. *Phytochem.* (2004) 65: 3033-3039.
- (48) Zhang L, Liao CC, Huang HC, Shen YC, Yang LM and Kuo YH. Antioxidant phenylpropanoid glycosides from *Smilax bracteata*. *Phytochem.* (2008) 69: 1398-1404.
- (49) Chang CL, Zhang LJ, Chen RY, Kuo LMY, Huang JP and Huang HC. Antioxidant and anti-inflammatory phenylpropanoid derivatives from *Calamus quiqueseinervius*. *J. Nat. Prod.* (2010) 73: 1482-1488.
- (50) Huang WH, Lee AR and Yang CH. Antioxidative and anti-inflammatory activities of polyhydroxyflavonoids of *Scutellaria baicalensis* Georgi. *Biosci. Biotech. Biochem.* (2006) 70: 2371-2380.
- (51) Cole IB, Cao J, Alan AR, Saxena PK and Murch SJ. Comparisons of *Scutellaria baicalensis*, *Scutellaria lateriflora* and *Scutellaria racemosa*: genome size, antioxidant potential and phytochemistry. *Planta Medica.* (2008) 74: 474-81.
- (52) Gao Z, Huang K, Yang X and Xu H. Free radical scavenging and antioxidant activities of flavonoids extracted from the radix of *Scutellaria baicalensis* Georgi. *BBA- Gen Subjects* (1999) 1472: 643-650.
- (53) Gabrielska J, Oszmiański J, Zylka R and Komorowska M. Antioxidant activity of flavones from *Scutellaria baicalensis* in lecithin liposomes. *J. Biosci.* (1997) 52: 817.
- (54) Su YL, Leung LK, Bi YR, Huang Y and Chen ZY. Antioxidant activity of flavonoids isolated from *Scutellaria rehderiana*. *J. Am. Oil Chem. Soc.* (2000) 77: 807-813.
- (55) Shieh DE, Liu LT and Lin CC. Antioxidant and free radical scavenging effects of baicalein, baicalin and wogonin. *Anticancer Res.* (2000) 20: 2861.
- (56) Ersoz T, Tasdemir D, Calis I and Ireland C. Phenylethanoid glycosides from *Scutellaria galericulata*. *Turk J. Chem.* (2002) 26: 465-472.

- (57) Shang X, He X, He X, Li M, Zhang R and Fan P. The genus *Scutellaria* an ethnopharmacological and phytochemical review. *J. Ethnopharmacol.* (2010) 128: 279-313.
- (58) Benavente-Garcia O, Castillo J, Lorente J, Ortuno A and Del Rio J. Antioxidant activity of phenolics extracted from *Olea europaea* L. leaves. *Food Chem.* (2000) 68: 457-462.
- (59) Vertuani S, Beghelli E, Scalambra E, Malisardi G, Copetti S and Toso RD. Activity and stability studies of verbascoside, a novel antioxidant, in dermo-cosmetic and pharmaceutical topical formulations. *Molecules* (2011) 16: 7068-7080.
- (60) Wang P, Kang J, Zheng R, Yang Z, Lu J and Gao J. Scavenging effects of phenylpropanoid glycosides from *Pedicularis* on superoxide anion and hydroxyl radical by the Spin trapping method. *Biochem. Pharmacol.* (1996) 51: 687-691.
- (61) Papoutsis Z, Kassi E, Mitakou S, Aligiannis N, Tsiapara A and Chrousos GP. Acteoside and martynoside exhibit estrogenic/antiestrogenic properties. *J. Steroid Biochem. Mol. Biol.* (2006) 98: 63-71.
- (62) Lu Y and Yeap FL. Antioxidant activities of polyphenols from sage (*Salvia officinalis*). *Food Chem.* (2001) 75: 197-202.
- (63) Penido C, Costa KA, Futuro DO, Paiva SR, Kaplan MAC and Figueiredo MR. Anti-inflammatory and anti-ulcerogenic properties of *Stachytarpheta cayennensis* (LC Rich) Vahl. *J. Ethnopharmacol.* (2006) 104: 225-233.
- (64) Kanchanapoom T, Kasai R, Picheansoonthon C and Yamasaki K. Megastigmane, aliphatic alcohol and benzoxazinoid glycosides from *Acanthus ebracteatus*. *Phytochem.* (2001) 58: 811-817.
- (65) Vo TNN, Phi LT, Lam TP, Lawrence MV, Phung NN and Kim PP. Lignans and Triterpenes from the Root of *Pseuderanthemum carruthersii* var. *atropurpureum*. *Chem. Pharm. Bull.* (2012) 60: 1125-1133.
- (66) Kim HPS, Kun HC, Hyeun WK and Sam S. Anti-inflammatory plant flavonoids and cellular action mechanisms. *J. Pharm. Sci.* (2004) 96: 229-245.
- (67) Azas N, Laurencin N, Delmas F, Di Giorgio C, Gasquet M and Laget M. Synergistic *in-vitro* antimalarial activity of plant extracts used as traditional herbal remedies in Mali. *Parasitol. Res.* (2002) 88: 165-171.
- (68) Najib Nik A Rahman N, Furuta T, Takane K and Ali Mohd M. Antimalarial activity of extracts of Malaysian medicinal plants. *J. Ethnopharmacol.* (1999) 64: 249-54.
- (69) Ancolio C, Azas N, Mahiou V, Ollivier E, Di Giorgio C and Keita A. Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome. *Phytother. Res.* (2002) 16: 646-649.
- (70) Addae-Kyereme J, Croft SL, Kendrick H and Wright CW. Antiplasmodial activities of some Ghanaian plants traditionally used for fever/malaria treatment and of some alkaloids isolated from *Pleiocarpa mutica*; *in-vivo* antimalarial activity of pleiocarpine. *J. Ethnopharmacol.* (2001) 76: 99-103.
- (71) Koch A, Tamez P, Pezzuto J and Soejarto D. Evaluation of plants used for antimalarial treatment by the Maasai of Kenya. *J. Ethnopharmacol.* (2005) 101: 95-99.
- (72) Lazari D, Gabrieli C, Papi R, Tsoleridis K and Kyriakidis D. Biological activities of iridoids from *Scutellaria rupestris* ssp. *adenotricha*. *Planta Medica* (2008) 74: 145.
- (73) Skaltsa HD, Lazari DM, Kyriazopoulos P, Golegou S, Triantaphyllidis S and Sokovic M. Composition and antimicrobial activity of the essential oils of *Scutellaria sieberia* Benth. and *Scutellaria rupestris* Boiss. et Heldr. ssp. *adenotricha* (Boiss. et Heldr.) Greuter et Burdet from Greece. *J. Essent. Oil Res.* (2005) 17: 232-235.
- (74) Movahhedini N, Barar J, Fathi FA, Barzegari A and Nazemiyeh H. Phytochemistry and biologic activities of *Caulerpa peltata* native to oman sea. *Iran. J. Pharm. Res.* (2014) 13: 515-21.
- (75) Firuzi O, Miri R, Asadollahi M, Eslami S and jasbi AR. Cytotoxic, Antioxidant and antimicrobial activities and phenolic contents of eleven *salvia* species from Iran. *Iran. J. Pharm. Res.* (2013) 12: 801-810.

This article is available online at <http://www.ijpr.ir>

**Search full text articles?
Visit <http://www.ijpr.ir>
or
[http:// ijpr.sbm.ac.ir](http://ijpr.sbm.ac.ir)**