

# Analysis of *Plasmodium vivax* Merozoite Surface Protein-1 Gene Sequences

MR Razavi<sup>1</sup>, \*SR Naddaf<sup>1</sup>, M Assmar<sup>1</sup>, A Raesi<sup>2</sup>, Sh Arshi<sup>3</sup>

<sup>1</sup>Dept. of Parasitology, Pasteur Institute of Iran, Tehran, Iran

<sup>2</sup>National Program of Malaria Control DCD/ MOH&ME, No. 68, Iranshahr Ave., Tehran, Iran

<sup>3</sup>Infectious Disease Research Center, Ardebil University of Medical Sciences, Iran

(Received 13 Jan 2006; accepted 20 Aug 2006)

---

---

## Abstract

The emergence of *Plasmodium vivax* in Dashte Moghan in northwestern Iran has become a major concern for Iranian's health officials. Knowledge of genetic make up of the *P. vivax* populations in this area would give us an insight into the origin of the prevalent infections and the possible routes they are introduced. This paper reports the analysis of a variable region between the two interspecies conserved blocks (ICBs) of 5 and 6 of *MSP-1* gene in 18 isolates from Dashte Moghan. The results revealed that all the 18 isolates were similar to an Azari Belem-like type with 21 glutamine (Q) in the repeated residues. Our results may give a clue that the resurgent malaria in Dashteh Moghan might have primarily been introduced from Azerbaijan. However, much more molecular and epidemiological evidence are needed to confirm this hypothesis.

**Keywords:** Malaria, *Plasmodium vivax*, Merozoite surface antigen (MSP-1), Belem allele (Bel), Iran

---

---

## Introduction

*Plasmodium vivax* was the most prevalent human malaria parasite throughout Iran in the past and accounted for the majority of 5 million cases reported annually in then 13 million populated Iran (Jalili Moslem GH, 1952, unpublished). With the beginning of malaria campaign at late 1950s this species was eradicated from the most northern parts of the country and confined to some areas in south and southeastern provinces, where it often co-occur with *Plasmodium falciparum*. The reemerge of *P. vivax* malaria In Dashte Moghan in northwestern Iran, which has been reported malaria-free since early 1960s (1) became a major concern for the public health officials. This area borders with Azerbaijan Republic, a newly independent state in central Asia with some experiences of *P. vivax* malaria outbreaks since early 1990s (2). Knowledge of the population genetic structure of the malaria parasite in Dashte Moghan would allow us to gain

an insight into origin of the parasite and update our understanding of epidemiology of the disease, which would finally lead us to adoption of proper control measures. The growing evidences on capability of *MSP-1* in *P. falciparum* (*PfMSP-1*) as a useful genetic maker for discriminating variation within and among populations of this malaria parasite (3- 5) prompted many researches to use the same tool for studying genetic structure of *P. vivax* populations. *P. vivax* major surface protein 1 (*PvMSP-1*) like *PfMSP-1* exhibits antigenic polymorphism (6). Complete gene sequence of *PvMSP-1*, a 200 kDa polymorphic glycoprotein, was characterized in two isolated strains: Brazilian Belem (Bel) and Salvador (Sal1) (7, 8). Later, analysis of specific gene regions derived from parasite isolates from Srilanka (9) Thailand (10), and Colombia (11) revealed new sequence types of *MSP-1* polymorphic region, most of them resulting from recombination between the sequence of the two original Belem

and Sal 1 strains (11). *PvMSP-1* gene consists of seven interallelic conserved blocks flanked by six variable blocks. The variable blocks, particularly the one between interspecies conserved blocks 5 and 6 show extensive variation in repeats and nonrepeats unique sequences among *P. vivax* isolates (10). In particular, the Belem strain can easily be characterized by a variable poly-glutamine (poly Q) region absent in Sal 1 strain.

This study reports the genetic make up of *P. Vivax* population in Dashte Moghan by analyzing the variable block between the ICB5 and ICB6 of *MSP-1* gene.

## Materials and Methods

**Study area and sample collection** Blood samples were collected from 18 febrile patients, who were seeking treatment from public health centers after their verbal consent. All the blood samples were shown to harbor *P. vivax* by both microscopy and rDNA-Polymerase chain reaction. After identification whole blood samples were frozen at -70 °C until used.

**DNA extraction** 300 µl of whole blood was mixed with three volumes of cold double distilled water (ddH<sub>2</sub>O) and kept on ice for 10 min. The parasite and unlysed erythrocytes were recovered by centrifugation at 8000 rpm for 7 min. The supernatant was discarded and the pellet was resuspended in 1 ml cold PBS containing saponin to a final concentration of 0.05%. Immediately after the lysis was observed, parasites and white blood cells were recovered by centrifugation as above. DNA extraction from the pellet was accomplished using a DNA isolation kit (Amersham Bioscience, USA) according to the manufacturer's instructions. The recovered DNA was resuspended in 50 µl of ddH<sub>2</sub>O.

### Polymerase chain reaction and sequencing

The highly polymorphic region with the flanking ICB5 and ICB6 of *MSP-1* gene was amplified using Pv1 and Pv2 primers and PCR reaction conditions of Putaporntip et al., (10). These primers would allow amplification of a nucleo-

tide region corresponding to basepairs 1813 to 2827 in Sal1 allele. 100 µl of each amplified sample was run on a 0.8% agarose gel and fragment of interest was cut and purified using a gel band purification kit (Amersham Pharmacia Biotech, Inc) according to manufacturer's instructions. The concentration of recovered DNAs was estimated by running them on an agarose gel along with a known concentration of DNA molecular weight marker (MWM III, Roche). Sequencing of fragments was performed in both directions using an automated sequencer (SeqLab Laboratory, Germany).

DNA alignment was done using a Clustal X (version 1.8) multiple sequence alignment program.

## Results

PCR amplification of the variable region and the flanking ICB5 and ICB6 of the *PvMSP-1* gene revealed a limited size variation among isolates ranging from 862 to 894.

The sequencing result of DNA and corresponding amino acids revealed a distinct Belem-like type in all 18 isolates. There was a consensus deletion of 6 nucleotides (CAACAA) in all isolates, corresponding to nucleotides 2334-2339 in Belem allele leading to two lesser number of Q in poly-Q repeats i.e. 21Q as compared to 23 in Belem allele. There were also some synonymous substitutions in the polymorphic region which could be classified into two types. Both types shared a substitution of *g* for *a* at position 2327, but type 2 had three more substitutions including *t* for *c* at position 2736, *a* for *g* at position 2738 and *c* for *t* at position 2747 (Fig. 1).



## Discussion

Dashte Moghan has been reported malaria-free since the implementation of an extensive national malaria eradication program throughout Iran beginning in 1958 (1). However this disease has reemerged in this area since 1994 with more than 3126 reported patients diagnosed with *P. vivax* up to 2004. Analysis of genetic make up of *P. vivax* populations in this area and their comparison with already published data available in Gebank from Iran and neighboring countries would give us some clues on the origin of this resurgent malaria. In the neighboring country of Azerbaijan, analysis of the highly variable region between ICB5 and ICB6 of this gene showed that both Belem-like and Sal1-like types together with some recombinant types were prevalent in the five districts of the country (2). We studied the same fragment in 18 isolates collected from Dashte Moghan in Ardebil Province of Iran. This area borders with Azerbaijan from north and northeastern parts, where many people move between two countries without government control. Our study revealed two types similar to an Azeri Belem-like type in all the 18 isolates, with the same poly-Q region of 21 repeats as compared to 23 in Belem allele. It is noteworthy that all the Belem-like types from Azerbaijan had also 21Q in the poly-Q repeats while the poly-Q repeats in the five Azari recombinant types ranged between 12-18 Q. Genetic analysis of 16 plasmodial isolates from southeastern Iran using the same region of *P. vivax* MSP-1 revealed 12 Belem-like types with number of poly-Q repeats ranging from 17-23. The only isolate with 21 Q was proved a recombinant type (13).

Our results may give a clue that the resurgent malaria in Dashteh Moghan might have primarily been introduced from Azerbaijan. However, much more molecular and epidemiological studies are needed to confirm this hypothesis.

## Acknowledgments

We acknowledge the colleagues of the Center

for Disease Control and Management (Malaria Division) in Iran for their assistance in collecting blood samples. This project was founded by Center for Disease Control, ministry of Health and Medical Education, Iran

## References

1. Manouchehri AV, Zaim M, Emadi AM (1992). A review of malaria in Iran 1975-1990. *J Am Mosq Control Assoc*, 8: 381-5.
2. Leclerc MC, Menegon M, Cligny A, Noyer JL, Mammadov S, Aliyev N et al. (2004). Genetic diversity of *Plasmodium vivax* isolates from Azerbaijan: <http://www.malariajournal.com/content/3/1/40>
3. Tanabe K, Mackay M, Goman M, Scaife JG (1987). Allelic dimorphism in a surface antigen gene of malaria parasite *Plasmodium falciparum*, *J Mol Biol*, 195: 273-87.
4. Jongwutiwes S, Tanabe K, Nakazawa S, Yanagi T, Kanbara H (1992). Sequence variation in the tripeptide repeats and T cell epitopes in p190 (MSA-1) of *Plasmodium falciparum* from field isolates. *Mol Biochem Parasitol*, 51: 81-90.
5. Kerr PJ, Ranford-Cartwright LC, Walliker D (1994). Proof of intragenic recombination in *Plasmodium falciparum*. *Mol Biochem Parasitol*, 66: 241-48.
6. Udagama PV, Gamage-Mendis AC, David PH, Peiris JSM, Perera, KLRL, Mendis KN, Carter R (1990). Genetic complexity of *Plasmodium vivax* parasites in individual human infections analyzed with monoclonal antibodies against variant epitopes on a single parasite protein. *Am J Trop Med Hyg*, 42: 104-10.
7. del Portillo HA, Longacre S, Khouri E, David PH (1991). Primary structure of the merozoite surface antigen 1 of *Plasmodium vivax* reveals a sequence conserved between different *Plasmodium* species. *Proc Natl Aca Sci USA*, 88: 4030-52.

8. Gibson HL, Tucker JE, Kaslow DC, Krettli AU, Collins WE, Kiefer MC, Bathurst IC, Barr PJ (1992). Structure and expression of the gene for Pv200, a major blood stage surface antigen of *Plasmodium vivax*. *Mol Biochem Parasitol*, 50: 325-34.
9. Premawansa S, Snewin VA, Khouri E, Mendis K, David PH (1993). *Plasmodium vivax*: recombination between potential allelic types of the merozoite surface protein MSP-1 in parasites isolated from patients. *Exp Parasitol*, 76: 192-99.
10. Putaporntip C H, Jongwutiwes S, Tanabe K, Thaithong S (1997) Interallelic recombination in the merozoite surface protein 1 (*MSP-1*) gene of *Plasmodium vivax* from Thai isolates. *Mol Bioch Parasitol*, 84: 49-56.
11. Mancilla L, Levitus G, Kirchgatter K, Mertens F, Herrera S, del portillo H (1994). *Plasmodium vivax*: Dimorphic DNA sequences from the *MSP-1* gene code for regions that are immunogenic in natural infections. *Exp Parasitol*, 79: 148-58.
12. raig AA, Kain KC (1996). Molecular analysis of strains of *Plasmodium vivax* from paired primary and relapse infection. *J Infect Dis*, 174: 812-820.
13. Zakeri S, Djadid ND, Zeinali S (2003). Sequence heterogeneity of the merozoite surface protein 1 gene (*MSP-1*) of *Plasmodium vivax* wild isolates in Southeastern Iran. *Acta Trop*, 88: 91-97.