

Clinico-epidemiological study on metal-induced contact dermatitis from North India

Khushman Singh*, Sunil Kumar Gupta*, Veenu Gupta**, Sukhmani Kaur Brar*

* Department of Dermatology, Venereology & Leprology, Dayanand Medical College and Hospital, Ludhiana

** Department of Microbiology, Dayanand Medical College and Hospital, Ludhiana

Abstract

Objective To determine the prevalence of metal sensitization and different clinical patterns in suspected patients of metal induced contact dermatitis.

Methods This was an epidemiological study during which a total of 100 patients suspected and provisionally diagnosed as having contact dermatitis to metals were subjected to a patch test with Indian Standard Battery developed by CODFI (Contact and Occupational Dermatoses Forum of India). Results were read after 48hrs of application and a second reading, if required, was taken after 72-96hrs.

Results Most of the patients in this series presented with hand eczema (40%), which was followed by dermatitis localized to site of contact with metal (20%) and facial dermatitis including eyelid dermatitis (15%). Twenty-seven patients showed positive reactions to metal allergens. Cross-sensitivity was low and three patients or 11.11% showed sensitization to two metal allergens. No patient showed sensitization to all the three metals.

Conclusion Nickel was the commonest metal allergen identified and it was relevant in 100% of the cases. It showed a strong female preponderance. Patch testing is hence a helpful diagnostic aid in identifying the agents responsible for contact dermatitis and a sincere effort should be made to determine clinical relevance of the test results in every case.

Keywords

Contact dermatitis, metal, clinico-epidemiological study.

Introduction

Contact dermatitis is an inflammatory response of the skin as a result of exposure to an exogenous agent and the agent, which produces this type of dermatitis, is called the contact allergen or contactant. It is a common problem accounting for 4-7% of all dermatological consultations.¹ Metals, as a group, are the most

common contact allergens and nickel ranks as the most common of all screening agents.²⁻⁴ It has been rightly said by the International Nickel Company in a brochure titled "The Romance of Nickel" that "Nickel is with you and does things for you from the time you get up in the morning until you go to sleep at night". On account of its significant public health importance, the American Contact Dermatitis Society (ACDS) named nickel as the "Contact Allergen of the Year" in 2008.⁵ It is commonly found in consumer articles like jewellery, cutlery, kitchen equipment, hardware, sporting goods, wire screens, electrical equipment, metallic buttons, make-up products etc.⁶⁻⁸ Occupational exposure

Address for correspondence

Prof. Dr. Sunil Kumar Gupta
Department of Dermatology, Venereology & Leprology
Dayanand Medical College and Hospital,
Ludhiana, Punjab, India
Email: vsunilgupta@rediffmail.com

to nickel is common in metal plating industries, cashiers, kitchen workers, seamstresses, restaurant workers, and hairdressers.

Other metal allergens widely used in patch test screening kits are cobalt and potassium dichromate. Since cobalt coexists with nickel in nature, nickel based alloys used in consumer products also contain cobalt, resulting in concomitant exposure and sensitization to both the metals. Cobalt is used to produce jewellery,⁹ spectacle frames and wristwatches.¹⁰ People may also be exposed to cobalt in certain nonmetallic products, for example, detergents and pigments used in paint, tattoos, and shoes.¹¹ The main source of chromium is cement, although the content varies widely depending upon the manufacturer. Construction workers, artists and do-it-yourself homebuilders are exposed to cement and the hazards of cement dermatitis. Occupational contact dermatitis clinics in Germany conducted patch tests on construction workers, which showed that potassium dichromate was the commonest allergen (31.9%) among them whereas chromate sensitivity was found in less than 2% of patients attending the general patch test clinic.¹²

The present study was conducted with an aim to determine the prevalence of metal sensitization and different clinical patterns in suspected patients of metal induced contact dermatitis.

Methods

This was an epidemiological study conducted from January 2013 to December 2014. A total of 100 patients suspected and provisionally diagnosed, as having contact dermatitis to metals and attending the dermatology OPD at DMC&H Ludhiana were selected for the study. Detailed clinical history and cutaneous examination with special reference to pattern and morphology of dermatitis, exacerbating factors, sites involved

and other suspected allergies was performed in each case. Particular attention was paid to risk factors in the form of ear piercing, occupational triggers like jewellery related profession, industrial workers and other miscellaneous professions related to metal work. Patients presenting with acute dermatitis, those on steroids and other immunosuppressants were excluded from the study. The patients were then subjected to a patch test with Indian Standard Battery developed by CODFI (Contact and Occupational Dermatoses Forum of India). Metal allergens in the testing kit included potassium dichromate, nickel sulphate and cobalt sulphate.

The results were read after 48 hours of application. First reading was taken 45-60 minutes after removing patches allowing adequate time for erythema due to stripping of tape to settle down. Second reading was taken, if required, after 72 or 96 hours to confirm the presence of allergic reaction that persisted or increased while irritant reaction decreased. In all the patients, clinical relevance of positive allergens was determined based upon the history and examination to isolate all the probable contactants.

Results

In the series, 35 females tested positive as compared to 20 males. Overall, 55 of the 100 tested patients had a positive patch test reaction. The age group with the most patch test positive patients was 40-49 years (15 patients). Age and sex distribution of these patients is outlined in **Table 1**.

Twenty-seven patients showed positive reactions to metal allergens. 24 patients among these i.e. 88.29% showed a positive reaction to single metal allergen. Cross sensitivity was low and three patients or 11.11% showed sensitization to

Table 1 Age and sex distribution.

Age (years)	Males			Females			Total		
	No. of cases tested	Positive cases	%	No. of cases tested	Positive cases	%	No. of cases tested	Positive cases	%
10-19	3	2	66.67%	5	4	80.00%	8	6	75.00%
20-29	5	4	80.00%	19	7	36.84%	24	11	45.83%
30-39	9	3	33.33%	14	8	57.14%	23	11	47.83%
40-49	6	2	33.33%	19	13	68.42%	25	15	60.00%
50-59	13	7	53.85%	5	3	60.00%	18	10	55.56%
≥60	2	2	100.00%	0	0	0.00%	2	2	100.00%
Total	38	20	52.63%	62	35	56.45%	100	55	55.00%

Table 2 Distribution of patients according to clinical patterns

Clinical pattern	No. of cases tested	Patients with positive results		Patients with positive results to probable contactants (relevant)	
		No.	%	No.	%
Localized (to area of contact)	20	16	29.09%	14	35.90%
Face	15	9	16.36%	8	20.51%
Hand	40	18	32.73%	7	17.95%
Mixed	5	4	7.27%	4	10.26%
Foot	10	3	5.45%	3	7.69%
ABCD	2	2	3.64%	1	2.56%
Acrofacial	1	1	1.82%	1	2.56%
Palmoplantar	7	2	3.64%	1	2.56%
Total	100	55	100.00%	39	100.00%

ABCD = air-borne contact dermatitis

Table 3 Correlation of positive patch test results with occupation

Occupation	No. of cases tested	Patients with positive results		Patients with positive results to probable contactants (relevant)	
		No.	%	No.	%
Housewife	42	24	43.64%	17	43.59%
Student	15	9	16.36%	5	12.82%
Farmer	10	6	10.91%	4	10.26%
Teacher	6	3	5.45%	3	7.69%
Construction worker	3	3	5.45%	2	5.13%
Medical/Paramedical staff	4	2	3.64%	2	5.13%
Service/clerical work	4	2	3.64%	2	5.13%
Businessman/ shopkeeper	7	2	3.64%	1	2.56%
Mechanic	2	1	1.82%	1	2.56%
Police	1	1	1.82%	1	2.56%
Retired	2	2	3.64%	1	2.56%
Chemical engineer	1	0	0.00%	0	0.00%
Gardener	1	0	0.00%	0	0.00%
IT Engineer	1	0	0.00%	0	0.00%
Tailor	1	0	0.00%	0	0.00%
Total	100	55	100.00%	39	100.00%

two metal allergens. Among them, two patients had a positive reaction to nickel and cobalt both while one patient had a positive reaction to

nickel and chromate. No patient showed sensitization to all the three metals. Cross sensitization was not seen in males. The highest




Figure 1 Etiological profile of various allergens established by patch testing



Figure 1 Dermatitis at site of contact with belt buckle.

number of positive responses was seen to nickel sulphate (21%). Details of the results obtained




Figure 2 Earlobe dermatitis due to earrings

after testing with the twenty allergens in the kit are shown in Figure 1. Figure 2 and 3 show examples of the clinical presentations showing positive reactions to metal allergens.

Clinically relevant reactions or positivity to probable allergens was observed in 39% of the cases while 16% showed a positive reaction to an unrelated or irrelevant allergen.

The clinical involvement observed in the patients was grouped into eight clinical patterns according to the sites involved.

- Hand eczema
- Foot eczema
- Palmoplantar eczema
- Facial dermatitis (including eyelid dermatitis)
- Acrofacial dermatitis (involving hands and face)
- Dermatitis localized to area of contact with metal, such as involvement of earlobes due to earrings
- Airborne contact dermatitis like pattern (ABCD)
- Mixed pattern or generalized (involving two or more of the clinical patterns described above).

Distribution of positive reactions in these groups is depicted in **Table 2**. Occupational history of the patients formed an important part of their evaluation and this was correlated with their respective patch tests results (**Table 3**).

Discussion

In the present study, 27 patients (27%) tested positive to metal allergens. Nickel had the highest incidence of positivity (21 patients or 21%). Hence, incidence of nickel hypersensitivity in metal reactors was found to be 77.78%. Cobalt and chromium were positive in three and six patients each. Clinical relevance for nickel, cobalt and chromium was present in 100%, 33.33% and 66.67% of the patients respectively. On reviewing the literature, nickel has been reported as the most allergen in many studies (**Table 4**).

Some European studies have reported decreasing prevalence of nickel allergy following the European Union Nickel Directive, according to which nickel content and release rates have been limited to $<0.5 \mu\text{g}/\text{cm}^2/\text{week}$ for products intended to come into direct and prolonged contact with the skin. Johansen et al showed that the prevalence of nickel allergy decreased significantly from 24.8% in 1985-1986 to 9.2% in 1997-1998 in Danish children aged 0-18 years.²⁵ In Poland, the prevalence of nickel allergy decreased from 15.9% in 1995 to 10.0% in 2004 in female dermatitis patients aged under 20 years.²⁶ To the best of our knowledge, no such regulation has been made in India so far.

In the present study, twenty six females tested positive to a metal allergen as compared to four males (male:female ratio= 2:13). Among the twenty-one positive patients, females again outnumbered males by 20:1 and this difference was significant ($p = 0.000$). Higher incidence of nickel hypersensitivity in females has been reported in various other studies. A German study conducted by Uter *et al.*²⁷ utilizing patch test data from 74,940 patients showed a strong association between female gender and nickel allergy with a prevalence ratio of 3.74. Thyssen *et al.*²⁸ showed that the median prevalence of nickel allergy was higher among women than men (17.1%). Calnan²⁹ and Gaul³⁰ individually described that nickel allergy was much more common in female patients due to nickel in suspenders and jewellery. However, in 1969, Kanan reported that prevalence of nickel allergy was higher in male Kuwaiti dermatitis patients, mainly due to nickel in wristwatches and press-studs in underpants.³¹ Overall, the higher prevalence of nickel allergy in women can be explained by the higher prevalence of ear and body piercing and exposure to costume jewellery in women than in men.

Table 4 Most common metal allergen in published literature

<i>Authors</i>	<i>Year of publication</i>	<i>Most common metal allergen</i>
Davis <i>et al.</i> ¹³	2011	Nickel (22.5%)
García-Rabasco <i>et al.</i> ¹⁴	2014	Nickel (24.2%)
Khatami <i>et al.</i> ¹⁵	2013	Nickel (20%)
Akyol <i>et al.</i> ¹⁶	2005	Nickel (17.6%)
Goon <i>et al.</i> ¹⁷	2005	Nickel (19.9%)
Cheng <i>et al.</i> ¹⁸	2008	Nickel (17.7%)
Fransway <i>et al.</i> ¹⁹	2013	Nickel (19.5%)
Warshaw <i>et al.</i> ²⁰	2013	Nickel (15.5%)
Dou <i>et al.</i> ²¹	2011	Nickel (25.7%)
Handa <i>et al.</i> ²²	2011	Potassium dichromate (12.3%)
Hassan <i>et al.</i> ²³	2013	Nickel (16.1%) Potassium dichromate (16.1%)
Majid ²⁴	2014	Nickel (12.8%)
Present study	2014	Nickel (21%)

Most of the patients in this series presented with hand eczema (40%), which was followed by dermatitis localized to site of contact with metal (20%) and facial dermatitis including eyelid dermatitis (15%). On further investigation with patch testing highest frequency of nickel positivity was seen in localized pattern (61.90% of nickel sensitive patients) followed by hand eczema (28.57%). Hand eczema was also the most common presenting dermatitis in a study conducted by Fransway *et al.*¹⁹ and Ruff *et al.*³² while airborne contact dermatitis (ABCD) affecting face, neck, flexures of arms, and legs was the most common pattern, followed by localized allergic contact dermatitis, hand dermatitis and footwear dermatitis in a study conducted by Handa *et al.*²² Our findings differ from those obtained by Cheng *et al.*¹⁸ Among the metal-sensitized individuals, common sites of dermatitis were the upper extremities, face, and hands in decreasing order. Those patients who were nickel positive showed a predilection for face (37.6%), upper extremities (36.6%) and hands (17.9%). One explanation for this difference could be the fact that our study, all the patients presenting with dermatitis of limbs, abdomen etc. were clubbed together as a single group of patients i.e. localized dermatitis.

Potassium dichromate was the most common metal allergen reported from a contact dermatitis

clinic in North India,²² in which 12.3% patients showed a positive reaction to potassium dichromate followed by 11.9% to nickel sulphate. However, in their study males (54.1%) outnumbered females (45.9%) and this could account for higher positivity to chromate since chromate sensitivity is more common in males while nickel sensitivity is commoner in females. This difference could also be partly attributed to the fact that chromium sensitivity is primarily seen in construction workers who constituted only a minority of the patients in our study (3 patients).

Conclusion

Metal induced contact dermatitis is a frequently encountered problem in dermatology clinics and it can manifest in different clinical patterns. The main limitation in our study was that dimethylglyoxime test was not incorporated into the study, hence, source of exposure or evidence of nickel release from the patients' own objects could not be identified. The representation of metal workers (mechanics, plating industry workers etc.) was less. This was probably due to the fact that they comprise a lower income group and prefer going to government hospitals. Nickel was the commonest metal allergen identified and it was relevant in 100% of the cases. Nickel hypersensitivity showed a strong female

preponderance. Patch testing is hence a helpful diagnostic aid in identifying the agents responsible for contact dermatitis and a sincere effort should be made to determine clinical relevance of the test results in every case.

References

1. Coenraads PJ, Diepgen T, Uter W. Epidemiology. In: Frosch PJ, Menné T, Lepoittevin JP, editors. *Contact dermatitis. 4th ed.* Berlin: Springer; 2006. p. 135–63.
2. Warshaw EM, Belsito DV, DeLeo VA *et al.* North American Contact Dermatitis Group patch-test results, 2003-2004 study period. *Dermatitis.* 2008;19:129-36.
3. Hogan DJ, Hill M, Lane PR. Results of routine patch testing of 542 patients in Saskatoon, Canada. *Contact Dermatitis.* 1988;19:120-4.
4. Lee TY, Lam TH. Patch testing of 490 patients in Hong Kong. *Contact Dermatitis.* 1996;35:23-6.
5. Kornik R, Zug KA. Nickel. *Dermatitis.* 2008;19:3-8.
6. Berne B, Boström Å, Grahnén AF, Tammela M. Adverse effects of cosmetics and toiletries reported to the Swedish Medical Protection Agency 1989–94. *Contact Dermatitis.* 1996;34:359-62.
7. Malten KE, Spruit D. The relative importance of various environmental exposures to nickel in causing contact hypersensitivity. *Acta Derm Venereol.* 1969;49:14-9.
8. Brandrup F, Larsen FS. Nickel dermatitis provoked by buttons in blue jeans. *Contact Dermatitis.* 1979;5:148-50.
9. Perryman JH, Fowler JF Jr. A patch test study to evaluate the allergenicity of a metallic jewelry alloy in patients allergic to cobalt. *Cutis.* 2006;77:77-80.
10. Basketter DA, Briatico-Vangosa G, Kaestner W *et al.* Nickel, cobalt and chromium in consumer products: a role in allergic contact dermatitis? *Contact Dermatitis.* 1993;28:15-25.
11. Goossens A, Bedert R, Zimerson E. Allergic contact dermatitis caused by nickel and cobalt in green plastic shoes. *Contact Dermatitis.* 2001;45:172.
12. Geier J, Schnuck A. A comparison of contact allergens among construction and non-construction workers attending contact dermatitis clinics in Germany: results of the Information Network of Departments of Dermatology from November 1989 to July 1993. *Am J Contact Dermatitis.* 1995;6:86-94.
13. Davis MD, Wang MZ, Yiannias JA *et al.* Patch testing with a large series of metal allergens: findings from more than 1,000 patients in one decade at Mayo Clinic. *Dermatitis.* 2011;22:256-71.
14. García-Rabasco AE, Zaragoza-Ninet V, García-Ruiz R, de la Cuadra-Oyanguren J. Allergic contact dermatitis due to nickel: descriptive study in a tertiary hospital, 2000-2010. *Actas Dermosifiliogr.* 2014;105:590-6.
15. Khatami A, Nassiri-Kashani M, Gorouhi F *et al.* Allergic contact dermatitis to metal allergens in Iran. *Int J Dermatol.* 2013;52:1513-8.
16. Akyol A, Boyvat A, Peksari Y, Gürgey E. Contact sensitivity to standard series allergens in 1038 patients with contact dermatitis in Turkey. *Contact Dermatitis.* 2005;52:333-7.
17. Goon AT, Goh CL. Metal allergy in Singapore. *Contact Dermatitis.* 2005;52:130-2.
18. Cheng TY, Tseng YH, Sun CC, Chu CY. Contact sensitization to metals in Taiwan. *Contact Dermatitis.* 2008;59:353-60.
19. Fransway AF, Zug KA, Belsito DV *et al.* North American Contact Dermatitis Group patch test results for 2007-2008. *Dermatitis.* 2013;24:10-21.
20. Warshaw EM, Belsito DV, Taylor JS *et al.* North American Contact Dermatitis Group patch test results: 2009 to 2010. *Dermatitis.* 2013;24:50-9.
21. Dou X, Zhao Y, Ni C *et al.* Prevalence of contact allergy at a dermatology clinic in China from 1990-2009. *Dermatitis.* 2011;22:324-31.
22. Handa S, Jindal R. Patch test results from a contact dermatitis clinic in North India. *Indian J Dermatol Venereol Leprol.* 2011;77:194-6.
23. Hassan I, Rather PA, Jabeen Y *et al.* Preliminary experience of patch testing at Srinagar, Kashmir. *Indian J Dermatol Venereol Leprol.* 2013;79:813-6.
24. Majid I. Contact allergens in Kashmiri population: Results from a 6-year patch testing experience in 550 patients. *Indian J Dermatol Venereol Leprol.* 2014;80:62-4.

25. Johansen J, Menne T, Christophersen J, Kaaber K, Veien N. Changes in the pattern of sensitization to common contact allergens in Denmark between 1985–86 and 1997–98, with a special view to the effect of preventive strategies. *Br J Dermatol*. 2000;**142**:490-5.
26. Rudzki E, Rebandel P. Changes in the pattern of sensitization to nickel in different age groups in Poland. *Contact Dermatitis*. 2005;**53**:177.
27. Uter W, Pfahlberg A, Gefeller O *et al*. Risk factors for contact allergy to nickel – results of a multifactorial analysis. *Contact Dermatitis*. 2003;**48**:33-8.
28. Thyssen JP, Linneberg A, Menne T, Johansen JD. The epidemiology of contact allergy in the general population – prevalence and main findings. *Contact Dermatitis*. 2007;**57**:287-99.
29. Calnan CD. Nickel dermatitis. *Br J Dermatol*. 1956;**68**:229-36.
30. Gaul LE. Incidence of sensitivity to chromium, nickel, gold, silver and copper compared to reactions to their aqueous salts including cobalt sulfate. *Ann Allergy*. 1954;**12**:429-44.
31. Kanan MW. Contact dermatitis in Kuwait. *J Kwt Med Assoc*. 1969;**3**:129-44.
32. Ruff CA, Belsito DV. The impact of various patient factors on contact allergy to nickel, cobalt, and chromate. *J Am Acad Dermatol*. 2006;**55**:32-9.