

Research progress of the studies on the roots of *Peucedanum praeruptorum* dunn (Peucedani radix)

Yuelin Song, Wanghui Jing, Ru Yan and Yitao Wang*

State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

Abstract: As a commonly employed traditional Chinese medicine, Peucedani Radix (Qian-hu in Chinese), which consists of the dried roots of *Peucedanum praeruptorum* Dunn, has a long history of application for the treatment of cough with thick sputum and dyspnea, nonproductive cough and upper air-way infections in traditional medicinal practice. The current review aims to summarize the research progress on the botany, phytochemistry, chemical analysis, pharmacological assay, and pharmacokinetic profile of this famous herbal drug. All available information on this traditional medicine was obtained via electronic search (using ACS, PubMed, Web of Science, Google Scholar, Baidu Scholar, and CNKI). Phytochemical investigations revealed that angular-type pyranocoumarins (APs), mainly (\pm)-praeruptorin A (Pd-Ia), (+)-praeruptorin A, (\pm)-praeruptorin B, (+)-praeruptorin B (Pd-II) and (+)-praeruptorin E (Pd-III), were the main active components in Qian-hu, while some other types of ingredients were also identified from this herb. The crude extract and pure compounds from Peucedani Radix exhibited a wide spectrum of *in vitro* and *in vivo* pharmacological activities, including vasorelaxant, cardioprotective, hepatoprotective, anti-tumor and anti-platelet aggregative effects. Conversely to the well-defined chemical constituents and activities, the properties of absorption, pharmacokinetics, and metabolism were rarely characterized. However, further investigations are wistful for the development of new drugs and therapies for various diseases, especially cardiovascular disorders. Collectively, the present review on the phytochemistry, chemical analysis, pharmacological evaluation, and pharmacokinetic profile of Peucedani Radix will provide meaningful information for further studies and commercial exploitation of the herbal medicine.

Keywords: Peucedani Radix; angular-type pyranocoumarins; chemical constituents; absolute configuration; pharmacological activities; pharmacokinetics

INTRODUCTION

Peucedani Radix consists of the dried roots of perennial herbaceous plant *Peucedanum praeruptorum* Dunn of Apiaceae family and initially documented in Miscellaneous Records of Famous physicians (Ming Yi Bie Lu). This herbal medicine is named as Qian-hu in Chinese, and also known as: aunt dishes, luo ghost food, water Qian-Hu, udo, chicken feet, etc. (Rao *et al.*, 1995) in some regions of China. This herbal medicine is acrid and bitter in flavour and slightly cold in nature. It is documented as tropistic to the lung channel.

The original plant is widely distributed in Zhejiang, Anhui, Jiangxi, Fujian, Jiangsu, Henan and Hunan provinces in China. After the Ministry of Agriculture of agricultural products quality and safety center review and expert appraisal, Peucedani Radix cultured in Ningguo country, Anhui province, in accordance with “The measures for the administration of geographical indication of agricultural products” registration protection conditions, which was stipulated by the Ministry of Agriculture in May 2010, formally issued the “Ning hogfennel agricultural product geographical indications

registration certificate”, the implementation of protection by law (Xiang, 2006).

Following the theory of traditional Chinese medicine, the bitterness of this herbal medicine usually could send down abnormally ascending of Qi, thus resolving cough to stop cough; the acridness is dispersing; and slight coldness can clear heat. Hence, it can send down the adverse Qi, resolve phlegm to stop cough, and disperse wind-heat, indicated the prospects of treatment in cough due to phlegm-heat, and those with cough and sore-throat caused by invasion of lung by wind-heat.

Before 2005, the roots of *Peucedanum decursivum* Maxim. (*Angelica decursive*, Zi-hua Qian-hu) were also listed in Chinese Pharmacopoeia as one of the two original source of Peucedani Radix (Rao *et al.*, 1995). This plant was removed owing to the significant difference of appearance and active characteristics between *P. decursivum* and *P. praeruptorum*. Although the sole source was authorized, many kinds of plants are still sold as the counterfeits and substitutes of Peucedani Radix in current material market, such as *P. decursivum* Maxim. (Zi-hua Qian-hu), *P. rubricaulis* Shan et Sheh (Yun Qian-hu), *P. guangxiense* Shan et Sheh (Guang-xi Qian-hu), *P. Mashanense* Shan et Sheh (Ma-shan Qian-

*Corresponding author: e-mail: ytwang@umac.mo

hu), *P. turgeniifolium* Wolff (Chang Qian-hu), *P. japonicum* Thunb. (Bing-hai Qian-hu), *P. wawrae* (Wolff) Su (Tai-shan Qian-hu), *P. medicum* Dunn (Hua-zhong Qian-hu), *P. medicum* Dunn var. *gracile* Dunn ex Shan et Sheh (Yan Qian-hu), *P. dissolutum* (Diels) Wolff (Nan-chuan Qian-hu), *P. wulongense* Shan et Sheh (Wu-long Qian-hu), *P. formosanum* Hyata (Tai-wan Qian-hu), *P. terebinthaceum* (Fisch.) Fisch. ex Turcz (Shifangfeng), *P. diellisianum* Fedde ex Wolff (Zhu-jie Qian-hu), *P. harry-smithii* Fedde ex Wolff (Hua-bei Qian-hu), *P. harry-smithii* var. *suglabrum* Shan et Sheh (Shao-mao Bei-qian-hu), etc (Rao *et al.*, 1995). Consequently, it is critically important to review the research progress for the chemical evaluation, qualitative and quantitative analysis of ingredients in *Peucedani Radix* and metabolic and pharmacokinetic profiles of this herb.

Chemical constituents

A plenty of chemical components have been identified from *Peucedani Radix*, including coumarins, flavones, sugars, saponins, steroids and volatile oils, while coumarins, in particular angular-type pyranocoumarins (APs), are regarded as the main active constituents contributing to the Qian-hu effect (Kong, 2010). Up to now, more than forty coumarins categorized to this type were identified from the roots of *P. praeruptorum*, and the chemical structures of these APs were summarized in fig. 1.

In addition to those angular-type pyranocoumarins, other components including praeroside I (44) (Okuyama *et al.*, 1989), rutarin (45) (Kong *et al.*, 1994a), isorutarin (46) (Kong *et al.*, 1994a), psoralen (47) (Kong *et al.*, 1993c), 8-methoxypsoralen (48) (Kong *et al.*, 1993c), marmesinin (49) (Okuyama *et al.*, 1989), nodakenin (50) (Kong *et al.*, 1994a), nodakenetin (51) (Liu and Xu, 2002), decuroside V (52) (Takata *et al.*, 1990b), bergapten (5-methoxypsoralen, 53) (Kong *et al.*, 1993c), apiosylskimmin (54) (Takata *et al.*, 1990b), skimming (55) (Takata *et al.*, 1990b), (-)-peucedanol (56) (Kong *et al.*, 1993b), scopolin (57) (Takata *et al.*, 1990b), scopoletin (58) (Kong *et al.*, 1994b), umbelliferone (59) (Chang and Li, 1999b), vanillic acid (60) (Kong *et al.*, 1994a), gallic acid (61) (Kong *et al.*, 1994a), daucosterol (62), β -sitosterol (63) (Kong *et al.*, 1993c), galactitol (64) (Kong *et al.*, 1993b), acetylactyodiolin (65) (Zhang *et al.*, 2005b), tanshinone I (66) (Zhang *et al.*, 2005b), tanshinone II (66) (Zhang *et al.*, 2005b), (-)-sclerodin (67) (Zhang *et al.*, 2006), palmitic acid (68) (Zhang *et al.*, 2006), imperatorin (69) (Zhang *et al.*, 2006), 2,6-dimethyl quinoline (70) (Zhang *et al.*, 2006), tetracosanoic acid (71) (Zhang *et al.*, 2006), angelicin (72) (Chang and Li, 1999a), arnocoumarin (73) (Chang and Li, 1999a), 3,4-methylenedioxy-8-methoxy-9,10-dihydrophenanthric acid,9,10-dione (74) (Zhang *et al.*, 2010), praeroside VI (75) (Ishii *et al.*, 2008), hymexelsin (76) (Ishii *et al.*, 2008), isofraxidin (77) (Ishii *et al.*, 2008), 8-carboxy-7-


hydroxycoumarin (78) (Ishii *et al.*, 2008), praeroside VII (79) (Chang *et al.*, 2008), peucedanoside A (80) (Chang *et al.*, 2007b), peucedanoside B (81) (Chang *et al.*, 2007b), apterin (82) (Chang *et al.*, 2007b), 5,8-dimethoxypsoralen (83) (Kong *et al.*, 1996b), isoscopoletin (84) (Kong *et al.*, 1996b), anchoic acid (85) (Kong *et al.*, 1996b), isoboconin (86) (Chang and Li, 1999b), aegelinol (87) (Chang and Li, 1999b), faltarindiol (88) (Miyazawa *et al.*, 1996), qianhuocoumarin F (89) (Kong and Li, 1994), Pd-C-I (90) (Kong *et al.*, 1994a), baihuaqianhuoside (91) (Kong *et al.*, 1994a), etc., were also isolated and identified from the roots of *P. praeruptorum*.

Determination of absolute configuration of angular-type pyranocoumarins

In general, the angular-type pyranocoumarins (APs) consist of the *cis/trans*-kellactone skeletons and acyl substituents at C-3' and/or C-4' positions, which were always mentioned as two chiral centers. The identification of acyl groups was traditionally achieved using NMR spectroscopy, yet electron ionization-tandem mass spectrometry (EI-MS/MS) and electrospray ionization-tandem mass spectrometry (ESI-MS/MS) were widely introduced in recently years. The acyl moieties at C-3' and C-4' positions would be successively lost in EI-MS/MS (fig. 2) (Swager and Cardellina II, 1985), while the cleavage of C-3' acyl group was prior to that at C-4' position in ESI-MS/MS case (fig. 3) (Tao *et al.*, 2009).

It is crucial to determine the absolute configurations of these two carbons, since the activities of APs are governed by their absolute chemical structures. In 1980s, $^1\text{H-NMR}$ spectroscopy was widely employed to elucidate the relative configurations using some rules of thumb. For *cis*-kellactone derivatives, the coupling constant ($J_{3',4'}$) between the two protons at C-3' and C-4' position is around 5.0 Hz, and 5'-CH₃ and 6'-CH₃ share almost an identical chemical shift (δ). On the other side, the coupling constant ($J_{3',4'}$) for *trans*-kellactone derivatives locates at the range of 3.0 ~ 6.9 Hz, and the difference between the chemical shifts ($\Delta\delta$) of 5'-CH₃ and 6'-CH₃ ranges from 0.08 ppm to 0.20 ppm. However, Okuyama *et al.* (Okuyama and Shibata, 1981) proposed that the $^{13}\text{C-NMR}$ spectroscopy could provide more important information for relative configuration. The chemical shift difference ($\Delta\delta$) between the C-5' and C-6' of *trans*-kellactone derivatives is between 4.0 ppm and 6.0 ppm, while the C-5' and C-6' of *cis*-kellactone derivatives almost share an identical chemical shift (δ).

In order to determine the absolute configurations of APs, alkaline hydrolysis is generally adopted as a chemical correlation tool to afford both *cis*-kellactone and *trans*-kellactone (fig. 4), which are generated by epimerization at C-4' during alkaline treatment. The absolute configuration of (+)-*cis*-kellactone was determined as *R*-configurations for C-3' and C-4' using X-ray



No.	Name	Con ^b	R ₁	R ₂	Reference
1	(±)-praeruptorin A (Pd-Ia)	<i>cis</i>	angeloyloxy	acetyloxy	(Okuyama and Shibata, 1981, Chen <i>et al.</i> , 1979, Lu <i>et al.</i> , 2001)
2	(±)-praeruptorin B	<i>cis</i>	angeloyloxy	angeloyloxy	(Chen <i>et al.</i> , 1979, Lu <i>et al.</i> , 2001)
3	(+)-praeruptorin A	<i>cis</i>	angeloyloxy (S)	acetyloxy (S)	(Chen <i>et al.</i> , 1979)
4	(+)-praeruptorin B (Pd-II)	<i>cis</i>	angeloyloxy (S)	angeloyloxy (S)	(Okuyama and Shibata, 1981, Chen <i>et al.</i> , 1979)
5	(-)-praeruptorin A	<i>cis</i>	angeloyloxy (R)	acetyloxy (R)	(Xu <i>et al.</i> , 2010)
6	(-)-praeruptorin B	<i>cis</i>	angeloyloxy (R)	angeloyloxy (R)	(Song <i>et al.</i> , 2012a)
7	(+)-praeruptorin E (Pd-III)	<i>cis</i>	angeloyloxy (S)	isovaleryloxy (S)	(Okuyama and Shibata, 1981, Ye <i>et al.</i> , 1982)
8	Pd-Ib	-	angeloyloxy (R)	oxo	(Wang <i>et al.</i> , 2006, Okuyama and Shibata, 1981)
9	Qianhucoumarin I	<i>cis</i>	acetyloxy (S)	tigloyloxy (S)	(Kong, 1996)
10	(3'S,4'S)-3'-angeloyl-4'-acetylkhellactone	<i>cis</i>	angeloyloxy (S)	acetyloxy (S)	(Lou <i>et al.</i> , 2004)
11	(3'S,4'R)-3'-acetyl-4'-angeloylkhellactone	<i>trans</i>	acetyloxy (S)	angeloyloxy (R)	(Lou <i>et al.</i> , 2004)
12	(3'R,4'S)-3'-acetyl-4'-tigloylkhellactone	<i>trans</i>	acetyloxy (R)	tigloyloxy (S)	(Lou <i>et al.</i> , 2004)
13	(3'R,4'R)-3'-acetyl-4'-angeloylkhellactone (pteryxin)	<i>cis</i>	acetyloxy (R)	angeloyloxy (R)	(Lou <i>et al.</i> , 2004, Takata <i>et al.</i> , 1990a, Kong <i>et al.</i> , 1993c)
14	(3'S,4'R)-3'-acetyl-4'-isobutyrylkhellactone	<i>trans</i>	acetyloxy (S)	isobutyryloxy (R)	(Lou <i>et al.</i> , 2004)
15	peucedanocoumarin I	<i>trans</i>	isovaleryloxy (S)	acetyloxy (R)	(Takata <i>et al.</i> , 1990a)
16	peucedanocoumarin II	<i>trans</i>	acetyloxy (S)	angeloyloxy (R)	(Takata <i>et al.</i> , 1990a)
17	peucedanocoumarin III	<i>trans</i>	acetyloxy (S)	tigloyloxy (R)	(Takata <i>et al.</i> , 1990a)
18	qianhucoumarin H	<i>trans</i>	angeloyloxy (S)	isovaleryloxy (R)	(Kong <i>et al.</i> , 1996b)
19	(3'R,4'S)-3'-angeloylkhellactone	<i>trans</i>	angeloyloxy (R)	hydroxyloxy (S)	(Kong <i>et al.</i> , 1993a)
20	qianhucoumarin A	<i>cis</i>	hydroxyl (R)	tigloyloxy (R)	(Kong <i>et al.</i> , 1993c)
21	qianhucoumarin B	<i>cis</i>	acetyloxy (S)	hydroxyl (S)	(Kong <i>et al.</i> , 1993a)
22	qianhucoumarin C	<i>cis</i>	hydroxyl (S)	acetyloxy (S)	(Kong <i>et al.</i> , 1993a)
23	qianhucoumarin D	<i>cis</i>	acetyloxy (S)	acetyloxy (S)	(Kong <i>et al.</i> , 1994b)
24	qianhucoumarin E	-	tigloyloxy (R)	oxo	(Kong <i>et al.</i> , 1994b)
25	(+)-laserpitin	<i>cis</i>	hydroxyl (S)	angeloyloxy (S)	(Hou <i>et al.</i> , 2010)
26	qianhucoumarin J	<i>cis</i>	angeloyloxy (S)	propionyloxy (S)	(Hou <i>et al.</i> , 2010)
27	<i>cis</i> -3',4'-diseneciolykhellactone	<i>cis</i>	seneciolyloxy	seneciolyloxy	(Chang and Li, 1999b)
28	isobocconin	<i>trans</i>	acetyloxy	isobutyryloxy	(Chang and Li, 1999b)
29	aegelinol	-	hydroxyl (R)	dihydro	(Chang and Li, 1999b)
30	<i>trans</i> -khellactone	<i>trans</i>	hydroxyl (R)	hydroxyl (S)	(Chang and Li, 1999b)
31	<i>cis</i> -khellactone	<i>cis</i>	hydroxyl (R)	hydroxyl (S)	(Willettee and Soine, 1962)

No.	Name	Con ^b	R ₁	R ₂	Reference
32	3'-acetyl-4'-isovalerylkhellactone	<i>cis</i>	acetyloxy (R)	isovaleryloxy (R)	(Willettee and Soine, 1962)
33	<i>cis</i> -3'-seneciroyl-4'-angeloylkhellactone	<i>cis</i>	seneciroyloxy (R)	angeloyloxy (R)	(Chang and Li, 1999a)
32	<i>cis</i> -3',4'-diisovalerylkhellactone	<i>cis</i>	isovaleryloxy	isovaleryloxy	(Zhang <i>et al.</i> , 2005b)
35	3'-isovaleryl-4'-keto-khellactone ^a	-	isovaleryloxy	oxo	(Hou <i>et al.</i> , 2009)
36	3'-angeloyl-4'-propionylkhellactone ^a	-	angeloyloxy	propionyloxy	(Hou <i>et al.</i> , 2009)
37	praeroside II	<i>cis</i>	glucosyloxy (R)	hydroxyl (R)	(Takata <i>et al.</i> , 1988)
38	praeroside III	<i>trans</i>	glucosyloxy (S)	hydroxyl (R)	(Takata <i>et al.</i> , 1988)
39	praeroside IV	-	glucosyloxy (R)	dihydro	(Takata <i>et al.</i> , 1988)
40	praeroside V	-	dihydro	glucosyloxy (S)	(Takata <i>et al.</i> , 1988)
41	praeroside VI	<i>cis</i>	apiosyl(1→6)glucosyloxy	hydroxyl (R)	(Chang <i>et al.</i> , 2008)
42	<i>cis</i> -3'-isovaleryl-4'-seneciroylkhellactone	<i>cis</i>	isovaleryloxy (S)	seneciroyloxy (S)	(Jong <i>et al.</i> , 1992)
43	<i>cis</i> -3'-isobutyryl-4'-acetylkhellactone	<i>cis</i>	isobutyryloxy (R)	acetyloxy (R)	(Kong <i>et al.</i> , 2003)

Fig. 1: The chemical structures of angular-type pyranocoumarins (APs, 1-43) identified from the roots of *Peucedanum praeruptorum* Dunn.

^athe chemical structure was determined using LC-MS/MS and the absolute configurations haven't been indicated.

^brelative configuration

spectrometry and the identity of (-)-*cis*-khellactone was thus characterized as (-)-(3'S, 4'S)-*cis*-khellactone. The hydrolyzed products were purified and compared with (+)-*cis*-khellactone and (-)-*cis*-khellactone. If (+)-*cis*-khellactone was identified as one of the hydrolyzed product, the configuration of C-3' should be characterized as *R*, while (-)-*cis*-khellactone only could be yielded by (3'S)-angular-type pyranocoumarin. Further, the configuration of C-4' was determined by the different NMR spectroscopic data between *cis*- and *trans*-type pyranocoumarins.

In addition, circular dichroism (CD) spectroscopy was recently subjected to carry out the determination of absolute configurations (Lou *et al.*, 2004, Xu *et al.*, 2010). A practical rule relating the position and absolute stereochemistry of the khellactone derivatives to the behavior of their Cotton effects in CD curves was proclaimed by Lou *et al.* (Lou *et al.*, 2004).

Chemical analysis and quality control of *peucedani radix*

Due to the different original herbs in current market, it is significant to identify the germplasm origin of Qian-hu. The microscopic features of the radix of *P. praeruptorum* Dunn were proposed in 1980s. As a widely adopted technique, thin-layer chromatography (TLC) was introduced to authorize this herbal drug by Song *et al* and Zhou *et al.* (Song and Xing, 2000, Zhou, 2008). In previous literatures, ¹H-NMR spectroscopy has been employed for chemotaxonomic evaluation of this traditional Chinese herbal drug (Wu *et al.*, 1996, Liu *et*

al., 1999, Wang *et al.*, 1996, Ye *et al.*, 1995), and the results revealed that ¹H-NMR spectroscopy is a quick, simple and reliable means to distinct Qian-hu from other similar herbal drugs, including *P. decursivi*, *P. medicum*, *P. rubicanle* and so on, while praeruptorin A and some other angular-type pyranocoumarins were adopted as the chemical indicators for recognition. In addition, a methodology of sequence characterized amplified regions (SCAR) markers was developed to distinguish of the three similar species of medicinal herbs, including *Angelica decursiva* (*P. decursivum*), *P. praeruptorum* and *Anthriscus sylvestris*, based on the random amplified polymorphic DNA (RAPD) and internal transcribed spacer (ITS) sequence, and this SCAR marker could also provide valuable information for verifying whether the official drug was mixed with an adulterate (Choo *et al.*, 2009).

High performance liquid chromatography coupled with diode array detection and tandem mass spectrometry (HPLC-DAD-MS/MS) was also subjected as a preferable technique for rapid identification and chemical profiling of *P. praeruptorum* (Tao *et al.*, 2009, Zhu *et al.*, 2004). In most cases, praeruptorin A, praeruptorin B and praeruptorin E were adopted the indicators for quality control of Peucedani Radix using HPLC-UV (Zhang *et al.*, 2005a, Kong *et al.*, 1996a, Xu *et al.*, 2001a, Wu *et al.*, 2009, Xu *et al.*, 2007, Wang, 2004) or gas chromatography (GC) (Xu *et al.*, 2001b). The quality standard of Peucedani Radix was proposed by Zhang *et al.* adopting HPLC-UV (Zhang *et al.*, 2005a).

Pharmacological activities of *Peucedani Radix* and the chemical components

Pharmacological activities of the crude extract

Extensive pharmacological activities were revealed for this herbal medicine, which was traditionally adopted the treatments of cough with thick sputum and dyspnea, upper respiratory infections, during screening on modern pharmacological models.




Fig. 2: The proposed ESI-MSⁿ fragmentation scheme for angular-type pyranocoumarins.

As an anciently employed agent for the treatment of asthma, the crude extract of Qian-hu can significantly restrain OVA-induced airway inflammation, airway hyperreactivity and Th2 predominant response on mice mode (Xiong *et al.*, 2012c), and can attenuated the contractions of isolated rabbit trachea smooth muscles by acetylcholine and potassium chloride (Jin *et al.*, 1994).

More interesting, this herbal drug shows a great prospect for the treatment of cardiovascular disease. Thus, there are an increasing number of the pharmacological evaluations concerning on the antihypertensive activity of this herbal medicine. The active components were proved mainly distributed in the petroleum ether extract and could relax pulmonary artery smooth muscle directly as a Ca²⁺ antagonist (Wei *et al.*, 1994). The crude extract was shown to prevent and reverse hypertrophy of renovascular hypertensive left ventricular hypertrophy (LVH) by decreasing the concentration of [Ca²⁺]_i, up-regulating the ATPase activity, improving cardiac function and myocardial compliance (Rao *et al.*, 2002b, Rao *et al.*, 2002c, Ji and Rao, 1996, Zhou *et al.*, 2001a, Zhou *et al.*, 2001b, Zhao *et al.*, 1994). Wang *et al.* mentioned that the therapeutic effect of Qian-hu extract on pulmonary hypertension in patients suffered chronic obstructive pulmonary disease (COPD) may be related with the inhibition on the synthesis or secretion endothelin-1 (ET-1), abnormality of which may provide important contribution in the development of chronic hypoxic pulmonary hypertension (Wang *et al.*, 2001). Wang *et al.*

regarded that Qian-hu extract could down-regulate the pulmonary artery pressure without predominant influencing the systemic artery pressure, decrease the right heart index and reduce the thickness of small pulmonary artery media significantly. The composition of tenascin-C (TN-C) decreased obviously in pulmonary vasculature in rats after treatment of the extract (Wang *et al.*, 2000).




Fig. 3: The proposed ESI-MSⁿ fragmentation scheme for angular-type pyranocoumarins.

The anti-myocardial ischemia action was also revealed in existing reports. Administration of *P. praeruptorum* extract could result in a decrease of the size of acute myocardial ischemia injury (AMI), and the activities of lactate dehydrogenase (LDH), aspartate amino transferase (AST), superoxide dismutase (SOD), methyl diaidehyde (MDA), creatine kinase MB (CK-MB) and creatine kinase (CK), suggesting that the crude extract of Qian-hu exhibited protective effects on AMI (Jiang *et al.*, 2004a, Jiang *et al.*, 2004c, Jiang *et al.*, 2002). The expression of IL-6, Fas, bax and bcl-2 could be modify by the extract of *Peucedani Radix* (Chang *et al.*, 2003, Liu *et al.*, 2002). In addition, significant protection of the microstructure and ultrastructure from reperfusion injury was also observed for Qian-hu extract (Jiang *et al.*, 2004b).

The low polar fraction of *Peucedani Radix* showed cytotoxic activity on *Artemia salina* test and antimicrobial activity on *Streptococcus agalactiae*, *Staphylococcus aureus*, *Escherichia coli*, *Shigella flexneri* and *Salmonella typhi*, while the *n*-butanol and aqueous extracts exhibited none or low activities on these models (Lu *et al.*, 2001, Chen *et al.*, 2002). Moreover, herbal extract was proved pharmacological actions including suppression of hepatic microsomal drug-metabolism enzymes activity in mice (Wang *et al.*, 2004), definite cytotoxic and anti-proliferative action on SGC7901 cells (Liang *et al.*, 2010), oxygen radicals elimination and lipid peroxidation inhibition (Wang *et al.*, 2008).

Pharmacological properties of the major angular-type pyranocoumarins in *Qian-hu*

Angular-type pyranocoumarins, that were observed as the major constituent of *P. praeruptorum* Dunn, in particular praeruptorin A (PA) and praeruptorin B (PB), were

universally regarded to be responsible for the effects of Qian-hu extract mentioned above.




Fig. 4: Representative alkaline hydrolysis pathways of angular-type pyranocoumarins

(±)-Praeruptorin A (PA) could relax the tracheal smooth muscle isolated from rabbits through blocking the potential-dependent Ca^{2+} channels and inhibiting the release of Ca^{2+} from intracellular Ca^{2+} -pool (Guan *et al.*, 1994). This active component could also suppress respiratory tract inflammation, remodel hyperresponsive airway, decrease the levels of IL-4 and IL-13 in BALF along with IgE in serum, increase the level of $\text{INF-}\gamma$ in BALF, down-regulate the expression of pSmad2/3 and TGF- β 1, and up-regulate the expression of Smad7 in pulmonary tissue, as well as inhibit eotaxin protein and mRNA expression, $\text{I}\kappa\text{B}\alpha$ degradation, NF- κB nuclear translocation, NF- κB DNA-binding activity and RelA/p65 phosphorylation in lung. Above all, PA significantly suppressed airway inflammation and airway remodeling induced by ovalbumin stimulation, thus showing great therapeutic potential for allergic asthma (Xiong *et al.*, 2012a, Xiong *et al.*, 2012b).

PA and (+)-praeruptorin A (*d*PA) were widely highlighted as anti-hypertensive drug candidates. PA was usually documented cardiovascular pharmacological activities as a Ca^{2+} antagonist and a K^{+} channel opener (Zhang *et al.*, 2001, Li *et al.*, 1994, Wang *et al.*, 1997, Wang *et al.*, 1995), while *d*PA was commonly mentioned due to the Ca^{2+} antagonistic activity (Rao *et al.*, 1988, Rao *et al.*, 2002a, Yang and Rao, 1997). Interestingly, *d*PA was found to increase coronary blood flow on isolated guinea pig heart (Chen *et al.*, 1979), yet the corresponding effect was absence for PA. Distinct relaxant effects of the levorotatory and dextrorotatory enantiomers of PA was proved by Xu *et al.* on the isolated rat aorta rings, which might be mainly attributed to nitric oxide synthesis mediated by endothelial nitric oxide synthase (Xu *et al.*, 2010).

PA and *d*PA could also prevent and treat ischemic reperfusion injury that was related to the effect of Qian-hu extract, typically through blocking the calcium channels and/or opening the potassium channels (Liu and Chang, 2002, Yang *et al.*, 1995, Chang *et al.*, 2007a, Chang *et al.*, 1994, Chang, 2002).

Moreover, pyranocoumarins, such as PA, *d*PA, (+)-praeruptorin B and (+)-praeruptorin E, exerted anti-inflammatory effects in LPS-stimulated RAW264.7 macrophages through inhibiting NF- κB and STAT3 activation (Yu *et al.*, 2012, Yu *et al.*, 2011). PA could regulate the multiple drug resistant (MDR), differentiation and apoptosis of cancer cells, when it was screened on a couple of cell lines (Fong *et al.*, 2004, Zhang *et al.*, 2003, Wu *et al.*, 2003).

Absorption, Distribution and metabolism of angular-type pyranocoumarins from peucedani radix

In contrast to extensive pharmacologic studies, the pharmacokinetic properties of the coumarin-type components of *Peucedani Radix* have scarcely been studied, hence their contribution to the herbal action and active forms *in vivo* remain unclear.

As an herbal medicine, *Peucedani Radix* is usually taken orally by convention, indicating that it is needed to clarify the intestinal absorption of those active constituents. A couple of *in vitro* models have been developed for permeability and absorption screening, of which the Caco-2 cell (human colon adenocarcinoma cell line) monolayer model is widely mentioned as a preferable tool (Yee, 1997). The transport of (±)-praeruptorin A (PA), (+)-praeruptorin B (*d*PB) and (3'R,4'R)-3',4'-diangeloylkhellactone (*i*PB, anomalin) was evaluated on this model (Zhao *et al.*, 2011, Yue *et al.*, 2012, Jing *et al.*, 2011). The transport property and metabolism of *dl*-PA in Caco-2 cells was characterized under the assistance of a well-developed chiral HPLC-UV method. PA showed a rapid transport across the Caco-2 monolayers, partially bound to cell membranes and underwent hydrolysis during transport. The hydrolysis of PA catalyzed by carboxylesterases was demonstrated, and it implicates extensive first-pass intestinal and hepatic hydrolysis of the tested compound. Slight enantioselectivity was observed in the transport process (Jing *et al.*, 2011). Anomalin [(3'R,4'R)-3',4'-diangeloyl khellactone] was demonstrated as a well-absorbed compound, and the transport mechanism was indicated as passive diffusion (Zhao *et al.*, 2011). At the meanwhile, the P_{app} values of *d*PB for apical (AP) to basolateral (BL) or BL to AP were between 2.0×10^{-6} cm/s and 5.0×10^{-6} cm/s, along with $P_{app}(\text{BL-AP})/P_{app}(\text{AP-BL})$ less than 1.5. As a consequence, *d*PB was proved as a drug candidate with relatively bad absorption, which absorbed mainly by passive diffusion approach through intestinal tract (Yue *et al.*, 2012). The data displayed in literatures also indicated enantiospecific absorption of praeruptorin B enantiomers.

In vitro metabolic models were introduced to characterize the metabolic features of APs in rat liver microsomes and human liver microsomes (RLMs and HLMs) (Jing *et al.*, 2013, Ruan *et al.*, 2011, Song *et al.*, 2012b, Song *et al.*, 2011, Song *et al.*, 2012a). Metabolic pathways including

hydrolysis, oxidation and intra-molecular acyl migration were detected as the main reaction types for *d*PA, *d*PA, *d*PB and *d*PB (Song *et al.*, 2012b, Song *et al.*, 2011). Extensive metabolism was observed for all the screened components in RLMs or HLMs and CYP3A4 was demonstrated to be the main isoenzyme mediating both hydrolysis and oxidation of *d*PA in HLMs, yet CYP 3A1 and/or 3A2 for PA metabolism in RLMs (Jing *et al.*, 2013, Ruan *et al.*, 2011).

PA was revealed that it was rapidly distributed and eliminated from rat plasma and demonstrated linear dynamics in dose range of 5-20 mg/kg following i.v. administration. The mean elimination half-life ($t_{1/2}$) of PA dosing of 5, 10 and 20 mg/kg were approximal 1.0 h. Spleen, heart and lung were observed as the major distribution tissues in rats, and the existence of PA in brain was also revealed since low polarity enabled PA to cross the blood-brain barrier (BBB). No long-term accumulation was detected for PA in all tissues. Low total recoveries were demonstrated for this compound within 24 h (0.120% in urine, 0.097% in bile and 0.009% in feces), which might be caused by significant liver-mediated first pass effect. When PA was i.v. administrated to liver cirrhosis (LC) rats at a single dose of 5 mg/kg, the area under curve (AUC) was significantly greater than that of the control group which could be owing to the slower hepatic blood flow rate and subsequently significant slower hepatic Cl_{int} in LC rats. In addition, the decrease of PA metabolic clearance might be at least partly arouse by the decreased expression and/or activity levels of CYP3A1 and 3A2 in LC rats, which were responsible for PA metabolism (Zhang *et al.*, 2011). Pteryxin [(3'R,4'R)-3'-acetyl-4'-angeloylhellactone], which is the regio-isomer of IPA, also showed rapidly distribution and elimination from mouse plasma ($t_{1/2}$ = 1.463 h). Surprisingly, liver was detected as the major distribution tissue for pteryxin in mice, and the transport of the BBB was observed for pteryxin owing to its low polarity. As was expected, long-term accumulation wasn't observed for pteryxin in all mouse tissues (Wang *et al.*, 2012). Moreover, in the case of *d*PB, the pharmacokinetic profile fitted well into a typical two-compartment model including a fast distribution phase coupled with a relative slow elimination phase. Tissue distribution results proved that the highest concentration of proto drug was detected in pulmonary tissue, followed by tissues of heart, liver and kidney, successively. As a low polar component, *d*PB was detectable in the brain after i.v. administration without surprise, indicating transport across the BBB for *d*PB (Liang *et al.*, 2012).

CONCLUSION

Overall, although many studies were performed on this famous, there are still some shortcomings in these literatures. Enantioselective activities, absorption and

metabolism were observed for the main components in Qian-hu, suggesting that it is crucial to develop enantiospecific method to monitor enantiomers separately for the purpose of quality control and pharmacokinetic study. Moreover, it is also necessary to characterize the chemical profile of this herbal medicine and characterized the herb-related components *in vivo* after oral administration of this plant.

ACKNOWLEDGMENTS

The research was supported by the National Basic Research Program of China 973 program (Grant No. 2009CB522707), the Macao Science and Technology Development Fund (077/2011/A3, 074/2012/A3) and the Research Fund of the University of Macau (MYRG 208 (Y3-L4)-ICMS11-WYT, and MRG012/WYT/2013/ICMS, MRG013/WYT/2013/ICMS).

REFERENCES

- Chang H, Chu XY, Zou J and Chang TH (2007a). Effect of *dl*-praeruptorin A on desmin and vimentin content in rat ischemiareperfusion myocardiocytes. *Chin. Med. J.*, **120**: 2256.
- Chang H, Okada Y, Okuyama T and Tu P (2007b). ¹H and ¹³C NMR assignments for two new angular furanocoumarin glycosides from *Peucedanum praeruptorum*. *Magn. Reson. Chem.*, **45**: 611.
- Chang HT and Li X (1999a). Chemical study of *Peucedanum praeruptorum* Dunn (V). *Zhong Cao Yao*, **30**: 414.
- Chang HT and Li X (1999b). Coumarins from *Peucedanum praeruptorum* Dunn. *Shenyang Yaokexue Xuebao*, **16**: 103.
- Chang HT, Okada Y, Ma TJ, Okuyama T and Tu PF (2008). Two new coumarin glycosides from *Peucedanum praeruptorum*. *J. Asian Nat. Prod. Res.*, **10**: 577.
- Chang TH (2002). Effects of *dl*-praeruptorin A on interleukin-6 level and Fas, bax, bcl-2 protein expression in ischemia-reperfusion myocardium. *Acta Pharmacol Sin.*, **23**: 769.
- Chang TH, Adachi H, Mori N, Saito I and Okuyama T (1994). Effects of a pyranocoumarin compound isolated from a Chinese medicinal plant on ischemic myocardial dysfunction in anesthetized dogs. *Eur. J. Pharmacol.*, **258**: 77.
- Chang TH, Liu XY, Zhang XH, Xing J and Wang HL (2003). Effects of *Peucedanum praeruptorum* Dunn and *dl*-praeruptorin A on IL-6 levels and Fas, bax, bcl-2 protein expressions in ischemia-reperfusion myocardium of rats. *Zhongguo Yike Daxue Xuebao*, **32**: 1.
- Chen BH, Wang MZ and Liu JQ (2002). Chemical constituents of the volatile oil from the roots of

- Peucedanum praeruptorum* and its antibacterial activities. *Redai Yaredai Zhiwu Xuebao*, **10**: 366.
- Chen ZX, Huang BS, She QL and Zeng GF (1979). The chemical constituents of Bai-Hua-Qian-Hu, the root of *Peucedanum praeruptorum* Dunn. (Umbelliferae)--four new coumarins. *Yao Xue Xue Bao*, **14**: 486.
- Choo BK, Moon BC, Ji Y, Choi G, Yoon T and Kim HK. (2009). Development of SCAR markers for the discrimination of three species of medicinal plants, *Angelica decursiva* (*Peucedanum decursivum*), *Peucedanum praeruptorum* and *Anthriscus sylvestris*, based on the internal transcribed spacer (ITS) sequence and random amplified polymorphic DNA (RAPD). *Biol. Pharm. Bull.*, **32**: 24.
- Fong WF, Zhang JX, Wu JY, Tse KW, Wang C, Cheung HY and Yang MS (2004). Pyranocoumarin (\pm)-4'-O-acetyl-3'-O-angeloyl-*cis*-khellactone induces mitochondrial-dependent apoptosis in HL-60 Cells. *Planta Med.*, **70**: 489.
- Guan FL, Jin WB, Zhang XH, Zhao NC and Okuyama T (1994). The effects of praeruptorin A on the contraction of isolated tracheal smooth muscle previously induced by high K^+ acetylcholine. *Zhongguo Yike Daxue Xuebao*, **23**: 549.
- Hou Z, Luo J, Wang J and Kong L (2010). Separation of minor coumarins from *Peucedanum praeruptorum* using HSCCC and preparative HPLC guided by HPLC/MS. *Sep. Purif. Technol.*, **75**: 132.
- Hou Z, Xu D, Yao S, Luo J and Kong L (2009). An application of high-speed counter-current chromatography coupled with electrospray ionization mass spectrometry for separation and online identification of coumarins from *Peucedanum praeruptorum* Dunn. *J. Chromatogr. B*, **877**: 2571.
- Ishii H, Okada Y, Baba K and Okuyama T (2008). Studies of Coumarins from the Chinese Drug Qianhu, XXVII Structure of a New Simple Coumarin Glycoside from Bai-Hua Qianhu, *Peucedanum praeruptorum*. *Chem. Pharm. Bull.*, **56**: 1349.
- Ji Y and Rao MR (1996). Effects of *Peucedanum praeruptorum* extractum on blood pressure, left ventricular hypertrophy and hemodynamic changes in left ventricular hypertrophied hypertensive rats. *Zhongguo Zhongxiyi Jiehe Zazhi*, **16**: 676.
- Jiang MY, Chang TH and Xu YJ (2004a). The protective effect of *Peucedanum praeruptorum* on acute myocardial infarction in anesthetized cats. *Zhongguo Yike Daxue Xuebao*, **33**: 22.
- Jiang MY, Xu YJ, Du X and Pan Q (2004b). Change of myocardial ultrastructure and the protective effect of *Peucedanum praeruptorum* Dunn and its effective component (praeruptorin A, Pd-Ia) against acute myocardial ischemia-reperfusion injury in rats. *Shenyang Yaoke Daxue Xuebao*, **21**: 59.
- Jiang MY, Xu YJ and Shen J (2002). Effects of *Peucedanum praeruptorum* Dunn and *dl*-praeruptorin A on serum SOD and MDA in ischemia-reperfusion myocardium of rats. *Zhongguo Yaoxue Zazhi*, **37**: 623.
- Jiang MY, Xu YJ and Shen J (2004c). Effects of *Peucedanum praeruptorum* Dunn and *dl*-PA on the activity of serum enzyme in ischemia-reperfusion myocardium of rats. *Zhongguo Yaoxue Zazhi*, **39**: 475.
- Jin X, Zhang XH and Zhao NC (1994). Effects of petroleum ether extract of *Peucedanum praeruptorum* Dunn on rabbit trachea smooth muscles. *Zhongguo Zhong Yao Za Zhi*, **19**: 365.
- Jing WH, Song YL, Yan R, Bi HC, Li PT and Wang YT (2011). Transport and metabolism of (+/-)-praeruptorin A in Caco-2 cell monolayers. *Xenobiotica*, **41**: 71.
- Jing WH, Song YL, Yan R and Wang YT (2013). Identification of cytochrome P450 isoenzymes involved in metabolism of (+)-praeruptorin A, a calcium channel blocker, by human liver microsomes using ultra high-performance liquid chromatography coupled with tandem mass spectrometry. *J. Pharm. Biomed. Anal.*, **77**: 175.
- Jong TT, Hwang HC and Jean MY (1992). An antiplatelet aggregation principle and X-ray structural analysis of *cis*-khellactone diester from *Peucedanum japonicum*. *J. Nat. Prod.*, **55**: 1396.
- Kong LY (2010). Systematic research of the material base of Radix Peucedani (Qianhu). *Zhongguo Yaoke Daxue Xuebao*, **41**: 203.
- Kong LY and Li X (1994). Structure elucidation of qianhu coumarin F by 2D NMR. *Bopuxue Zazhi*, **11**: 245.
- Kong LY, Li X, Pei YH, Yu RM, Min ZD and Zhu TR (1994a). Isolation and structure elucidation of baihuaqianhuoside and Pd-C-I from *Peucedanum praeruptorum*. *Yao Xue Xue Bao*, **29**: 276.
- Kong LY, Li X, Pei YH and Zhu TR (1994b). Isolation and structural elucidation of qianhu coumarin D and qianhu coumarin E from *Peucedanum praeruptorum*. *Yao Xue Xue Bao*, **29**: 49.
- Kong LY, Li Y and Min ZD (1996a). Study on analysis of Pd-Ia in *Peucedanum praeruptorum* and Pd-C-I in *Peucedanum decursivum* by RP-HPLC. *Zhongguo Yaoke Daxue Xuebao*, **27**: 215.
- Kong LY, Li Y, Min, ZD, Li X and Zhu TR (1996b). Coumarins from *Peucedanum praeruptorum*. *Phytochemistry*, **41**: 1423.
- Kong LY, Li Y and Niwa M (2003). A new pyranocoumarin from *Peucedanum praeruptorum*. *Heterocycles*, **60**: 1915.
- Kong LY, Min ZD, Li Y, Li X and Pei YH (1996). Qianhu coumarin I from *Peucedanum praeruptorum*. *Phytochemistry*, **42**: 1689.
- Kong LY, Pei YH, Li X, Wang SX, Hou BL and Zhu TR (1993a). The isolation and identification of qianhu coumarin B and qianhu coumarin C from *Peucedanum praeruptorum*. *Yao Xue Xue Bao*, **28**: 772.
- Kong LY, Pei YH, Li X and Zhu TR (1993b). chemical study of *Peucedanum praeruptorum* Dunn (V). *Zhong*

- Cao Yao, **24**: 401.
- Kong LY, Pei YH, Li X, Zhu TR and Okuyama T (1993c). Isolation and structural elucidation of qianhu coumarin A. *Yao Xue Xue Bao*, **28**: 432.
- Li JM, Chang TH, Sun XD, Hao LY, Wang YP, Yu YF and Zhang KY (1994). Effect of *dl*-praeruptorin A on calcium current in ventricular cells of guinea pig. *Acta Pharmacol. Sin.*, **15**: 525.
- Liang T, Yue W, Du X, Ren L and Li Q (2012). Pharmacokinetics and tissue distribution study of praeruptorin d from radix peucedani in rats by high-performance liquid chromatography (HPLC). *Int. J. Mol. Sci.*, **13**: 9129.
- Liang T, Yue W and Li Q (2010). Chemopreventive effects of *Peucedanum praeruptorum* DUNN and its major constituents on SGC7901 gastric cancer cells. *Molecules*, **15**: 8060.
- Liu BM and Xu Q (2002). Analysis of *dl*-praeruptorin A, *dl*-praeruptorin B and nodakenetin in Qianhu by GC/MS and GC. *Guangxi Kexue*, **9**: 294.
- Liu QX, Wu MY, Rao GX, Ye JS and Hui H (1999). Chemotaxonomy of Ferulineae based on the coumarins identified by ¹H-NMR. *J. Plant Resour. & Environ.*, **8**: 46.
- Liu XY and Chang TH (2002). Effect of Pd-Ia on expression of myocardial Fas protein and serum IL-6 level of ischemic reperfusion rats. *Shenyang Yaokexue Xuebao*, **19**: 348.
- Liu XY, Chang TH, Zhang XH and Xing J (2002). Effect of aqua-ethanol extract of *Peucedanum praeruptorum* Dunn on rats with ischemic reperfusion. *Zhongguo Yike Daxue Xuebao*, **31**: 255.
- Lou HX, Sun LR, Yu WT, Fan PH, Cui L, Gao YH, Ma B, Ren DM and Ji M (2004). Absolute configuration determination of angular dihydrocoumarins from *Peucedanum praeruptorum*. *J. Asian Nat. Prod. Res.*, **6**: 177.
- Lu M, Nicoletti M, Battinelli L and Mazzanti G (2001). Isolation of praeruptorins A and B from *Peucedanum*. *Farmaco*, **56**: 417.
- Miyazawa M, Shimamura H, Bhuvra R, Nakamura S and Kameoka H (1996). Antimutagenic activity of falcariindiol from *Peucedanum praeruptorum*. *J. Agric. Food Chem.*, **44**: 3444.
- Okuyama T and Shibata S (1981). Studies on Coumarins of a Chinese Drug "Qian-Hu". *Planta Med.*, **42**: 89.
- Okuyama T, Takata M and Shibata S (1989). Structure of linear furano- and simple-coumarin glycosides of Bai-Hua Qian-Hu. *Planta Med.*, **55**: 64.
- Rao GX, Liu QX, Dai ZJ, Yang Q and Dai WS (1995). Textual research and discussing modern variety of *Peucedanum praeruptorum* Dunn, Chinese herb. *Yunnan Zhongyi Xueyuan Xuebao*, **18**: 1.
- Rao MR, Liu WB, Liu PQ and Wei EH (2002a). Effects of Praeruptorin C on vascular hypertrophy, [Ca²⁺]_i, Collagen. *J. Chin. Pharma. Sci.*, **11**: 110.
- Rao MR, Shen XH and Zou X (1988). Effects of praeruptorin C and E isolated from 'Qian-Hu' on swine coronary artery and guinea-pig atria. *Eur. J. Pharmacol.*, **155**: 293.
- Rao MR, Sun L and Zhang XW (2002b). Effect of praeruptorin coumarin on cardiac mass, myocardial [Ca²⁺]_i and Na⁺,K⁺-ATPase, Ca²⁺,Mg²⁺-ATPase activity in renovascular hypertensive rats. *Yao Xue Xue Bao*, **37**: 401.
- Rao MR, Sun L and Zhang XW (2002c). Effects of praeruptorin coumarin on heart hemodynamics, myocardial compliance and collagen content in heart hypertrophy rats. *Zhongguo Yaoli Xue Yu Duli Xue Zazhi*, **16**: 265.
- Ruan H, Zhang Z, Liang XF, Fu Y, Su MQ, Liu QL, Wang XM and Zhu X (2011). Metabolism of *dl*-praeruptorin A in rat liver microsomes using HPLC-electrospray ionization tandem mass spectrometry. *Arch. Pharm. Res.*, **34**: 1311.
- Song JL and Xing HZ (2000). Identification of Qian-hu using thin-layer chromatography. *Shizhen Guoyi Guoyao*, **11**: 143.
- Song YL, Zhang Qw, Li YP, Yan R and Wang YT (2012a). Enantioseparation and absolute configuration determination of angular-type pyranocoumarins from Peucedani Radix using enzymatic hydrolysis and chiral HPLC-MS/MS analysis. *Molecules*, **17**: 4236.
- Song YL, Jing WH, Zhao HY, Yan R, Li PT and Wang YT (2012b). Stereoselective metabolism of (+/-)-praeruptorin A, a calcium channel blocker from Peucedani Radix, in pooled liver microsomes of rats and humans. *Xenobiotica*, **42**: 231.
- Song YL, Yan R, Jing WH, Zhao HY and Wang YT (2011). Characterization of metabolism of (+)-praeruptorin B and (+)-praeruptorin E in human and rat liver microsomes by liquid chromatography coupled with ion trap mass spectrometry and time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.*, **25**: 719.
- Swager TM and Cardellina JH (1985). Coumarins from *Museion divaricatum*. *Phytochemistry*, **24**: 805.
- Takata M, Okuyama T and Shibata S (1988). Studies on Coumarins of a Chinese Drug, "Qian-Hu"; VIII. Structures of New Coumarin-Glycosides of "Bai-Hua Qian-Hu". *Planta Med.*, **54**: 323.
- Takata M, Shibata S and Okuyama T (1990a). Structures of angular pyranocoumarins of Bai-Hua Qian-Hu, the Root of *Peucedanum praeruptorum*. *Planta Med.*, **56**: 307.
- Takata M, Shibata S and Okuyama T (1990b). Structures of glycosides of Bai-Hua Qian-Hu, the root of *Peucedanum praeruptorum*. *Planta Med.*, **56**: 64.
- Tao YY, Luo JG, Lu YY, Xu DR, Hou ZG and Kong LY (2009). Rapid Identification of two species of *Peucedanum* by high-performance liquid chromatography-diode array detection-electrospray ionization tandem mass spectrometry. *Nat. Prod. Commun.*, **4**: 1079.

- Wang CH (2004). Primary determination of content of coumarin in different species from radix *Peucedani* by HPLC. *China Pharmacist*, **7**: 361.
- Wang DC, Zhang XZ and Feng L (2008). Antioxidative effects of total coumarins from *Peucedanum praeruptorum* Dunn *in vitro*. *Yiyao Daobao*, **27**: 899.
- Wang DC, Zhao XM, Li TD, Ma J, Kong ZK and Gao Ys (2004). Effect of total coumarins from *Peucedanum praeruptorum* Dunn on the activity of hepatic drug-metabolizing enzymes in mice. *Yiyao Daobao*, **23**: 522.
- Wang J, Ma Y, Li W, Hu F, Chen T, Shen X and Feng S (2012). Study on pharmacokinetics and tissue distribution of pteryxin in mice by ultra-pressure liquid chromatography with tandem mass spectrometry. *Biomed. Chromatogr.*, **26**: 802.
- Wang JY, Wang HL, Zhang XH and Xing J (2000). Effects of *Peucedanum praeruptorum* Dunn on the monocrotaline-induced pulmonary hypertension in rats. *Zhongguo Yaoxue Zazhi*, **35**: 899.
- Wang LJ, Li JM, Chang TH, Hao LY, Wang YP and Zhang KY (1995). Effect of *dl*-praeruptorin A on delayed outward potassium current in single ventricular cell of guinea pig. *Zhongguo Yaoli Xue Yu Duli Xue Zazhi*, **9**: 90.
- Wang LJ, Li JM, Chang TH, Wang Y, Meng WF and Zhang KY (1997). The preliminary research of *dl*-praeruptorin A in mechanism of action on delayed outward potassium current. *Zhongguo Yaoli Xue Tongbao*, **13**: 450.
- Wang NH, Baba K and Taniguchi M (1996). Simple and convenient application of ¹H-NMR method in the chemotaxonomy of Apioideae. *J. Plant Resour. & Environ.*, **5**: 40.
- Wang QY, Kang J, Li ER and Yu RJ (2001). Effect of extract from Baihua Qianhu on the plasma endothelin-1 in patients with hypoxic pulmonary hypertension. *Zhongguo Yike Daxue Xuebao*, **30**: 28.
- Wang XB, Shi YR and Kong LY (2006). (+)-(3'R)-Angeloxylxy-4'-oxo-3',4'-dihydroseselin (Pd-Ib). *Acta. Crystallogr. E.*, **62**: 2505.
- Wei MJ, Zhang XH and Zhao NC (1994). Effects of whiteflower Hogfennel (*Peucedanum praeruptorum*) on isolated rabbit pulmonary arteries. *Zhong Cao Yao*, **25**: 137.
- Willette R and Soine T (1962). Coumarins I. Isolation, purification, and structure determination of pteryxin and suksdorfins. *J. Pharma. Sci.*, **51**: 149.
- Wu JYC, Fong WF, Zhang JX, Leung CH, Kwong HL, Yang MS, Li D and Cheung HY (2003). Reversal of multidrug resistance in cancer cells by pyranocoumarins isolated from Radix *Peucedani*. *Eur. J. Pharmacol.*, **473**: 9.
- Wu MY, Ye JS and Liu QX (1996). Identification and studies on chemotaxonomy of Chinese crude drug "Qian-hu" by NMR spectroscopy. *J. Plant Resour. & Environ.*, **5**: 14.
- Wu WL, Chen JJ, Liu SJ, Fang CW and Zhang QQ (2009). Simultaneous determination of three coumarin constituents in roots of *Peucedanum praeruptorum* by RP-HPLC. *Zhongguo Zhong Yao Za Zhi.*, **34**: 1121.
- Xiang JR (2006). Study on bionics wild cultivation technique of Ning-guo *Peucedanum praeruptorum*. *Xiandai Zhongyao Yanjiu Yu Shijian*, **20**: 18.
- Xiong Y, Wang J, Wu F, Li J, Zhou L and Kong L (2012a). Effects of (+/-)-praeruptorin A on airway inflammation, airway hyperresponsiveness and NF-kappaB signaling pathway in a mouse model of allergic airway disease. *Eur. J. Pharmacol.*, **683**: 316.
- Xiong YY, Wang JS, Wu FH, Li J and Kong LY (2012b). The effects of (+/-)-Praeruptorin A on airway inflammation, remodeling and transforming growth factor-beta1/Smad signaling pathway in a murine model of allergic asthma. *Int. Immunopharmacol.*, **14**: 392.
- Xiong YY, Wu FH, Wang JS, Li J and Kong LY (2012c). Attenuation of airway hyperreactivity and T helper cell type 2 responses by coumarins from *Peucedanum praeruptorum* Dunn in a murine model of allergic airway inflammation. *J. Ethnopharmacol.*, **141**: 314.
- Xu GB, Liu SJ, Liang YM, Fang CW, Yu NJ and Peng DY (2007). Determination of the content of praeruptorin C in radix *Peucedani* using HPLC. *Zhongguo Zhong Yao Za Zhi*, **32**: 2069.
- Xu Q, Deng LD and Liu BM (2001a). Determination of Pd-Ia and Pd-II in *Peucedanum praeruptorum* by RP-HPLC. *Huaxi Yaoxue Zazhi*, **16**: 215.
- Xu Q, Liu BM and Zhang ZX (2001b). Determination of Pd-Ia and Pd-II in *Peucedanum praeruptorum* by GC. *Zhongguo Yaoxue Zazhi*, **36**: 122.
- Xu Z, Wang X, Dai Y, Kong L, Wang F, Xu H, Lu D, Song J and Hou Z (2010). (+/-)-Praeruptorin A enantiomers exert distinct relaxant effects on isolated rat aorta rings dependent on endothelium and nitric oxide synthesis. *Chem. Biol. Interact.*, **186**: 239.
- Yang J and Rao Mr (1997). Calcium-antagonistic effects of praeruptorin C on tailartery in two models of hypertensive rats. *Zhongguo Yaoli Xue Tongbao*, **13**: 216.
- Yang JR, Li QP and Rao MR (1995). Effects of praeruptorin C on hemodynamics of isolated rat heart with ischemia-reperfused damage. *Zhongguo Yaoli Xue Yu Duli Xue Zazhi*, **9**: 271.
- Ye JS, Wu MY and Liu QX (1995). Identification and chemotaxonomy of 3 Chinese crude drug "Qian-hu" by NMR spectroscopy. *J. Plant Resour. & Environ.*, **4**: 61.
- Ye JS, Zhang HQ and Yuan CQ (1982). Isolation and identification of coumarin praeruptorin E from the root of the Chinese drug *Peucedanum praeruptorum* Dunn (Umbelliferae). *Yao Xue Xue Bao*, **17**: 431.
- YEE, S (1997). *In vitro* permeability across Caco-2 cells (colonic) can predict *in vivo* (small intestinal) absorption in man--fact or myth. *Pharm Res*, **14**, 763.
- Yu PJ, Ci W, Wang GF, Zhang JY, Wu SY, Xu W, Jin H, Zhu ZG, Zhang JJ, Pang JX and Wu SG (2011).

- Praeruptorin A inhibits lipopolysaccharide-induced inflammatory response in murine macrophages through inhibition of NF-kappaB pathway activation. *Phytother. Res.*, **25**: 550.
- Yu PJ, Jin H, Zhang JY, Wang GF, Li JR, Zhu ZG, Tian YX, Wu SY, Xu W, Zhang JJ and Wu SG (2012). Pyranocoumarins isolated from *Peucedanum praeruptorum* Dunn suppress lipopolysaccharide-induced inflammatory response in murine macrophages through inhibition of NF-kappaB and STAT3 activation. *Inflammation*, **35**: 967.
- Yue WY, Liang TG, Du X and Li QS (2012). The absorption and transport characteristics of praeruptorin in caco-2 cell model. *Zhongguo Yaowu Yu Linchuang*, **12**: 851.
- Zhang C, Li L, Geng LD, Liu YY, Lin N and Xiao YQ (2005a). Studies on quality standard of *Peucedanum praeruptorum*. *Zhongguo Yaowu Zazhi*, **30**: 177.
- Zhang C, Li L, Xiao YQ, Li W, Yin XJ, Tian GF, Chen DD and Wang Y (2010). A new phenanthraquinone from the roots of *Peucedanum praeruptorum*. *Chin. Chem. Let.*, **21**: 816.
- Zhang C, Xiao YQ, Taniguchi M and Baba K (2005b). Studies on chemical constituents in roots of *Peucedanum praeruptorium*. *Zhongguo Zhong Yao Za Zhi*, **30**: 675.
- Zhang C, Xiao YQ, Taniguchi M and Baba K (2006). Studies on chemical constituents from the roots of *Peucedanum praeruptorum* II. *Zhongguo Zhong Yao Za Zhi*, **31**: 1333.
- Zhang JX, Fong WF, Wu JY, Yang MS and Cheung HY (2003). Pyranocoumarins Isolated from *Peucedanum praeruptorum* as Differentiation Inducers in Human Leukemic HL-60 Cells. *Planta Med.*, **69**: 223.
- Zhang SL, Li JM, Xiao QH, Wu AH, Zhao Q, Yang GR and Zhang KY (2001). Effect of *dl*-praeruptorin A on ATP sensitive potassium channels in human cortical neurons. *Acta. Pharmacol. Sin.*, **22**: 813.
- Zhang Z, Liang XF, Su MQ, Yang Q, Li LP, Zhang XH, Wang XM and Zhu X (2011). Pharmacokinetics of *dl*-praeruptorin A after single-dose intravenous administration to rats with liver cirrhosis. *DARU*, **19**: 210.
- Zhao B, Yang XB, Yang XW, Wu Q, Wang Y, Zhang LX and Xu W (2011). Intestinal permeability of the constituents from the roots of *Saposhnikovia divaricata* in the human Caco-2 cell monolayer model. *Planta Med.*, **77**: 1531.
- Zhao NC, Wang HL, Jin X, Zhang XH, Liu KJ and Guan FL (1994). The effects of petroleum ether extract of *Peucedanum praeruptorum* Dunn (Bai-Hua Qian-Hu) on hypoxic pulmonary hypertension in dogs. *Zhongguo Yike Daxue Xuebao*, **23**: 197.
- Zhou R, Wang HL and Zhang XH (2001a). Effects of PPD on hemorrheology, hemodynamics of pulmonary circulation and correlation analysis in pulmonary hypertension in rats. *Zhongguo Xiandai Yingyong Yaowu Zazhi*, **18**: 263.
- Zhou R, Wang HL, Zhang XH, Xing J and Li CM (2001b). Effects of *Peucedanum* Dunn on hemorrheology and hemodynamics in pulmonary circulation of pulmonary hypertensive rats. *Zhongguo Yike Daxue Xuebao*, **30**: 325.
- Zhou Y (2008). Identification of Radix *Peucedani* and its adulterant Radix *Peucedani* Terebinthaceum. *Zhongguo Yaoye*, **17**: 62.
- Zhu GY, Chen GY, Li QY, Shen XL and Fang HX (2004). HPLC/MS/MS Method for Chemical Profiling of Radix *Peucedani* (Baihua Qianhu). *Chin. J. Nat. Med.*, **2**: 304.