

Factors related to good death in the Eastern Mediterranean Region: a systematic review

Armin Fereidouni,¹ Maryam Rassouli,² Toktam Kianian,³ Maryam Elahikhah,⁴ Hossein Souri,⁴ Shilan Azizi⁵ and Salman Barasteh^{6,7}

¹Department of Operating Room Technology, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. ²Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Islamic Republic of Iran. ³Department of Community Health and Geriatrics, School of Nursing and Midwifery, Iran University of Medical Sciences, Tehran, Islamic Republic of Iran. ⁴Student Research Committee, Baqiyatallah University of Medical Sciences, Tehran, Islamic Republic of Iran. ⁵Department of Nursing, Bandar Abbas Branch, Islamic Azad University, Bandar Abbas, Islamic Republic of Iran. ⁶Health Management Research Center, Baqiyatallah University of Medical Sciences, Tehran, Islamic Republic of Iran (Correspondence: S. Barasteh: s.barasteh@gmail.com). ⁷Nursing Faculty, Baqiyatallah University of Medical Sciences, Tehran, Islamic Republic of Iran.

Abstract

Background: Considerations for palliative care and quality of death has significantly increased over the past 10 years in the Eastern Mediterranean Region (EMR). Recent trends in ageing and increasing chronic disease burden have drawn attention to the need to pay attention to the concept of good death and related factors from the perspective of the local population.

Aims: To assess the factors related to good death in the EMR.

Methods: We searched PubMed, Embase, Scopus, Web of Science, and ProQuest on 22 October 2021 for English language articles, with no time limit, using keywords “quality of death”, “good death”, “quality of dying”, “good dying”, “Middle Eastern”, and countries in the Region. The quality of articles was evaluated using the Hawker criterion and based on the PRISMA guidelines. From the thematic analysis, the factors influencing good death were extracted. EndNote X8 software was used for data management.

Results: The search yielded 55 articles, and 14 were included in the study, with a total of 3589 participants. Factors related to good death were classified into 2 main categories: patient preferences and end-of-life care. The former was divided into 4 groups: symptom management, psychological support, social support, and spiritual care. The second category included 2 subcategories: death control and patient autonomy, and end-of-life care.

Conclusion: Although patients’ beliefs about good death are personal, unique, and different, perception about good death in the EMR depends on the extent to which patients’ preferences are met and end-of-life care is provided. More research on good death is recommended in the context of Islamic countries in EMR, and to empower patients and their families to better manage the dying process and create educational programmes.

Keywords: palliative care, end-of-life, good death, Eastern Mediterranean Region

Citation: Fereidouni A; Rassouli M; Kianian T; Elahikhah M; Souri H; Azizi S; et al. Factors related to good death in the Eastern Mediterranean Region: a systematic review. *East Mediterr Health J.* 2022;28(08):601–609. <https://doi.org/10.26719/emhj.22.069>

Received: 30/08/21; accepted: 01/08/22

Copyright © World Health Organization (WHO) 2022. Open Access. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license (<https://creativecommons.org/licenses/by-nc-sa/3.0/igo>)

Introduction

Death is a natural and inevitable phenomenon (1,2) that plays an important role in all religions, societies and cultures (3). Death reflects the sociocultural identity and belonging of the individual to society (4). People, regardless of their cultural background, view death differently depending on their beliefs, attitudes, fears and aspirations (3). Like most phenomena, death is often experienced well or badly (5). A good death is a multidimensional concept used by various disciplines, including medicine, psychology, theology, sociology and anthropology (6,7). There is no clear and constant definition of what constitutes a good death (4). The concept of a good death was first used in Ancient Greece as a synonym for euthanasia and a condition in which life ended deliberately (8). O’Neil in 1983 described good death as the ability to control one’s independence over death. Weisman defines good death as resolving the

physical, social and emotional concerns of the dying person (9).

A good death is influenced by the culture of communities, religion, and the individual’s experience of death. Some of the factors that affect a good death vary in different societies (4,10). Awareness of death, patient preference, acceptance of death, and patient independence in treatment decisions are highlighted as key factors of good death in western countries (11). In the United States of America and Europe, the concept of good death was widely used in the 1960s and 1970s as a key element for palliative and hospice care (12). American researchers have cited respect for individual independence and open communication between patients and their families (13). Van der Greest in Ghana mentioned death at home and being surrounded by relatives (14), and Ruland and Moore in Norway have expressed painlessness, dignity, respect and calm (2) as the factors that influence good

death. Despite the progress made in different countries to conceptualize good death and related factors, paying attention to this concept in Middle Eastern countries only dates back to the last decade (15–17). The dominant religion in the Middle East is Islam, and death in Islam is described as the “will of God. It is part of the process of life and rebirth. The concept of death is so important in Islam that in the Quran, the word is repeated 84 times (18, 19). Several studies have been conducted on good death in the Middle East. For example, in Turkey, Gurdogan et al. mentioned death as being calm, fulfilling the patient’s wishes, and the presence of loved ones at the time of death (1). However, studies on good death have not been systematically considered, and the factors influencing good death in the Middle East are unclear. As a result, this systematic review was carried out to identify the characteristics that contribute to good death in the Eastern Mediterranean Region (EMR).

Methods

Study design

This systematic review was based on the guidelines of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA), which consists of a 17-item checklist intended to facilitate the preparation and reporting of a robust protocol for systematic review. Detailed, well-described protocols can facilitate understanding and appraisal of the review methods (20). All studies selected were required to specify quality of death or provide data that could be used to calculate this measure. Abstract studies published at conferences, case studies, reviews, historical studies, grey literature, and letters to editors were not included. Participants in the studies were from any race, ethnicity or sex. Embase, PubMed/Medline, Scopus, Web of Science and ProQuest were searched on 22 October 2021 without any time limitation. Keyword selection was a combination of MeSH terms and free text words.

Inclusion criteria

Included articles were original research papers, published in English, without time limitations, and focused on quality of death. Duplicate articles were removed using EndNote X8 software. In the screening phase, the titles and abstracts of the articles were reviewed. Selected studies were divided into 3 categories: relevant, irrelevant and unspecified. Articles classified as irrelevant by 2 researchers were excluded. In the selection phase, the full text of the articles was independently reviewed by 2 researchers. The data were qualitatively analysed and the quality of the studies was assessed by 2 researchers. After reviewing the purpose of the studies and the inclusion criteria, the quality of 14 studies was separately evaluated. The criteria of Hawker et al. were used to evaluate study quality (21). Studies were assessed on a scale ranging from 4 to 1 (4, good; 3, fair; 2, poor; and 1, very poor). The selected articles were analyzed using content analysis. Data were separately extracted

using a researcher-made sheet. At first, a paper was evaluated as a pilot with the sheet; then, it was used for other articles. Any disagreement among the 2 researchers were discussed and resolved.

Identified studies

The search identified 55 articles. EndNote software version 8 was used to organize the information. According to review of the titles and abstracts, 22 duplicate articles were removed. Then, 20 irrelevant articles were removed that were in contrast with the objectives of the study. Finally, the full text of 13 articles was searched. To ensure the retrieval of all articles, the list of references of the final articles were manually searched and 1 more article added to the final total. Fourteen articles that discussed factors affecting good death were finally included (Figure 1 PRISMA flowchart). Based on the quality appraisal criteria of Hawker et al. (21), 10 articles were good quality, and 4 were fair quality; therefore, all articles had appropriate quality (Table 1).

Results

Study characteristics

The 14 studies included a total of 3589 individuals. Table 2 lists the study features, including authors, year of publication, techniques, study population and size, country, age, sex and study goal. Four studies were conducted in 2010–2015, 5 in 2016–2018 and 5 in 2019–2021. Six studies were conducted in Turkey, 5 in the Islamic Republic of Iran, 2 in Saudi Arabia, and 1 in Jordan. Participants included cancer patients, bereaved family members, nurses, social workers, and nursing and medical students. There were 11 cross-sectional, 1 cohort and 2 qualitative studies. After extracting the results of the studies and analysing them, the factors affecting good death were classified into 2: patient preferences and provision of end-of-life care.


Patient preferences

One particularly influential factor in good death was valuing patients’ preferences at end-of-life (24,25). After analysing the study findings, patients’ preferences were categorized into 2: death control and patient autonomy.

Death control

Four studies showed that preplanned death and patient control over what happened in the last days of life influenced good death (15,17,26,27). These studies mentioned factors such as saying goodbye to loved ones (15,17,27), knowing the time of death, dying after a long life without unfulfilled dreams, and doing all the things one loves (3,17,27). The choice of preferred place of death was another factor related to good death (3,16,17,28,29). Three studies showed that home was the preferred place of death (3,17,29). Three studies reported access to hospice care in the last days of life as affecting good death (15,17,27).

Figure 1 PRISMA flowchart



Patient autonomy

Patient autonomy influences good death in different cultures (30). Three studies reported patients' reluctance to impose suffering and hardship on others (3,16,27). Two studies reported disease awareness and the ability to easily access medical professionals (17,31). Death without depending on others was mentioned in 1 study (3). Five studies reported the patients' ability to access advance care directives, such as whether or not to use life-sustaining treatment, or do-not-resuscitate orders, and the desire to access euthanasia (15,17,27,32,33).

Provision of end-of-life care

Provision of end-of-life care had 4 subcategories: symptom management, psychological support, social support, and spiritual care.

Symptom management

Symptom management affects good death in different cultures (4), and includes pain relief (3,17,31).

Psychological support

The emphasis on psychological and social aspects influences good death worldwide (4). Three studies mentioned psychological comfort, absence of stress, and hope in the last days of life (1,3,16). In 2 studies in Turkey and Saudi Arabia, social workers, patients and caregivers reported patient access to emotional support (15,25). Four studies reported the

Social support

Social support entails helping and supporting others (31). Because cultural values are described as an integrated system of patterns of conduct developed by a group of

individuals in society, there may be cultural disparities in the idea of a good death (4). Respect for the patient's cultural values was mentioned by social workers in Turkey (27), and participants in 4 other studies cited dignity and patient's reputation (15,17,27,34). Respect for the patient's privacy at end-of-life and after death (15, 25) and the efforts of professional caregivers to take care of the appearance and health of patients to maintain the patients' dignity (15). One study mentioned not having pressure on the family regarding the costs of care, death and burial (3). Three studies emphasized having a good relationship with family members (15,16,31), patient satisfaction with their families, family security after death, and the patient's readiness to face death (31).

Spiritual care

Religion and spirituality are essential concepts related to well-being, health, disease and death (4). The participants of 4 studies expressed spiritual care in the last days of life as important for good death (15,17,24,27). A study in Saudi Arabia mentioned the remembrance of God, the ability to utter religious expressions, respect for the religious rites of death (34), and the presence of a clergy at the patient's bedside to pray and ask for forgiveness at end-of-life was mentioned in another study (3). Three studies considered resolution of patient conflicts related to faith and belief before death (25,34) and the ability to ask for forgiveness (3).

Discussion

This review aimed to identify the factors related to good death in the EMR. Paying attention to patient preferences and end-of-life care were mentioned as important factors

Table 1 Assessments of articles included in the review

Author, Year	Abstract and title	Introduction and aims	Method and data	Sampling	Data analysis	Ethics and bias	Results	Transferability generalizability	Implications usefulness	Total	Average	Grade
Moghadam et al. (2019)	4	3	3	2	2	2	3	2	3	24	2.7	Fair
Alawneh et al. (2021)	3	2	3	2	3	3	4	2	3	25	2.8	Fair
Şahin et al. (2017)	3	3	3	2	4	2	4	3	3	25	2.8	Fair
Tayeb et al. (2010)	3	3	4	2	4	1	3	3	3	26	2.9	Fair
Gurdoğan et al. (2020)	3	4	3	2	3	4	4	3	2	28	3.1	Good
Razban et al. (2016)	4	2	4	4	3	3	4	3	3	30	3.3	Good
Duyan et al. (2016)	2	4	4	3	4	2	4	4	4	31	3.4	Good
Ceyhan et al. (2018)	2	4	4	4	3	4	4	3	3	31	3.4	Good
Iranmanesh et al. (2011)	3	4	4	4	4	2	4	4	3	32	3.5	Good
Hammami et al. (2015)	4	4	4	4	4	2	4	3	3	32	3.5	Good
Estebnsari et al. (2017)	4	3	4	4	2	4	4	4	4	33	3.7	Good
Duru Aşiret et al. (2020)	3	4	3	3	4	4	4	4	4	33	3.7	Good
Demir et al. (2017)	4	4	4	4	4	4	4	3	3	34	3.8	Good
Moslemi et al. (2020)	4	4	4	4	4	3	4	3	4	34	3.8	Good
Total	3,2	3,4	3,6	3,1	3,4	2,8	3,8	3,1	3,2	31	3,4	Good

of good death (35). Few studies specifically evaluated good death, and the available research was confined to 4 nations in the Region (Turkey, Islamic Republic of Iran, Jordan, and Saudi Arabia).

Patient preferences are the result of individual experiences that are influenced by multidimensional factors and are important in achieving a good death (4,36–38). Patients in Arabic Islamic societies often prefer to reaffirm their religious beliefs before death, say goodbye to their loved ones, and determine how to hold their funeral (1,3,15). Gerritsen et al. found that spending time with loved ones was important for American families and saying goodbye to loved ones was important for Dutch families (39). Cottrell et al. consider these as coping strategies (40) rooted in religious beliefs (16) and ignoring them can cause patients to feel that they have lost control of their lives and their dignity (37).

In this study, dying in a preferred place was one of the factors influencing good death, and 3 studies reported that the preferred place of death was at home. Many patients in the final stages of life attach importance to the preferred place of death, and this place has a significant impact on their quality of life, death and care (40,41). Similarly, a recent systematic review showed that 55% of patients preferred to die at home (44).

Patients consider death to be good when they die without the feeling of overwhelming and disturbing others. The fact that they no longer have control over their bodies and can no longer carry out their personal and daily activities causes a feeling of despair and worthlessness (45). In this regard, Rodríguez-Pra et al. reported that patients at the end of life are more afraid of dependence and lower self-esteem than pain and physical problems, which can potentially make the dying process more painful (46). For this reason, they

Table 2 Characteristics of the studies included in the review

Author/ year	Country	Study type	Participants (n)	Age (yr), mean (SD)	Sex distribution (%)	Aim
Gurdoğan et al. ¹ 2020	Turkey	Cross-sectional	Family caregivers (182)	44.02 (13.02)	Male: 41.8 female: 58.2	To examine the importance of the concept of a good death and the contributing factors from the perspectives of family caregivers of advanced cancer patients.
Duru Aşiret et al. ²⁶ 2020	Turkey	Cross-sectional	Nursing students (224)	20.7 (1.2)	Male: 69.2 female: 30.8	To identify the relationship between the nursing students' attitudes towards spiritual care and the principles of a good death.
Estebansari et al. ³ 2017	Islamic Republic of Iran	Qualitative	Nursing students (300)	No report	Male: 60 female: 40	To explain the views of Iranians about the elements of good death
Duyan et al. ²⁷ 2015	Turkey	Cross-sectional	Social workers (195)	34 (9.65)	Male: 44 female: 56	To be the basis of discussions and further studies on such sensitive topics in Turkey.
Moslemi et al. ³¹ 2020	Islamic Republic of Iran	Cross-sectional	NICU nurses (130)	34.2 (2.3)	Male: 0 female: 100	To assess the psychometric features of the quality of dying and death questionnaire in NICU nurses in Tehran.
Hammami et al. ³⁴ 2015	Saudi Arabia	Qualitative	Saudi men (120)	32.1 (9.8)	Male: 100 female: 0	To explore Saudi male opinions regarding end-of life priorities.
Ceyhan et al. ²² 2018	Turkey	Cross-sectional	Nurses (122)	31.2 (5.39)	No report	To determine the attitude of nurses regarding the concept of a good death and terminal phase was conducted to determine the effect on patient care.
Demir et al. ¹⁷ 2017	Turkey	Cross-sectional	Nurses (856)	30.49 (6.12)	Male: 7.7 female: 92.3	To determine nurses' perceptions and experiences with futile medical care and their opinions about principles of good death.
Iranmanesh et al. ¹⁶ 2011	Islamic Republic of Iran.	Cross-sectional	Bereaved family members (150)	33	Male: 19 female: 81	To evaluate a good death concept from the Iranian bereaved family members' perspective.
Moghadam et al. ³² 2019	Islamic Republic of Iran.	Cross-sectional	Medical students (152)	22.12 (2.35)	Male: 39.5 female: 60.5	To investigate medical students' viewpoints regarding euthanasia.
Razban et al. ³³ 2016	Islamic Republic of Iran.	Cross-sectional	ICU nurses (104)	No report	Male: 7.1 female: 92.9	To evaluate the attitude of critical care nurses towards life-sustaining treatments in South East Iran.
Şahin et al. ²³ 2017	Turkey	Cross-sectional	Emergency - ICU nurses (140)	32.72 (7.35)	Male: 7.1 female: 86.4	To define and measure attitudes toward good death and death anxiety in nurses working at emergency service and intensive care.
Tayeb A. et al. ¹⁵ 2010	Saudi Arabia	Cross-sectional	(Muslim patients and healthcare providers) (284)	37	Male: 58 female: 42	To review the Future of Health and Care of Older People good death perception to identify and describe other components of the Muslim good death perspective.
Alawneh et al. ²⁹ 2021	Jordan	Retrospective cohort study	Cancer patients (630)	No report	Male: 48 female: 52	To examine the place of death of cancer patients in Jordan and the determinants of home death among this patient group.

NICU = neonatal intensive care unit.

should be involved in care activities, and their preferences should be recognized.

The nature of terminal care provided to dying patients and their families is another determinant of good death. Effective symptom management is a critical part of end-of-life care, which minimizes discomfort and anxiety, enabling patients to safely and comfortably pass away (44,45). Saphire et al. reported that inattention to symptom management reduces the quality of life and emotional well-being at end-of-life (49). Baillie et al. attributed access to specialized palliative care to achieving a peaceful death at the patient's preferred place, thus reducing emotional and economic burdens (50). It seems that in terms of the many benefits of palliative care at end-of-life, it is necessary to do more research and make better use of existing research.

Other factors influencing good death include painless death, patients' ability to make independent decisions, and maintenance of patients' dignity. The perception of good death, to some extent, includes timely and practical preparation for death and life after death, which may in part be related to social and financial issues (51). According to Gurdogan et al., patients at end-of-life suffer from major psychological issues due to fear and uncertainty about the future, unresolved issues, and worries about their loved ones (1). Therefore, the inclusion of psychological support in end-of-life care can help manage this suffering by respecting values and creating a relaxed environment for expressing ideas and feelings (1).

If end-of-life care is comprehensive and can incorporate various dimensions of social support, it can lead to good death by conveying a sense of worth to the patient and reducing their concerns (3,53). Financial support reduces the mental suffering of the patient and family and leads to better well-being at end-of-life (i). Sudore et al. stated that not receiving support and lack of family involvement may deprive patients of quality care

because, in most cases, patients cannot make appropriate decisions at end-of-life (55).

According to Tayeb et al., people in the Middle East believe that death is closely linked to faith and spirituality (15). Spiritual support can be used as part of end-of-life care to alleviate the emotional isolation and stress (56,57). This highlights the necessity to empower the healthcare provider's team, especially nurses, to provide spiritual care and, by employing clergy, to convey comfort to the patient (58). One review showed that family support in the end-of-life care process is an essential dimension to good death.

Modern medical institutions have been preparing and debating end-of-life options, variables affecting good death, and overcoming problems and hurdles to obtaining a good and acceptable death for patients.

Our study had several limitations. The most important limitation was the small number of studies that addressed good death in the EMR. Another limitation of the study was the scattering of studies, such that the studies included patients, their caregivers and healthcare providers.

Conclusion

The current study found that 2 core elements of patient preferences and end-of-life care shape the concept of good death in the EMR. By taking these elements into consideration, patients and healthcare providers can make better decisions at end-of-life. It is thought to be important to provide information about death and organize educational programmes to empower patients and family caregivers to better manage the dying process. Therefore, we suggest the need to have more studies and strategies on good death in the context of Islamic countries in the Region.

Funding: None

Competing interests: None declared.

Facteurs liés à la bonne mort dans la Région de la Méditerranée orientale : revue systématique

Résumé

Contexte : Les considérations relatives aux soins palliatifs et à la qualité du décès ont considérablement augmenté au cours des 10 dernières années dans la Région de la Méditerranée orientale. Les tendances récentes du vieillissement et de l'augmentation du fardeau des maladies chroniques ont mis en évidence la nécessité de prêter attention au concept de bonne mort et aux facteurs associés du point de vue de la population locale.

Objectifs : Évaluer les facteurs liés à la bonne mort dans la Région de la Méditerranée orientale.

Méthodes : Le 22 octobre 2021, nous avons effectué des recherches sur PubMed, Embase, Scopus, Web of Science et ProQuest afin de trouver des articles en anglais, sans limitation de période, en utilisant les mots clés « qualité du décès », « bonne mort », « qualité du processus de la mort », « bien mourir », « Moyen-Orient » et les pays de la Région. La qualité des articles a été évaluée à l'aide des critères utilisées par Hawker et en suivant les directives PRISMA. L'analyse thématique a permis l'extraction des facteurs influençant la bonne mort. Le logiciel EndNote X8 a été utilisé pour la gestion des données.

Résultats : La recherche a produit 55 articles, dont 14 ont été inclus dans l'étude, avec un total de 3589 participants. Les facteurs liés à une bonne mort ont été classés en deux catégories principales : les préférences du patient et

les soins de fin de vie. La première a été divisée en quatre groupes : prise en charge des symptômes, soutien psychologique, soutien social et soins spirituels. La seconde catégorie comprenait deux sous-catégories : contrôle sur la mort et autonomie du patient, ainsi que les soins de fin de vie.

Conclusion : Bien que les croyances des patients sur la bonne mort soient personnelles, uniques et différentes, la perception de la bonne mort dans la Région de la Méditerranée orientale dépend de la mesure dans laquelle les préférences des patients sont satisfaites et les soins de fin de vie fournis. Il est recommandé de mener davantage de recherches sur la bonne mort dans le contexte des pays islamiques dans la Région de la Méditerranée orientale, ainsi que de donner aux patients et à leurs familles les moyens de mieux gérer le processus de la mort et de créer des programmes éducatifs.

العوامل المتعلقة "بالاحتضار في سلام" في إقليم شرق المتوسط: استعراض منهجي

آرمين فريدوني، مريم رسولي، تكتم كيانيان، مريم الهى خواه، حسين سوري، شيلان عزيزي، سلمان برسته

الخلاصة

الخلفية: على مدى السنوات العشر الماضية، زاد الاهتمام بالرعاية الملطفة وظروف الاحتضار زيادة كبيرة في إقليم شرق المتوسط. وقد لفتت الاتجاهات الحديثة في الشيخوخة وتزايد عبء الأمراض المزمنة الانتباه إلى ضرورة الاهتمام بمفهوم "الاحتضار في سلام" والعوامل المتعلقة به من منظور السكان المحليين.

الأهداف: هدفت هذه الدراسة إلى تقييم العوامل المتعلقة "بالاحتضار في سلام" في إقليم شرق المتوسط.

طرق البحث: في يوم 22 أكتوبر/ تشرين الأول 2021 بحثنا في قواعد أبحاث: Web of Science, Scopus, Embase, PubMed, ProQuest، عن مقالات باللغة الإنجليزية، بدون تحديد إطار زمني، باستخدام الكلمات المفتاحية التالية: "quality of death" (ظروف الاحتضار)، أو: "good death" (الاحتضار في سلام)، أو: "quality of dying" (ظروف الاحتضار)، أو: "good dying" (الاحتضار في سلام)، أو: "Middle Eastern" (شرق أوسطي) وأسماء بلدان الإقليم. وقِيم الباحثون المقالات باستخدام معيار هوكر، واستنادًا إلى المبادئ التوجيهية للعناصر الموصى بها لإعداد تقارير الاستعراض المنهجي والتحليلات التلوية (PRISMA). ومن خلال التحليل الموضوعي، استُخرجت العوامل التي تؤثر على "الاحتضار في سلام". وأديرَت البيانات ببرنامج EndNote X8.

النتائج: أسفر البحث بالكلمات الرئيسية عن 55 مقالة، وأدرجت 14 منها في الدراسة، ولقد بلغ مجموع المشاركين فيها 3589 مشاركًا. وصُنفت العوامل المتعلقة "بالاحتضار في سلام" إلى فئتين رئيسيتين: تفضيلات المرضى والرعاية في مرحلة الاحتضار. وبعد ذلك قُسمت الفئة الأولى إلى 4 مجموعات، هي: علاج الأعراض، والدعم النفسي، والدعم الاجتماعي، والرعاية الروحية. أما الفئة الثانية فقد اندرجت تحتها فئتان فرعيتان: التحكم في مرحلة الاحتضار واستقلالية المريض، والرعاية في مرحلة الاحتضار.

الاستنتاجات: على الرغم من أن اعتقادات المرضى بخصوص "الاحتضار في سلام" اعتقادات شخصية ومتفردة ومختلفة، فإن فكرة الناس في إقليم شرق المتوسط عن "الاحتضار في سلام" تعتمد على مدى تحقيق تفضيلات المرضى وتوفير الرعاية في مرحلة الاحتضار. ويُوصى بإجراء المزيد من البحوث عن "الاحتضار في سلام" في البلدان الإسلامية في إقليم شرق المتوسط، وتمكين المرضى وأسرها من إدارة مرحلة الاحتضار على نحو أفضل وإعداد برامج للتوعية في هذا الصدد.

References

1. Gurdogan EP, Aksoy B, Kinici E. The concept of a good death from the perspectives of family caregivers of advanced cancer patients. *Omega*. 2020 Jun;85(2):303–16. DOI: 10.1177/0030222820945082 PMID: 32703073
2. Ruland CM, Moore SM. Theory construction based on standards of care: a proposed theory of the peaceful end of life. *Nurs Outlook*. 1998 Jul–Aug;46(4):169–75. PMID: 9739534 DOI: 10.1016/s0029-6554(98)90069-0
3. Estebarsari F, Taghdisi MH, Mostafaei D, Rahimi Z. Elements of healthy death: A thematic analysis. *Med J Islam Repub Iran*. 2017 Mar 23;31:24. DOI: 10.18869/mjiri.31.24 PMID: 29445653
4. Krikorian A, Maldonado C, Pastrana T. Patient's perspectives on the notion of a good death: a systematic review of the literature. *J Pain Symptom Manage*. 2020 Jan;59(1):152–64. DOI: 10.1016/j.jpainsymman.2019.07.033 PMID: 31404643
5. Scarre G. Can there be a good death? *J Eval Clin Pract*. 2012 Oct;18(5):1082–6. DOI: 10.1111/j.1365-2753.2012.01922.x PMID: 22995013
6. Counts DA, Counts D. The good, the bad, and the unresolved death in Kaliai. *Soc Sci Med*. 2004 Mar;58(5):887–97. DOI: 10.1016/j.socscimed.2003.10.040 PMID: 14732603
7. Flaskerud JH. Individual and dynamic: western views of a good death. *Issues Ment Health Nurs*. 2017 Oct;38(10):865–8. DOI: 10.1080/01612840.2017.1295492 PMID: 28350477
8. Kehl KA. Moving toward peace: an analysis of the concept of a good death. *Am J Hosp Palliat Med*. 2006;23(4):277–86.

9. Weisman AD. Appropriate death and the hospice program. *Hosp J*. 1988;4(1):65–77. DOI: 10.1080/0742-969x.1988.11882615 PMID: 3391538
10. Kastbom L, Milberg A, Karlsson M. A good death from the perspective of palliative cancer patients. *Support Care Cancer*. 2017 Mar;25(3):933–9. DOI: 10.1007/s00520-016-3483-9 PMID: 27837324
11. Kuczmarski TM, Odejide OO. Goal of a “good death” in end-of-life care for patients with hematologic malignancies - are we close? *Curr Hematol Malig Rep*. 2021 Apr;16(2):117–25. DOI: 10.1007/s11899-021-00629-1 PMID: 33864180
12. Saunders Y, Ross J, Riley J. Planning for a good death: responding to unexpected events. *BMJ*. 2003 Jul 26;327(7408):204–6. DOI: 10.1136/bmj.327.7408.204 PMID: 12881264
13. Winland-Brown J. John and Mary Q. Public’s perceptions of a good death and assisted suicide. *Issues Interdiscip Care*. 2001;3(2):137–44.
14. Van der Geest S. Dying peacefully: considering good death and bad death in Kwahu-Tafo, Ghana. *Soc Sci Med*. 2004 Mar;58(5):899–911. DOI: 10.1016/j.socscimed.2003.10.041 PMID: 14732604
15. Tayeb MA, Al-Zamel E, Fareed MM, Abouellail HA. A “good death”: Perspectives of Muslim patients and health care providers. *Ann Saudi Med*. 2010 May–Jun;30(3):215–21. DOI: 10.4103/0256-4947.62836 PMID: 20427938
16. Iranmanesh S, Hosseini H, Esmaili M. Evaluating the “good death” concept from Iranian bereaved family members’ perspective. *J Support Oncol*. 2011 Mar–Apr;9(2):59–63. DOI: 10.1016/j.suponc.2010.12.003 PMID: 21542412
17. Demir A, Sançar B, Yazgan EÖ, Özcan S, Duyan V. Intensive care and oncology nurses’ perceptions and experiences with ‘futile medical care’ and ‘principles of good death’. *Turk Geriatri Dergisi*. 2017;20(2):116–24.
18. Zahedi F, Larijani B, Tavakoly J. End of life ethical issues and Islamic views. *Iran J Allergy Asthma Immunol*. 2007 Feb;6(Suppl. 5):5–15
19. Rassouli M, Sajjadi M. Cancer care in countries in transition: the Islamic Republic of Iran. *Springer*; 2016:317–36.
20. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015 Jan 1;4(1):1–9. DOI: 10.1186/2046-4053-4-1 PMID: 25554246
21. Hawker S, Payne S, Kerr C, Hardey M, Powell J. Appraising the evidence: reviewing disparate data systematically. *Qual Health Res*. 2002 Nov;12(9):1284–99. DOI: 10.1177/1049732302238251 PMID: 12448672
22. Ceyhan Ö, Özen B, Zincir H, Şimşek N, Başaran M. How intensive care nurses perceive good death. *Death Stud*. 2018 Nov–Dec;42(10):667–72. DOI: 10.1080/07481187.2018.1430084 PMID: 29393820
23. Sahin DS, Onal O, Inanc BB. Attitudes of nurses working emergency and intensive care units toward good death and death anxiety. *J Clin Anal Med*. 2017;8(Suppl. 2):75–9.
24. Waller A, Sanson-Fisher R, Nair BR, Evans T. Preferences for end-of-life care and decision making among older and seriously ill inpatients: a cross-sectional study. *J Pain Symptom Manage*. 2020 Feb;59(2):187–96. DOI: 10.1016/j.jpainsymman.2019.09.003 PMID: 31539600
25. Sanjo M, Miyashita M, Morita T, Hirai K, Kawa M, Akechi T, et al. Preferences regarding end-of-life cancer care and associations with good-death concepts: a population-based survey in Japan. *Ann Oncol*. 2007 Sep;18(9):1539–47. DOI: 10.1093/annonc/mdm199 PMID: 17660496
26. Duru Aşiret G, Kütmeç Yılmaz C, Gökşin İ. Relationship between the nursing students’ attitudes towards spiritual care and the principles of a good death. *Perspect Psychiatr Care*. 2020 Oct;56(4):913–9. DOI: 10.1111/ppc.12511 PMID: 32285477
27. Duyan V, Serpen AS, Duyan G, Yavuz S. Opinions of social workers in Turkey about the principles on die with dignity. *J Relig Health*. 2016 Dec;55(6):1938–53. DOI: 10.1007/s10943-015-0090-x PMID: 26239506
28. Ali M, Capel M, Jones G, Gazi T. The importance of identifying preferred place of death. *BMJ Support palliat Care*. 2019 Mar;9(1):84–91. DOI: 10.1136/bmjspcare-2015-000878 PMID: 26408428
29. Alawneh A, Anshasi H. Place of death for patients treated at a tertiary cancer center in Jordan. *Support Care Cancer*. 2021 Apr;29(4):1837–42. DOI: 10.1007/s00520-020-05677-6 PMID: 32779008
30. Cheng SY, Lin WY, Cheng YH, Huang CH, Chen CY, Hwang SJ, et al. Cancer patient autonomy and quality of dying—a prospective nationwide survey in Taiwan. *Psychooncology*. 2016 Feb;25(2):179–86. DOI: 10.1002/pon.3901 PMID: 26179714
31. Moslemi M, Nikfarid L, Nourian M, Nasiri M, Rezayi F. Translation, cultural, and age-related adaptation and psychometric properties of Persian version of “quality of dying and death” in nurses working in neonatal intensive care units. *Indian J Palliat Care*. 2020 Jan–Mar;26(1):34–9. DOI: 10.4103/IJPC.IJPC_119_19 PMID: 32132781
32. Moghadam HM, Nakhaee S, Talebi M, Ataei AA, Mehrpour M. Iranian medical students’ attitude towards euthanasia. *Indian J Forensic Med Toxicol*. 2019;13(4):612–7.
33. Razban F, Iranmanesh S, Aliabadi HE, Forouzi MA. Critical care nurses’ attitude towards life-sustaining treatments in South East Iran. *World J Emerg Med*. 2016;7(1):59–64. DOI: 10.5847/wjem.j.1920-8642.2016.01.011 PMID: 27006741
34. Hammami MM, Al Gaai E, Hammami S, Attala S. Exploring end of life priorities in Saudi males: usefulness of Q-methodology. *BMC Palliat Care*. 2015 Nov 26;14(1):1–16. DOI: 10.1186/s12904-015-0064-5 PMID: 26611147
35. Barasteh S, Parandeh A, Rassouli M, Zaboli R, Vahedian-Azimi A, Khaghanizadeh M. Integration of palliative care into the primary health care of Iran: a document analysis. *Mid East J Cancer*. 2021;12(2):292–300.

36. Hou X, Lu Y, Yang H, Guo R, Wang Y, Wen L, et al. Preferences for a good death: a cross-sectional survey in advanced cancer patients. *BMJ Support Palliat Care*. 2019. <https://doi.org/10.1136/bmjspcare-2018-001750>
37. Meier EA, Gallegos JV, Thomas LPM, Depp CA, Irwin SA, Jeste DV. Defining a good death (successful dying): literature review and a call for research and public dialogue. *Am J Geriatr Psychiatry*. 2016 Apr;24(4):261–71. DOI: 10.1016/j.jagp.2016.01.135 PMID: 26976293 PMCID: PMC4828197
38. Gerritsen RT, Koopmans M, Hofhuis JG, Curtis JR, Jensen HI, Zijlstra JG, et al. Comparing quality of dying and death perceived by family members and nurses for patients dying in US and Dutch ICUs. *Chest*. 2017 Feb;151(2):298–307. DOI: 10.1016/j.chest.2016.09.003 PMID: 27660153
39. Cottrell L, Duggleby W. The “good death”: an integrative literature review. *Palliat Support Care*. 2016 Dec;14(6):686–712. DOI: 10.1017/S1478951515001285 PMID: 26732508
40. van Gorp J, Soyannwo O, Odeunmi K, Dania S, van Selm M, van Leeuwen E, et al. Telemedicine’s potential to support good dying in Nigeria: a qualitative study. *PLoS One*. 2015 Jun 1;10(6):e0126820. DOI: 10.1371/journal.pone.0126820 PMID: 26030154
41. Minamiguchi Y. Decision-making about the place of death for cancer patients: a concept analysis. *Asia J Oncol Nurs*. 2019 Dec 5;7(1):103–12. DOI: 10.4103/apjon.apjon_38_19 PMID: 31879691
42. Skorstengaard MH, Neergaard MA, Andreassen P, Brogaard T, Bendstrup E, Lokke A, et al. Preferred place of care and death in terminally ill patients with lung and heart disease compared to cancer patients. *J Palliat Med*. 2017 Nov;20(11):1217–24. DOI: 10.1089/jpm.2017.0082 PMID: 28574737
43. Fereidouni A, Rassouli M, Salesi M, Ashrafzadeh H, Vahedian-Azimi A, Barasteh S. Preferred place of death in adult cancer patients: a systematic review and meta-analysis. *Front Psychol*. 2021 Aug 27;12:704590. DOI: 10.3389/fpsyg.2021.704590 PMID: 34512460
44. Östlund U, Brown H, Johnston B. Dignity conserving care at end-of-life: a narrative review. *Eur J Onco Nurs*. 2012 Sep;16(4):353–67. DOI: 10.1016/j.ejon.2011.07.010 PMID: 21917517
45. Rodríguez-Prat A, Monforte-Royo C, Porta-Sales J, Escribano X, Balaguer A. Patient perspectives of dignity, autonomy and control at the end of life: systematic review and meta-ethnography. *PLoS One*. 2016 Mar 24;11(3):e0151435. DOI: 10.1371/journal.pone.0151435 PMID: 27010323
46. Albert RH. End-of-life care: managing common symptoms. *Am Fam Physician*. 2017 Mar 15;95(6):356–61. PMID: 28318209
47. Saphire ML, Prsic EH, Canavan ME, Wang S-YJ, Presley CJ, Davidoff AJ. Patterns of symptom management medication receipt at end-of-life among medicare beneficiaries with lung cancer. *J Pain Symptom Manage*. 2020 Apr;59(4):767–77.e1. DOI: 10.1016/j.jpainsymman.2019.11.015 PMID: 31778783
48. Baillie J, Anagnostou D, Sivell S, Van Godwin J, Byrne A, Nelson A. Symptom management, nutrition and hydration at end-of-life: a qualitative exploration of patients’, carers’ and health professionals’ experiences and further research questions. *BMC Palliat Care*. 2018 Apr 16;17(1):60. DOI: 10.1186/s12904-018-0314-4 PMID: 29656713
49. Fu C, Glasdam S. The ‘good death’ in Mainland China—a scoping review. *Int J Nurs Stud Adv*. 2022 Dec;4. <https://doi.org/10.1016/j.ijnsa.2022.100069>
50. Fan S-Y, Lin I-M, Hsieh J-G, Chang C-J. Psychosocial care provided by physicians and nurses in palliative care: a mixed methods study. *J Pain Symptom Manage*. 2017 Feb;53(2):216–23. DOI: 10.1016/j.jpainsymman.2016.08.019 PMID: 27756622
51. Dobříková P, Macková J, Pavelek L, AlTurabi LK, Miller A, West DJ. The effect of social and existential aspects during end of life care. *Nur Palliat Care*. 2016;1(3):47–51. DOI: 10.15761/NPC.1000113
52. Sudore RL, Casarett D, Smith D, Richardson DM, Ersek M. Family involvement at the end-of-life and receipt of quality care. *J Pain Symptom Manage*. 2014 Dec;48(6):1108–16. DOI: 10.1016/j.jpainsymman.2014.04.001 PMID: 24793077
53. Definition of palliative care. Geneva: World Health Organization; 2017 (<http://www.who.int/cancer/palliative/definition/en/2017>, accessed 14 August 2017).
54. Batstone E, Bailey C, Hallett N. Spiritual care provision to end-of-life patients: a systematic literature review. *J Clin Nurs*. 2020;29(19–20):3609–24. <https://doi.org/10.1111/jocn.15411>
55. Khorrami Markani A, Vahidi M. [Spiritual care of end of life patients in Iran: a neglected nursing practice]. *Nurs Midwifery J*. 2020;17(10):767–70 (in Persian).