

# Risk assessment of road traffic accidents related to sleepiness during driving: a systematic review

Shehzad Saleem<sup>1</sup>

<sup>1</sup>Department of Community Medicine, King Edward Medical University, Lahore, Punjab, Pakistan (Correspondence to: S. Saleem: drshehzad786@gmail.com).

## Abstract

**Background:** Injuries due to accidental crash are the 8th leading cause of death worldwide. Sleepiness results in disrupted neurological function and is a major risk factor for road traffic accidents.

**Aims:** This systematic review assessed the relationship between sleepiness during driving and road traffic accidents.

**Methods:** A systematic review was conducted using online databases such as Wiley Online Library, JSTOR, Medline, and PubMed. Full-text, English language articles published between May 2000 and November 2020 were retrieved. Road traffic accident was set as the outcome of interest and sleepiness during driving as the exposure. The review included studies containing adjusted risk estimates (95% confidence interval). Ten cross-sectional studies (N = 55,945), 5 case-control studies (N = 3821), and 2 cohort studies (N = 16,875) were included.

**Results:** Over 50% of the participants in the different studies experienced sleep deprivation ranging from 3.5% to 67.3%. Abe et al. reported the highest (58%) frequency of sleepiness during driving in their cross-sectional study in Japan, and Nabi et al. reported the lowest (1.1%) in their cohort study in France.

**Conclusion:** Sleepiness and sleep deprivation were related to road traffic accidents; and sleep deprivation was the main contributor to drowsiness while driving.

Keywords: road traffic accidents, sleepiness, driving, observational study, risk assessment

Citation: Saleem S. Risk assessment of road traffic accidents related to sleepiness during driving: a systematic review. *East Mediterr Health J.* 2022;28(9):695–700. <https://doi.org/10.26719/emhj.22.055>

Received: 09/10/21; accepted: 11/05/22

Copyright © World Health Organization (WHO) 2022. Open Access. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license (<https://creativecommons.org/licenses/by-nc-sa/3.0/igo>)

## Introduction

Sleep is a dynamic process that affects the way our bodies function (1). Sleepiness can be defined as difficulty remaining awake even while carrying out activities (2). Sleep deprivation is defined as a state caused by inadequate quantity or quality of sleep, including voluntary or involuntary sleeplessness and circadian rhythm sleep disorders (3). Sleepiness results in disrupted brain functioning, such as reduced reaction time or decreased ability for decision-making. It is a major contributor to road traffic accidents, which often occur when a driver experiences drowsiness at the wheel, or due to sleep abnormalities, lack of sleep, alcohol consumption or medication (4). About 1.3 million deaths occur each year as a result of road traffic accidents globally, causing a 3% loss of the gross domestic product of most countries (5). The US National Highway Traffic Safety Administration has estimated that worldwide every year, about 100 000 road accidents are caused by drowsiness, accounting for > 1500 deaths and > 70,000 injuries (6).

In every country, road traffic accidents are a major public health problem and cause huge societal and financial burdens (7). Sleepiness causes disruption of neurological functions (8,9). Factors that contribute to the incidence of road traffic accidents range from continued driving even when feeling drowsy, having a physical condition, fewer sleeping hours, more working hours,

and nutritional imbalances (10). Several studies during the last 20 years have suggested that sleepiness is among the main factors that cause road traffic accidents (11–15). Sleepiness while driving contributes to 3% to > 30% of all road traffic accidents globally (16–18), which may involve a variety of sleep conditions but also may be caused by sleep deprivation (19–20). More than 20% of the drivers feel a need to stop driving at least once due to sleepiness (21). A religious lifestyle was found to be negatively associated with the risk of road traffic accidents, as were younger drivers (22,23). This systematic review was designed for a better understanding of the relationship between sleepiness and risk of road traffic accidents.

## Methods

### Study protocol

This systematic review protocol was developed keeping in view the requirements of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and guidelines 2020, without execution of a meta-analysis. The PRISMA statement is a guideline developed by an international group of 29 methodologists and experienced researchers in 2005. It comprises 27 checklists and 4-phase flow diagram to ensure transparent reporting of a systematic review [24].

## Search strategy

The reviewer searched for articles published between January 2000 and December 2020 in Wiley Online Library, JSTOR, Medline and PubMed databases. The keywords used were: road traffic accidents, sleepiness while driving, and observational study. PRISMA information flow during the phases of this systematic review is presented in Figure 1.

## Search eligibility criteria

The reviewer included observational studies with adjusted risk estimates and outcome measure of road traffic accidents. Due to expected difficulties of quantification, excluded studies were experimental studies, case series without comparison groups and case reports.

## Data extraction


The potentially relevant articles were screened by title and abstract and full text of the articles were retrieved from databases. Final eligibility was assessed independently keeping in view the PRISMA guideline checklist 2020, and the reviewer extracted the following details from the included studies: study design (cross-sectional, case-control and cohort), number of road accidents, frequency of sleepiness while driving, and the adjusted risk estimates of accidents resulting from sleepiness. For quality assessment purposes, the Newcastle Ottawa Scale was used with a total score for quality ranging from 0 to

9 (25). Participants' characteristics such as age, gender, and sleep disorders and lack of sleep due to workload/ other causes were extracted. The overall frequency of sleepiness during driving was calculated using adjusted risk estimates.

## Study selection

A total of 5651 articles were identified from Wiley Online Library, JSTOR, Medline and PubMed databases, where sleepiness while driving was among the causes of road traffic accidents. A total of 1132 duplicates (20.03%) were removed including 717 (63.33%) from JSTOR, and 415 from Medline and PubMed (36.66%). Out of the 4519 remaining articles (79.96%), 4283 were found to be unrelated (94.77%) and were excluded. The remaining 236 articles (5.22%) were fully reviewed and this led to a total of 17 observational studies (7.20%) that qualified for the systematic review. The 17 eligible studies consisted of a total of 76,641 participants worldwide (Table 1). There were 10 cross-sectional studies (55,945 participants; 72.99%); 5 case-control studies (3821 participants; 4.98%), and 2 cohort studies (16,875 participants; 22.01%). Sixteen studies included both sexes while 2 Saudi Arabian studies were based on men due to the previous ban on female drivers. The ages of participants were < 30 years in 3 studies, 30–50 years in 8 studies, and > 50 years in 6 studies.

Figure 1 PRISMA flow data



**Table 1** Estimates of risk associated with sleepiness during driving

Study type (ref)	Country	Sample size (accidents)	Age, yr	Women	Sleepiness during driving (accidents)	Frequency of sleepiness during driving	Sleep disorders	Sleep deprivation	OR (95% CI)
Cross-sectional (10)	Japan	2462 (21)	30–50	22.0%	1429 (20)	58.0	20.5	42.8	12.90 (1.72–97.69)
Cross-sectional (2)	Saudi Arabia	1219 (773)	30–50	0	307 (228)	25.1	1.1	67.3	1.19 (0.85–1.67)
Case-control (4)	New Zealand	1159 (571)	<30	36.6%	71 (63)	6.1	1.4	9.0	8.20 (3.40–19.70)
Case-control (9)	USA	399 (114)	30–50	31.0%	158 (NR)	39.5	38.8	15.0	1.60 (1.0–2.70)
Cross-sectional (16)	New Zealand	5368 (644)	30–50	48.0%	NR	NR	NR	–	1.52 (1.15–2.02)
Cohort (23)	USA	3201 (222)	40–89	45.4	222	6.9	1.3	69.0	
Cross-sectional (17)	Greece	1366 (742)	30–50	40.0	NR	NR	NR	NR	1.41 (1.14–1.76)
Cross-sectional (18)	USA	506 (202)	<30	46.0	103 (55)	20.3	NR	36.7	1.79 (1.07–2.99)
Case-control (21)	China	844 (406)	>50	3.9	19 (8)	2.25	4.9	3.5	0.63 (0.22–1.82)
Cross-sectional (24)	Spain	229 (60)	>50	5.0	81 (NR)	35.3	82.5	NR	5.05 (2.30–10.90)
Cohort (8)	France	13 674 (260)	>50	23.0	160 (NR)	1.1	NR	NR	2.90 (1.30–6.32)
Case-control (7)	France	544 (272)	30–50	49.0	20 (17)	3.6	16.2	23.5	9.97 (1.57–63.50)
Cross-sectional (15)	France	35 004 (2520)	>50	26.0	20 236 (131)	57.8	5.2	NR	9.48 (4.14–21.72)
Cross-sectional (25)	Italy	339 (80)	<30	42.0	135 (45)	39.8	5.8	NR	2.06 (1.19–3.56)
Cross-sectional (26)	Saudi Arabia	4679 (474)	<50	13.63	115	10.2	NR	NR	1.90 (1.38–2.60)
Cross-sectional (27)	France	4774 (278)	>50	54.0	1411 (138)	29.5	2.2	NR	2.03 (1.57–2.64)
Case-control (28)	USA	874 (467)	30–50	NR	292 (169)	33.4	47.8	18.3	8.25 (4.53–15.05)

CI = confidence interval; NR = not recorded; OR = odds ratio; USA = United States of America.

## Results

Results from all 17 studies showed that sleepiness and sleep deprivation were major contributors to road traffic accidents. The high frequency of sleepiness reported while driving, with significant odd ratios, makes this a significant risk factor for road traffic accidents. In these studies, >50% (3.5–67.3%) of the participants agreed that they experienced sleep deprivation. Abe et al. reported the highest (58%) frequency of sleepiness during driving in their cross-sectional study (OR 12.90) in Japan, and Nabi et al. reported the lowest (1.1%) in a cohort study in France (OR 2.90) (Figure 2).

Liu et al concluded that a significant decrease in injuries related to road traffic accidents can be attained if fewer people drive when they are sleepy (26). Gottlieb et al. associated sleep apnoea with a 123% greater risk of road traffic accidents than apnoea unrelated to sleep (27). This shows that sleeping for 6 hours daily is connected to a 33% greater risk of accidents than sleeping for 7 or 8 hours per night. Comparatively, Cummings et al. reported such an accident risk to be 39.5% in a case-control study in the United States of America [14]. Lloberes et al found in a cross-sectional study in Spain that 35.3% of drivers fell asleep while driving (28). Pizza et al. reported 1.9 times greater risk of accidents in individuals with poor sleep quality making them to fall asleep while driving (29).

AlShareef et al. in a population-based analysis in Saudi Arabia showed the correlation between sleep and sleepiness during driving and reported that the strongest sleep predictor while driving was being a male driver (30). Most drivers in this study were men (86.5%) as women have only been allowed to drive in Saudi Arabia since June 2018. Sagaspe et al. found that 28% of drivers had at least 1 incident of uncontrollable sleepiness during driving, and about 5% of drivers had an accident or near miss due to sleepiness (31). Stutts et al. reported that almost 8% of the drivers in road traffic accidents admitted consuming alcohol before causing a crash. However, this study highlighted potential bias because the data depended on self-reports and the drivers could have blamed longer work hours and poor sleep habits for their accidents (32).

Limitations of this review include the possibility of selection bias because of the methods used or participants involved. There may have been a residual confounding effect because studies were based on observational

methods. Other possible risk factors for accidents, age, body mass index, medical conditions, alcohol and drug abuse, and sleep duration were not calculated in terms of ORs (adjusted) in all of the studies used for this review.

## Conclusion

Driver fatigue or drowsiness is a road transport safety hazard. The risk of road traffic accidents increases proportionately when drivers experience sleepiness. Among the frequent explanations for sleepiness during driving were sleep disorders such as sleep apnoea and some behavioural factors, most importantly sleep deprivation. The risk factors for sleep deprivation were found to be driving at night, not getting enough sleep, and working or staying awake for long periods. Other factors were young age, male sex, office worker, smoker, shorter sleep duration, poor subjective sleep quality, moderate or severe excessive daytime sleepiness, and

alcoholism. A minimum of 6 hours of sleep every day could significantly decrease the number of road traffic accidents.

To reduce the incidence of accidents related to sleepiness during driving, it is important to conduct safety checks before driving, monitor sleeping patterns, record and track driving hours, and conduct psychological assessments and behavioural training. The results show that road traffic accidents are consistently associated with sleepiness during driving. Therefore, awareness campaigns and strengthening of road safety programmes should be implemented to reduce the increasing number of road traffic accidents related to sleepiness during driving. Further studies will be required for a more in-depth analysis of this subject.

**Funding:** None

**Competing interests:** None declared.

## Évaluation des risques d'accidents de la route liés à la somnolence au volant : une revue systématique

### Résumé

**Contexte :** Les traumatismes dus à un accident représentent la huitième cause de décès dans le monde. La somnolence entraîne une perturbation des fonctions neurologiques et constitue un facteur de risque majeur d'accidents de la circulation.

**Objectifs :** La présente revue systématique a évalué la relation entre la somnolence au volant et les accidents de la circulation.

**Méthodes :** Une analyse systématique a été réalisée en utilisant des bases de données en ligne telles que Wiley Online Library, JSTOR, Medline et PubMed. Des articles complets, en langue anglaise, publiés entre mai 2000 et novembre 2020 ont été extraits. L'accident de la circulation a été fixé comme le résultat d'intérêt et la somnolence au volant comme l'exposition. L'analyse comprenait des études contenant des estimations de risque ajustées (intervalle de confiance à 95 %). Dix études transversales ( $n = 55\,945$ ), cinq études cas-témoins ( $n = 3821$ ) et deux études de cohorte ( $n = 16\,875$ ) ont été incluses.

**Résultats :** Plus de 50 % des participants aux différentes études ont connu une privation de sommeil allant de 3,5 à 67,3 %. Dans leur étude transversale au Japon, Abe et al. ont rapporté la fréquence de somnolence la plus élevée (58 %) et Nabi et al. la plus faible (1,1 %) dans leur étude de cohorte en France.

**Conclusion :** La somnolence et la privation de sommeil étaient liées aux accidents de la circulation, et la privation de sommeil était le principal facteur de somnolence au volant.

## تقييم مخاطر الحوادث المرورية بسبب النعاس أثناء القيادة: استعراض منهجي

شهزاد سليم

### الخلاصة

**الخلفية:** الإصابات الناجمة عن التصادمات هي السبب الرئيسي الثامن للوفاة في العالم. والنعاس، بما يسببه من تعطيل للوظائف العصبية، عامل خطر أساسي للحوادث المرورية.

**الأهداف:** هدفت هذه الدراسة في هذا الاستعراض المنهجي إلى تقييم العلاقة بين النعاس أثناء القيادة والحوادث المرورية.

**طرق البحث:** أجرى الباحث استعراضاً منهجياً باستخدام قواعد البيانات المتاحة على الإنترنت مثل Wiley Online Library وكذلك JSTOR ونظام استرجاع المعلومات البيولوجرافية الطبية والبيولوجية (قاعدة بيانات مدلاين) وقاعدة البيانات الطبية PubMed، حيث استرجع النص الكامل للمقالات المنشورة باللغة الإنجليزية بين مايو/ أيار 2000 ونوفمبر / تشرين الثاني 2020. وحدد الباحث الحوادث المرورية بوصفها النتيجة محل الاهتمام، والنعاس أثناء القيادة بوصفه التعرض. وتضمن الاستعراض دراسات تحتوي على تقديرات مخاطر معدلة (بفترة

ثقة تبلغ 95٪). وشملت الدراسة عشر دراسات مقطعية (العدد = 55,945)، و5 دراسات من دراسات الحالات والشواهد (العدد = 3821)، ودراستين من دراسات الأتراب (العدد = 16,875).

النتائج: كان أكثر من 50٪ من المشاركين في الدراسات المختلفة يعانون من حرمان من النوم، وتراوحت النسبة بين 3.5٪ و67.3٪. وذكر آبي وآخرون أعلى معدل تواتر للنعاس (58٪) في أثناء القيادة في دراستهم المقطعية في اليابان، بينما ذكر نابي وآخرون أدنى معدل (1.1٪) في دراستهم، وهي من دراسات الأتراب في فرنسا.

الاستنتاجات: ارتبط النعاس والحرمان من النوم بحوادث المرور على الطرق، وكان الحرمان من النوم المساهم الرئيسي في النعاس في أثناء القيادة.

## References

1. Brain basics: understanding sleep [website]. National Institute of Neurological Disorders and Stroke (<https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Understanding-Sleep>, accessed 14 June 2022).
2. Sleep deprivation [website]. ResSleep; 2021 <https://www.ressleep.com.au/sleep-deprivation>, accessed 14 June 2022).
3. Sleep deprivation [website]. Better Health Channel; 2021 (<https://www.betterhealth.vic.gov.au/health/conditionsandtreatments/sleep-deprivation>, accessed 14 June 2022).
4. Miyama G, Fukumoto M, Kamegaya R, Hitosugi M. Risk factors for collisions and near-miss incidents caused by drowsy bus drivers. *Int J Environ Res Public Health*. 2020 Jun 18;17(12):4370. <https://doi.org/10.3390/ijerph17124370>. PMID:32570777
5. Road traffic injuries [website]. Geneva: World Health Organization (<https://www.who.int/news-room/fact-sheets/detail/road-traffic-injuries>, accessed 14 June 2022).
6. Preidt, R. Sleepy drivers involved in 100,000 crashes a year [website]. WebMD; 2018 (<https://www.webmd.com/sleep-disorders/news/20181107/sleepy-drivers-involved-in-100000-crashes-a-year>, accessed 14 June 2022).
7. Prevalence of motor vehicle crashes involving drowsy drivers, United States, 1999–2008. *Accid Anal Prev*. 2012 Mar;45:180–6. <https://doi.org/10.1016/j.aap.2011.05.028> PMID:22269499
8. Medic G, Wille M, Hemels ME. Short- and long-term health consequences of sleep disruption. *Nat Sci Sleep*. 2017 May 19;9:151–61. <https://doi.org/10.2147/NSS.S134864> PMID:28579842
9. Connor J, Norton R, Ameratunga S, Robinson E, Civil I, Dunn R, et al. (2002). Driver sleepiness and risk of serious injury to car occupants: population based case control study. *BMJ* 2002 May 11;324(7346):1125. <https://doi.org/10.1136/bmj.324.7346.1125> PMID:12003884
10. Philip P, Chaufton C, Orriols L, Lagarde E, Amoros E, Laumon B, et al. Complaints of poor sleep and risk of traffic accidents: a population-based case-control study. *PLoS One*. 2014 Dec 10;9(12):e114102. <https://doi.org/10.1371/journal.pone.0114102>. PMID:25494198
11. Nabi H, Gueguen A, Chiron M, Lafont S, Zins M, Lagarde E. Awareness of driving while sleepy and road traffic accidents: prospective study in GAZEL cohort. *BMJ*. 2006 Jul 8;333(7558):75. <https://doi.org/10.1136/bmj.333.7558.75> PMID:16798754
12. Ohayon MM. Determining the level of sleepiness in the American population and its correlates. *J Psychiatr Res*. 2012;46(4):422–7. <https://doi.org/10.1016/j.jpsychires.2011.06.008>. PMID:22297274
13. Ellen RL, Marshall SC, Palayew M, Molnar FJ, Wilson KG, Man-Son-Hing M. Systematic review of motor vehicle crash risk in persons with sleep apnea. *J Clin Sleep Med*. 2006 Apr 15;2(2):193–200. PMID:17557495
14. Cummings P, Koepsell TD, Moffat JM, Rivara FP. Drowsiness, counter-measures to drowsiness, and the risk of a motor vehicle crash. *Inj Prev*. 2001 Sep;7(3):194–9. <https://doi.org/10.1136/ip.7.3.194>. PMID:11565983
15. Abe T, Komada Y, Nishida Y, Hayashida K, Inoue Y. Short sleep duration and long spells of driving are associated with the occurrence of Japanese drivers rear-end collisions and single-car accidents. *J Sleep Res*. 2010 Jun;19(2):310–316. <https://doi.org/10.1111/j.1365-2869.2009.00806.x>. PMID:20337905
16. Carter, N., Ulfberg, J., Nyström, B., & Edling, C. (2003). Sleep debt, sleepiness and accidents among males in the general population and male professional drivers. *Accident; analysis and prevention*, 35(4), 613–617. [https://doi.org/10.1016/s0001-4575\(02\)00033-7](https://doi.org/10.1016/s0001-4575(02)00033-7). PMID:12729824
17. Drake CL, Roehrs T, Richardson G, Walsh JK, Roth T. Shift work sleep disorder: prevalence and consequences beyond that of symptomatic day workers. *Sleep*. 2004 Dec 15;27(8):1453–1462. <https://doi.org/10.1093/sleep/27.8.1453>. PMID:15683134
18. Philip P, Sagaspe P, Lagarde E, Leger D, Ohayon MM, Bioulac B, et al. Sleep disorders and accidental risk in a large group of regular registered highway drivers. *Sleep Med*. 2010 Dec;11(10):973–979. <https://doi.org/10.1016/j.sleep.2010.07.010>. PMID:20961809
19. BaHammam AS, Alkhunizan MA, Lesloum RH, Alshantqi AM, Aldakhil AM, Pandi-Perumal SR et al. Prevalence of-related accidents among drivers in Saudi Arabia. *Ann Thorac Med*. 2014 Oct;9(4):236–41. <https://doi.org/10.4103/1817-1737.140138> PMID:25276244 PMCID: PMC4166072
20. Czeisler CA, Wickwire EM, Barger LK, Dement WC, Gamble K, Hartenbaum N, et al. Sleep-deprived motor vehicle operators are unfit to drive: a multidisciplinary expert consensus statement on drowsy driving. *Sleep Health*, 2016 Jun;2(2):94–9. <https://doi.org/10.1016/j.sleh.2016.04.003> PMID:28923267



21. Gander PH, Marshall NS, Harris RB, Reid P. Sleep, sleepiness and motor vehicle accidents: a national survey. *Aust N Z J Public Health*. 2005 Feb;29(1):16–21. <https://doi.org/10.1111/j.1467-842x.2005.tb00742.x> PMID:15782866.
22. Gnardellis C, Tzamalouka G, Papadakaki M, Chliaoutakis JE. An investigation of the effect of sleepiness, drowsy driving, and lifestyle on vehicle crashes. *Transp Res Traff Psychol Behav*. 2008 Jul;11(4):270–81. (<https://www.sciencedirect.com/science/article/pii/S1369847808000107>)
23. Hutchens L, Senserrick TM, Jamieson PE, Romer D, Winston FK. Teen driver crash risk and associations with smoking and drowsy driving. *Accid Anal Prev*. 2008 May;40(3):869–76. <https://doi.org/10.1016/j.aap.2007.10.001>. PMID:18460353.
24. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *BMJ*. 2009;339:b2700. <https://doi.org/10.1136/bmj.b2700>
25. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandom studies in meta-analyses [website]. Ottawa Hospital Research Institute ([http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp), accessed 15 June 2022).
26. Liu GF, Han S, Liang DH, Wang FZ, Shi XZ, Yu J, et al. Driver sleepiness and risk of car crashes in Shenyang, a Chinese north-eastern city: population-based case-control study. *Biomed Environ Sci*. 2003 Sep;16(3):219–26. PMID:14631827.
27. Gottlieb DJ, Ellenbogen JM, Bianchi MT, Czeisler CA. Sleep deficiency and motor vehicle crash risk in the general population: a prospective cohort study. *BMC Med*. 2018 Mar 20;16(1):44. <https://doi.org/10.1186/s12916-018-1025-7> PMID:29554902
28. Lloberes P, Levy G, Descals C, Sampol G, Roca A, Sagales T, et al. Self-reported sleepiness while driving as a risk factor for traffic accidents in patients with obstructive sleep apnoea syndrome and in non-apnoeic snorers. *Respir Med*. 2000 Oct;94(10):971–6. <https://doi.org/10.1053/rmed.2000.0869> PMID:11059950
29. Pizza F, Contardi S, Antognini AB, Zagoraiou M, Borrotti M, Mostacci B, et al. Sleep quality and motor vehicle crashes in adolescents. *J Clin Sleep Med*. 2010 Feb 15;6(1):41–5. PMID:20191936
30. AlShareef SM. Excessive daytime sleepiness and associations with sleep-related motor vehicle accidents: results from a nationwide survey. *Sleep Breath*. 2021 Sep;25(3):1671–6. <https://doi.org/10.1007/s11325-020-02260-5>. PMID:33242183.
31. Sagaspe P, Taillard J, Bayon V, Lagarde E, Moore N, Boussuge J, et al. Sleepiness, near-misses and driving accidents among a representative population of French drivers. *J Sleep Res*. 2010 Dec;19(4):578–84. <https://doi.org/10.1111/j.1365-2869.2009.00818.x> PMID:20408921
32. Stutts JC, Wilkins JW, Scott Osberg J, Vaughn BV. Driver risk factors for sleep-related crashes. *Accid Anal Prev*. 2003 May;35(3):321–31. [https://doi.org/10.1016/s0001-4575\(02\)00007-6](https://doi.org/10.1016/s0001-4575(02)00007-6) PMID:12643949.