

Changes in lifestyle behaviour and dietary patterns among Iranian medical students during COVID-19 lockdown

Afsaneh Soltani,^{1,2,3} Noosha Samieefar^{1,2,3} and Meisam Akhlaghdoust^{3,4,5}

¹Student Research Committee, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ²USERN Office, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ³Functional Neurosurgery Research Center, Shohada Tajrish Comprehensive Neurosurgical Center of Excellence, Shahid Beheshti University of Medical Sciences, Tehran, Iran (Correspondence to: M. Akhlaghdoust: meisam_akhlagh@yahoo.com). ⁴USERN Office, Functional Neurosurgery Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ⁵Virtual Rete of Gifted and Talented Medical Students, Smart University of Medical Sciences, Iran.

Abstract

Background: The COVID-19 pandemic has affected many aspects of human life, including physical and mental health. The public health measures to prevent COVID-19 transmission led to social isolation, physical inactivity, anxiety, depression, sleep disorders, obesity, and consumption of unhealthy food.

Aims: To evaluate changes in lifestyle behaviour during the fifth wave of COVID-19 in Iranian medical students.

Methods: This was a cross-sectional descriptive study. In September 2021, an online questionnaire containing 32 questions was administered to 1143 medical students in Teheran universities. Some 286 completed copies of the questionnaire were excluded because of incomplete answers. Descriptive statistics of central tendency measures and dispersion were used and independent t-test and a Chi-square test between two variables were conducted. Statistical analysis was performed using SPSS version 24.

Results: A total of 462 (53.9%) students had COVID-19 during the fifth wave of the disease, 657 (76.7%) were deprived of social and sports activities, and 278 (32.4%) had difficulty completing their academic studies. Students who took vitamin supplements experienced lower levels of stress and 711 (83.0%) reported bodyweight change. There were significant relationships between bodyweight change, stress level and hospitalization due to COVID-19. There was a significant relationship between smoking, bodyweight change and hospitalization rate. A total of 426 (50.3%) students did not have a regular sleep-wake pattern, 701 (81.8%) spent more time using social media, 197 (23.0%) consumed more fast food, and 674 (78.6%) experienced stress or anxiety.

Conclusion: There was an increased tendency towards unhealthy lifestyle among medical students in Tehran during COVID-19 lockdown. These findings can be useful in assessing lifestyle behaviours among students and in planning for improvements.

Keywords: COVID-19, medical students, lifestyle, lockdown, Islamic Republic of Iran

Citation: Soltani A; Samieefar N; Akhlaghdoust M. Changes in lifestyle behaviour and dietary patterns among Iranian medical students during COVID-19 lockdown. *East Mediterr Health J.* 2022;28(12):896–903. <https://doi.org/10.26719/emhj.22.094>

Received: 01/06/21; accepted: 11/08/22

Copyright © Authors 2022; Licensee: World Health Organization. EMHJ is an open access journal. This paper is available under the Creative Commons Attribution Non-Commercial ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Introduction

Coronaviruses are a large family that can cause respiratory infections, ranging from the common cold to more serious illnesses, such as COVID-19, which is caused by the novel SARS-CoV-2 (1). The COVID-19 pandemic has affected many aspects of human life, including physical and mental health. To evaluate lifestyle, the World Health Organization recommends assessment of alcohol consumption, smoking, physical exercise, nutrition, and stress (2). Lifestyle is a complex concept that reflects people's awareness of their lives and personalities (3). Due to the COVID-19 pandemic, there has been a sudden change in people's lifestyles and habits. Quarantine and social distancing have led to students participating in their classes online, and staying at home has reduced physical activity. Listening to news about the increasing number of COVID-19 patients, as well as lack of physical activity, have caused stress, depression, and anxiety.

Decreased physical activity leads to obesity, which increases the risk of cardiovascular disease. Obesity can also be caused by being alone at home and under stress, which increases the desire to eat and consume sugar (4). A study on physicians and hospital nurses in Wuhan, China during the outbreak of COVID-19 showed that medical care workers had a high degree of depressive symptoms and anxiety (5).

Many countries implemented social distancing and stay-at-home policies to prevent transmission of COVID-19. These policies reduced morbidity and mortality, but they also had an adverse effect on mental health, emotional state, and lifestyle behaviour (6,7). According to Kumari et al., social isolation and staying at home increased overeating and inactivity, and disrupted sleep patterns (8). Another study of 584 young people, 2 weeks after the outbreak of COVID-19 in China, found that 40.4% of the participants were prone to mental

health problems and 14.4% had signs of developing post-traumatic stress disorder (9).

COVID-19 and social isolation have disrupted the education of medical students, according to a study in the United States of America, gradually affecting students' physical, mental, and emotional health, and increasing anxiety. Medical students worldwide are a vulnerable population, and higher rates of depression and suicidal ideation were reported among them during the COVID-19 pandemic, and they were less likely to seek support, than students in previous studies (10).

This study was to investigate the lifestyle patterns and their influencing factors among medical students in Tehran universities during the fifth wave of the COVID-19 pandemic in the Islamic Republic of Iran.

Methods

This was a web-based, cross-sectional descriptive study conducted during the fifth wave of the COVID-19 pandemic (delta variant) in the Islamic Republic of Iran. This wave peaked in August 2021 and was caused by the delta variant of SARS-CoV-2. Medical students at Shahid Beheshti University, Tehran University of Medical Sciences, Iran University of Medical Sciences, and Islamic Azad University, Tehran Medical Branch were surveyed online in September 2021 using the snowball sampling method. Data were collected using Google Forms through a self-reported questionnaire that was designed using a previously validated questionnaire of lifestyle patterns (11). The questionnaire was completed by 1143 students; 857 fully answered the questions and 286 incomplete questionnaires were excluded from the study. We also excluded students aged < 18 and > 30 years and first-semester students (because of confounding factors such as stress, lifestyle changes, and relocation to start university), as well as students with history of disability and psychiatric illness. Students' names and phone numbers were not collected to protect confidentiality.

The questionnaire consisted of 32 questions divided into 2 parts: the first included demographic information, and the second, information about the health and lifestyle of students during the COVID-19 pandemic. There were 5 questions about history of COVID-19, hospitalization, remdesivir injection, family members with COVID-19, and death of relatives from COVID-19.

Statistical analysis was performed using SPSS version 24. The mean and standard deviation were calculated as descriptive statistics. Independent *t* test and bivariate χ^2 test were used to assess the correlation between lifestyle aspects and COVID-19 infection rate, hospitalization, and stress level. $P < 0.05$ indicated statistical significance.

Results

Sociodemographic characteristics of survey respondents

There were 161 male (18.8%) and 696 female (81.2%) participants, 792 were single (92.4%) and 65 (7.6%) were married. The mean (standard deviation) age of students was 23.4 (1.8) years. The average age of female students was 21.8 (3.6) years, and the mean age of male students was 24.6 (4.4) years. Seven hundred and sixty-one (88.8%) students were unemployed, 75 (8.8%) were employed, and 21 did not answer this question. There were 647 (75.9%) undergraduates, 54 (3.6%) masters students, and 156 (18.2%) doctoral students. Most students ($n = 782$, 91.2%) had health insurance, and 104 (12.1%) had not been vaccinated as of the time of the survey.

Lifestyle changes during COVID-19 lockdown

Physical activity

Three hundred and eight (35.9%) students stated that they undertook physical activity at least once or twice, or ≥ 3 times weekly (Table 1). Physical distancing restricted physical activity, learning, and daily activities in 657 (76.7%) students (Table 2). Students who exercised less during the week had more difficulty in studying because of stress, anxiety, or depression ($P = 0.006$) (Table 3).

Medical history and drug treatment

Six hundred and sixty-eight (77.9%) students had no history of underlying disease (Table 4). Eighty-three (9.7%) students reported regular drug treatment, with levothyroxine being the most common ($n = 22$; 2.67%) (results not shown). Students who took vitamin supplements experienced significantly lower levels of stress ($P = 0.003$) (Table 5).

Rate of COVID-19 infection

Four hundred and sixty-two (53.9%) students acquired COVID-19 during the fifth wave of the disease, 182 (21.2%) required remdesivir injection, and 32 (3.7%) were hospitalized with severe COVID-19 (Table 4).

Body weight and obesity

Most students ($n = 625$; 72.9%) did not have a sudden change in health status during the COVID-19 pandemic, but 91 (10.6%) reported such a change (Table 2). A large number of students (83.0%, $n = 711$) had a change in body weight and 78.6% ($n = 674$) reported becoming overweight. Change in body weight in the last 6 months resulted in significant increases in stress levels ($P = 0.03$) and hospitalization because of COVID-19 ($P = 0.007$) (Table 5).

Smoking

There were 809 (94.4%) nonsmokers and only 47 (5.6%) smokers (Table 1). Most students ($n = 831$, 97.0%) either did not smoke at all or did not smoke during the COVID-19 pandemic (Table 2). Students who smoked had more academic problems ($P = 0.002$), and more difficulty in studying because of depression, anxiety, or stress

Table 1 Personal health habits of medical students in Tehran during the COVID-19 pandemic in September 2021

Weekly physical activity	Never	112 (13.1%)
	1 or 2 times	308 (35.9%)
	≥ 3 times	308 (35.9%)
	Every day	129 (15.1%)
Daily tooth brushing/mouth washing	Never	5 (0.6%)
	Occasionally	41 (4.8%)
	1 or 2 times	693 (80.7%)
	≥ 3 times	118 (13.9%)
Regular Sleep-wake pattern	No	431 (50.3%)
	Yes	426 (49.7%)
Average daily sleep duration	< 6 hours	117 (13.7%)
	6–8 hours	460 (53.7%)
	8–10 hours	262 (30.6%)
	≥ 10 hours	18 (2.1%)
Smoking	Yes	47 (6.5%)
	No	809 (94.4%)
	No answer	
Average daily time on computer, tablet or mobile	Never	26 (3.0%)
	< 2 hours	249 (29.1%)
	2–4 hours	297 (34.7%)
	> 4 hours	258 (30.1%)
	No answer	27 (3.2%)

($P = 0.002$) (Table 3). Smokers also had a higher rate of hospitalization because of COVID-19 ($P < 0.001$).

Sleeping

Approximately half of the students ($n = 431$, 50.3%) did not have a regular sleep-wake pattern (Table 1). Four hundred and sixty (53.7%) students slept between 6 and 8 hours daily, 262 (30.6%) between 8 and 10 hours, 117 (13.7%) < 6 hours, and 18 (2.1%) > 10 hours. Five hundred and eighty-four (68.1%) students experienced reduced sleep time (Table 2). Students with irregular sleep patterns and those who slept > 10 or < 6 hours were more likely to have COVID-19 and to be hospitalized because of it (Table 3).

Social media (screen time)

Computer, tablet, and mobile phone use was > 4 hours daily (258 students, 30.1%), 2–4 hours daily (297 students, 34.7%), and < 2 hours daily (249 students, 29.1%) (Table 1). A large number of students ($n = 701$, 81.8%) reported that they spent more time using social media during the COVID-19 pandemic (Table 2), and these students were more likely to develop COVID-19 ($P = 0.004$) (Table 3).

Eating habits and nutritional changes

Increased consumption of fast food during the COVID-19 pandemic was reported by 584 (68.1%) students (Table 2). There was a significant relationship between consumption of fast food and incidence of COVID-19 ($P = 0.001$) and related hospitalization ($P < 0.001$) (Table 5). Students who consumed more fruits and vegetables were less likely to have COVID-19 ($P = 0.009$).

Mental health and hygiene

Six hundred and seventy-four (78.6%) students experienced excessive stress or anxiety during the COVID-19 pandemic (Table 2). Most students ($n = 776$, 90.5%) reported that they had an obsession with washing

or hygiene during the pandemic. Weekly consumption of fast food was significantly associated with stress level ($P = 0.002$). Students who were less physically active ($P = 0.024$) and used more social media ($P = 0.02$) were more obsessed with hygiene and handwashing (Table 3). Six hundred and ninety-three (80.7%) students used toothbrushes and mouthwashes at least once or twice daily, 118 (13.9%) used them ≥ 3 times, and 42 (4.8%) used them occasionally (Table 1). Five (6.0%) students stated that they did not use a toothbrush or mouthwash at all.

Discussion

This study investigated the changes in lifestyle behaviour of medical students in Tehran during the fifth wave of COVID-19. At the time of the study, partial lockdown and some physical distancing restrictions were implemented, and consequently many social activities were cancelled. Our study demonstrated that several factors that affected students' lifestyles, including physical activity, sleep patterns, smoking, nutrition, depression, and anxiety, changed significantly during the COVID-19 pandemic, indicating that students experienced an unhealthy and sedentary lifestyle.

A systematic review by Ammar et al. (10), which investigated the lifestyles of 1047 adults found that, during the COVID-19 pandemic, physical activity decreased, sleep quality deteriorated, unhealthy dietary intake increased, and average time spent sitting at home increased from 5 to 8 hours. We found that ~78% of the study participants reported that their physical activity was limited during the COVID-19 pandemic. This is consistent with a study in Poland that reported reduced physical activity among adults during the pandemic (12), and a study in Canada that reported a significant reduction in physical activity among children and adolescents (13). In Northern Italy,

22% of participants increased their physical activity, 14% of adults maintained a constant level of physical activity, but 50% decreased their physical activity during the COVID-19 pandemic (14).

According to Dragun et al. (15), 32% of Croatian medical students reported weight loss and 19% reported weight gain during lockdown, while physical activity remained constant. These findings are inconsistent with our study because we found that 78.6% of students reported being overweight and 10.5% lost weight during the fifth COVID-19 wave in the Islamic Republic of Iran.

In our study, people with irregular sleep patterns and those who slept > 10 or < 6 hours per night were more likely to develop COVID-19 and to be hospitalized. They also reported sudden changes in their health during the COVID-19 pandemic. In the Republic of Korea, as people spent more time at home, their total sleep time increased, but sleep satisfaction decreased during the COVID-19 pandemic (16). This suggests that participants may have experienced poor and erratic sleep patterns.

Rawat et al. showed that high weight and stress led to a lack of sleep (17), which was similar to our study. Alena

Table 2 Effect of physical distancing restrictions on health among medical students in Tehran in September 2021

Did you have a sudden change in health status during the COVID-19 pandemic?	Yes	91 (10.6%)
	No	625 (72.9%)
	No answer	141 (16.5%)
Did your body weight change during the COVID-19 pandemic?	No	146 (17.0%)
	Yes	711 (83.0%)
How has your body weight changed during the COVID-19 pandemic?	Decreased	90 (10.5%)
	Increased	674 (78.6%)
	No change	93 (10.8%)
Did the lockdown and pandemic limit you from exercising, daily activities and learning?	No	32 (3.7%)
	Yes	657 (76.7%)
	No answer	168 (19.6%)
Did you have difficulty completing your academic duties because of COVID-19?	Yes	278 (32.4%)
	No	483 (56.4%)
	No answer	96 (11.2%)
Have you smoked or increased smoking during the COVID-19 pandemic?	Yes	26 (3.0%)
	No	831 (97%)
How have your sleep patterns changed during the COVID-19 pandemic?	Decreased	584 (68.1%)
	Increased	111 (12.9%)
	No change	162 (18.9%)
How has your media usage changed during the COVID-19 pandemic?	Decreased	65 (7.6%)
	Increased	701 (81.8%)
	No change	91 (10.6%)
How has your fast-food consumption changed during the COVID-19 pandemic?	Decreased	47 (5.5%)
	Increased	197 (23.0%)
	No change	584 (68.1%)
Have you experienced excessive stress or anxiety because of the COVID-19 pandemic?	Yes	674 (78.6%)
	No	183 (21.4%)
Have you been too obsessed with hygiene and washing during the COVID-19 pandemic?	Yes	81 (9.5%)
	No	776 (90.5%)

Table 3 Correlation between changes in lifestyle among medical students in Tehran during the COVID-19 pandemic in September 2021

	Weekly physical activity		Regular sleep pattern		Daily hours of sleep		Smoking		Daily hours of media usage	
	r ^a	P	r	P	r	P	r	P	r	P
Sudden changes in health during the COVID-19 pandemic	0.75	0.861	4.81	0.028	0.09	0.004	0.07	0.034	0.05	0.100
Weight changes during the COVID-19 pandemic	7.68	0.053	2.29	0.130	0.09	0.004	0.02	0.385	-0.08	0.804
Difficulty in studying because of stress, anxiety or depression	44.88	0.006	11.67	0.166	-0.04	0.111	-0.09	0.002	0.02	0.442
Hospitalization because of COVID-19	3.29	0.308	0.42	0.517	0.08	0.017	0.19	<0.001	0.03	0.250
Infected with COVID-19	3.59	0.308	12.82	<0.001	0.12	<0.001	0.05	0.084	0.09	0.004
Hygiene obsession	31.62	0.024	0.76	0.135	-0.10	0.849	-0.07	0.195	0.06	0.020

^ar refers to χ^2 statistics. Bold values are statistically significant.

Table 4 Medical history of medical students in Tehran in September 2021

Do you have a history of underlying disease?	Yes	189 (22.1%)
	No	668 (77.9%)
Did you acquire COVID-19 during this wave?	Yes	462 (53.9%)
	No	316 (36.9%)
	No answer	79 (9.2%)
Did you need to inject remdesivir?	Yes	182 (21.2%)
	No	632 (73.7%)
	No answer	43 (5.0%)
Were you hospitalized because of COVID-19?	Yes	32 (3.7%)
	No	825 (96.3%)
Did any of your first-degree family members acquire COVID-19?	Yes	58 (6.8%)
	No	662 (77.2%)
	No answer	137 (16.0%)
Did you lose your first-degree relatives because of COVID-19?	Yes	21 (2.4%)
	No	836 (97.6%)

et al. found that reduced physical activity and stress decreased sleep quality during the COVID-19 pandemic (18). In the study of Dragun et al. (15), the average sleep duration increased by 1.5 hours. However, among nursing students, the quality of sleep was lower despite their longer sleep time (19). This result is inconsistent with our study because 68% of participants experienced reduced sleep time. Ismail et al. found that stress and anxiety impaired the quality of sleep, and reduced energy levels during the day: 10.8% of participants had poor sleep quality and there was a 26.2% increase among people who felt tired during the pandemic (20).

Our study showed a significant relationship between healthy food intake and the rate of COVID-19 infection and hospitalization and stress levels. Our findings were consistent with the study of Rawata et al. (17), which revealed that when people experienced stress and anxiety, their cortisol and blood insulin levels increased; both of which led to increased appetite and likelihood of choosing unhealthy foods. Similarly, 68.1% of students in our study experienced more fast food consumption during the pandemic. However, 2 other studies (21, 22) indicated that fast food intake was reduced during the COVID-19 pandemic.

We found that 78.6% of the participants experienced anxiety and distress. Similarly, Liu et al. reported that 50.7% of mental health professionals experienced depressive symptoms and 44.7% reported anxiety (23). Giuntella et al. (24) found that depression increased as a result of physical distancing and reduced socialization during the COVID-19 pandemic. Socializing was < 30 minutes a day, and the time spent on mobile phones and laptops doubled. In our study, a significant number of students (81.7%) admitted that their screen time increased during the COVID-19 pandemic. Students reported that they had more stress and anxiety when social interactions were reduced. This was similar to the study of Kuang et al. (18), which found that reducing social interactions increased stress levels among the general population.

In our study, participants who smoked had more academic difficulties and mental health problems during the COVID-19 pandemic, as well as a higher rate of hospitalization for COVID-19. According to Di Renzo et al. (4), some smokers in Italy reduced the number of cigarettes smoked daily because of fear of respiratory distress and death from COVID-19, and some had quit smoking altogether during the pandemic.

In a study of northern Italians during the COVID-19 pandemic, 23% started taking supplements to boost their immune systems (24). In our study, 21% of participants took vitamin C (11.4%) and vitamin D (9.9%) supplements.

According to a Dutch study of 1004 respondents aged 18–88 years (20), about one fifth adjusted their lifestyle to healthier eating habits, exercise, and relaxation. However, 10% of the total population surveyed stated that they had acquired an unhealthy lifestyle because of the COVID-19 pandemic.

Some of the strengths of our study were the large sample size and online nature of the questionnaire because our target population was students and they all had access to the internet. However, the online nature of the questionnaire and the availability of self-reporting tools may have led to inaccurate reporting. One of the limitations of this study was that the number of female respondents was about 4 times that of men. In a study by Smith et al. (25), the number of female participants in online surveys was higher, and should be considered when reviewing the results. Another limitation was that

Table 5 Correlation between nutritional status and COVID-19 infection, COVID-19 hospitalization and stress in medical students in Tehran in September 2021

	COVID-19 infection rate		COVID-19 hospitalization		Stress level	
	r ^a	P	r	P	r	P
Daily consumption of fruits and vegetables	11.54	0.009	8.48	0.205	5.98	0.112
Fluid intake	0.10	0.992	22.18	0.001	0.12	0.988
Weekly consumption of fast food	16.66	0.001	31.19	<0.001	16.81	0.002
Daily consumption of dairy products	3.03	0.386	6.53	0.366	5.78	0.122
Taking vitamin supplements	4.07	0.254	0.72	0.695	15.81	0.003
Body weight change in the last 6 months	2.14	0.343	9.93	0.007	6.99	0.030

^ar refers to χ^2 statistics. Bold values are statistically significant.

we only examined medical students in Tehran, and did not compare them with students in other fields in other cities. We recommend that this study be performed with a wider range of variables and a larger population.

Conclusions

Our study showed a tendency towards unhealthy lifestyle among medical students in Tehran during the COVID-19 lockdown. These changes affected sleep, body weight, obesity, nutrition, use of social media, mental health, and physical activity. In the current situation in which the pandemic has stabilized to some extent as a result of vaccination, it is important to gather more research evidence in this area, to identify students prone

to psychological, behavioural, and nutritional disorders that may endanger their physical and mental health. The evidence could help in planning and determining priorities. It is also necessary to be able to maintain and promote medical students' mental health as the core of the country's medical staff, with appropriate psychological and behavioural strategies and techniques. The main purpose of this study was to assess the lifestyle changes in medical students in Tehran, which could be helpful for health policymakers and physicians. As the COVID-19 pandemic continues, our results should be validated in future larger population studies.

Funding: None.

Competing interests: None declared.

Changements dans les comportements liés au mode de vie et les habitudes alimentaires des étudiants en médecine iraniens pendant le confinement dû à la COVID-19

Résumé

Contexte : La pandémie de COVID-19 a touché de nombreux aspects de la vie humaine, y compris la santé physique et mentale. Les mesures de santé publique visant à prévenir la transmission de la COVID-19 ont conduit à l'isolement social, à la sédentarité, à l'anxiété, à la dépression, aux troubles du sommeil, à l'obésité et à la consommation d'aliments malsains.

Objectifs : Évaluer les changements dans les comportements liés au mode de vie des étudiants en médecine iraniens pendant la cinquième vague de COVID-19.

Méthodes : Il s'agissait d'une étude transversale descriptive. En septembre 2021, un questionnaire en ligne contenant 32 questions a été remis à 1143 étudiants en médecine dans les universités de Téhéran. Quelque 286 copies remplies du questionnaire ont été exclues en raison de l'incomplétude des réponses. Des statistiques descriptives des mesures de tendance centrale et de dispersion ont été utilisées et un test t indépendant et un test chi carré entre deux variables ont été effectués. Une analyse statistique a été réalisée à l'aide du logiciel SPSS version 24.

Résultats : Au total, 462 (53,9 %) étudiants avaient contracté la COVID-19 pendant la cinquième vague de la maladie, 657 (76,7 %) étaient privés d'activités sociales et sportives et 278 (32,4 %) avaient des difficultés à terminer leurs études universitaires. Les étudiants qui avaient pris des suppléments vitaminiques ont connu des niveaux de stress plus faibles et 711 (83,0 %) ont signalé un changement de poids corporel. Il existait des relations significatives entre l'évolution du poids corporel, le niveau de stress et l'hospitalisation due à la COVID-19. Il y avait une relation significative entre le tabagisme, l'évolution du poids corporel et le taux d'hospitalisation. Au total, 426 étudiants (50,3 %) n'avaient pas de sommeil ou d'éveil régulier, 701 (81,8 %) passaient plus de temps à utiliser les médias sociaux, 197 (23,0 %) consommaient davantage de fast-food et 674 (78,6 %) souffraient de stress ou d'anxiété.

Conclusion : Une tendance accrue à un mode de vie malsain a été constatée chez les étudiants en médecine à Téhéran pendant le confinement dû à la COVID-19. Ces résultats peuvent être utiles pour évaluer les comportements liés au mode de vie des étudiants et pour planifier des améliorations.

التغيرات التي طرأت على سلوكيات أنماط الحياة والأنماط الغذائية بين طلاب الكليات الطبية الإيرانية في أثناء فترة حظر كوفيد-19

عفسانة سلطاني، نوشة ساميعفار، ميسام أخلاجدوست

الخلاصة

الخلفية: أثرت جائحة كوفيد-19 على العديد من جوانب حياة الإنسان، ومنها الصحة البدنية والعقلية. ولقد أدت تدابير الصحة العامة الرامية إلى منع انتقال فيروس كوفيد-19 إلى العزلة الاجتماعية والخمول البدني والقلق والاكتئاب واضطرابات النوم والسمنة واستهلاك أغذية غير صحية.

الأهداف: هدفت هذه الدراسة إلى تقييم التغيرات التي طرأت على سلوكيات أنماط الحياة خلال الموجة الخامسة من مرض كوفيد-19 في جمهورية إيران الإسلامية.

طرق البحث: هذه دراسة وصفية مقطعية. وفي سبتمبر/ أيلول 2021، قُدم استبيان إلكتروني يحتوي على 32 سؤالاً إلى 1143 طالباً في كليات الطب بجامعة طهران. واستُبعد قرابة 286 نسخة من استبيانات المشاركين بسبب عدم اكتمال الإجابات. واستُخدمت إحصاءات وصفية لمقاييس الاتجاه المركزي والتشتت، وجرت الاستعانة باختبار "t-test" المستقل، واختبار مربع كاي بين متغيرين. واستُخدم الإصدار 24 من برنامج SPSS لإجراء التحليل الإحصائي.

النتائج: إجمالاً، أُصيب 462 طالباً (53.9%) بمرض كوفيد-19 خلال الموجة الخامسة من المرض، وحُرم 657 طالباً (76.7%) من الأنشطة الاجتماعية والرياضية، وجابه 278 طالباً (32.4%) صعوبة في إكمال دراساتهم الأكاديمية. وأما الطلاب الذين تناولوا مكملات الفيتامينات، فعانوا مستويات أقل من التوتر، وأفاد 711 طالباً (83.0%) بحدوث تغيير في وزن الجسم. ولقد خلص إلى وجود علاقات مهمة بين تغيير وزن الجسم ومستوى الإجهاد والإدخال إلى المستشفى بسبب كوفيد-19. كذلك تبين وجود علاقة مهمة بين التدخين وتغيير وزن الجسم ومعدل الإدخال إلى المستشفى. وافتقر 426 طالباً (50.3%) إلى نمط نوم - استيقاظ منظم، وقضى 701 طالب (81.8%) وقتاً أطول في استخدام وسائل التواصل الاجتماعي، وتناول 197 طالباً (23.0%) المزيد من الوجبات السريعة، وتعرّض 674 طالباً (78.6%) للتوتر أو القلق.

الاستنتاجات: تبين وجود اتجاه متزايد نحو نمط حياة غير صحي بين طلاب الطب في طهران في أثناء فترة الإغلاق لكوفيد-19. وهذه النتائج يمكن أن تكون مفيدة في تقييم سلوكيات نمط الحياة بين الطلاب من جهة، وفي التخطيط لتحسينات في هذا الشأن من جهة أخرى.

References

1. Rezaei N, Ashkevarian S, Fathi MK, Hanaei S, Kolahchi Z, Ladi Seyedian S-S, et al. Introduction on coronavirus disease (COVID-19) pandemic: the global challenge. *Adv Med Exp Biol*. 2021;1318:1–22. https://doi.org/10.1007/978-3-030-63761-3_1 PMID:33973169
2. Angelika S, Anja F, Merle C. Sleep problems in university students – an intervention. *Neuropsychiatr Dis Treat*. 2017 Jul 26;13:1989–2001. <https://doi.org/10.2147/NDT.S142067> PMID:28794633
3. Hershner SD, Chervin RD. Causes and consequences of sleepiness among college students. *Nat Sci Sleep*. 2014 Jun 23;6:73. <https://doi.org/10.2147/NSS.S62907> PMID:25018659
4. Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med*. 2020 Jun 8;18(1):1–15. <https://doi.org/10.1186/s12967-020-02399-5> PMID:32513197
5. Casigliani V, De Nard F, De Vita E, Arzilli G, Grosso FM, Quattrone F, et al. Too much information, too little evidence: is waste in research fuelling the covid-19 infodemic. *BMJ*. 2020;370:m2672. <https://doi.org/10.1136/bmj.m2672>
6. Odone A, Lugo A, Amerio A, Borroni E, Bosetti C, Carreras G, et al. COVID-19 lockdown impact on lifestyle habits of Italian adults. *Acta Biomed*. 2020 Jul 20;91(9-S):87–9. <https://doi.org/10.23750/abm.v91i9-S.10122> PMID:32701921
7. Ammar A, Trabelsi K, Brach M, Chtourou H, Boukhris O, Masmoudi L, et al. Effects of home confinement on mental health and lifestyle behaviours during the COVID-19 outbreak: insights from the ECLB-COVID19 multicentre study. *Biol Sport*. 2021 Mar;38(1):9–21. <https://doi.org/10.5114/biolsport.2020.96857> PMID:33795912
8. Kumari A, Ranjan P, Vikram NK, Kaur D, Sahu A, Dwivedi SN, et al. A short questionnaire to assess changes in lifestyle-related behaviour during COVID 19 pandemic. *Diabetes Metab Syndr*. 2020 Nov–Dec;14(6):1697–701. <https://doi.org/10.1016/j.dsx.2020.08.020> PMID:32911201
9. Liang L, Ren H, Cao R, Hu Y, Qin Z, Li C, et al. The effect of COVID-19 on youth mental health. *Psychiatric quarterly*. 2020 Sep;91(3):841–52. <https://doi.org/10.1007/s11126-020-09744-3> PMID:32319041
10. Halperin SJ, Henderson MN, Prenner S, Grauer JN. Prevalence of anxiety and depression among medical students during the Covid-19 pandemic: a cross-sectional study. *J Med Educ Curric Dev*. 2021 Feb 15;8:1–7. <https://doi.org/10.1177/2382120521991150> PMID:33644399
11. Mansouri M, Sharifi F, Varmaghani M, Yaghubi H, Tabrizi YM, Raznahan M, et al. Iranian university students lifestyle and health status survey: study profile. *J Diab Metab Disord*. 2017 Dec 20;16(1):1–10. <https://doi.org/10.1186/s40200-017-0329-z> PMID:29270393
12. Górnicka M, Drywień ME, Zielinska MA, Hamułka, J. Dietary and lifestyle changes during COVID-19 and the subsequent lockdowns among Polish adults: a cross-sectional online survey PLifeCOVID-19 study. *Nutrients*. 2020 Aug 3;12(8):2324. <https://doi.org/10.3390/nu12082324> PMID:32756458
13. Liu X, Liu J, Zhong X. Psychological state of college students during COVID-19 epidemic. *Psychol Schools*. 2022;1–11. <https://doi.org/10.1002/pits.2269>
14. Canello R, Soranna D, Zambra G, Zambon A, Invitti C. Determinants of the lifestyle changes during COVID-19 pandemic in the residents of Northern Italy. *Int J Environ Res Public Health*. 2020 Aug 28;17(17):6287. <https://doi.org/10.3390/ijerph17176287> PMID:32872336
15. Dragun R, Veček NN, Marendić M, Pribisalić A, Đivić G, Cena H, et al. Have lifestyle habits and psychological well-being changed among adolescents and medical students due to COVID-19 lockdown in Croatia? *Nutrients*. 2020 Dec 30;13(1):97. <https://doi.org/10.3390/nu13010097> PMID:33396690

16. Park KH, Kim AR, Yang MA, Lim SJ, Park JH. Impact of the COVID-19 pandemic on the lifestyle, mental health, and quality of life of adults in South Korea. *PLoS One*. 2021 Feb 26;16(2):e0247970. <https://doi.org/10.1371/journal.pone.0247970> PMID:33635897
17. Rawat D, Dixit V, Gulati S, Gulati S, Gulati A. Impact of COVID-19 outbreak on lifestyle behaviour: a review of studies published in India. *Diabetes Metab Syndr*. 2021 Jan–Feb;15(1):331–6. <https://doi.org/10.1016/j.dsx.2020.12.038> PMID:33493852
18. Kuang, J., Ashraf, S., Das, U., & Bicchieri, C.). Awareness, risk perception, and stress during the COVID-19 pandemic in communities of Tamil Nadu, India. *Int J Environ Res Public Health*. 2020 Sep 30;17(19):7177. <https://doi.org/10.3390/ijerph17197177> PMID:33007992
19. Romero-Blanco C, Rodríguez-Almagro J, Onieva-Zafra MD, Parra-Fernández ML, Prado-Laguna MdC, Hernández-Martínez A. Sleep pattern changes in nursing students during the COVID-19 lockdown. *Int J Environ Res Public Health*. 2020 Jul 20;17(14):5222. <https://doi.org/10.3390/ijerph17145222> PMID:32698343
20. Van der Werf ET, Busch M, Jong MC, Hoenders H. J. Lifestyle changes during the first wave of the COVID-19 pandemic: a cross-sectional survey in the Netherlands. *BMC Public Health*. 2021 Jun 25 21(1):1226. <https://doi.org/10.1186/s12889-021-11264-z> PMID:34172042
21. Husain W, Ashkanani F. Does COVID-19 change dietary habits and lifestyle behaviours in Kuwait: a community-based cross-sectional study. *Environ Health Prev Med*. 2020 Oct 12;25(1):61. <https://doi.org/10.1186/s12199-020-00901-5> PMID:33045996
22. Ruiz-Roso MB, de Carvalho Padilha P, Mantilla-Escalante DC, Ulloa N, Brun P, Acevedo-Correa D, et al. Covid-19 confinement and changes of adolescent's dietary trends in Italy, Spain, Chile, Colombia and Brazil. *Nutrients*. 2020 Jun 17;12(6):1807. <https://doi.org/10.3390/nu12061807> PMID:32560550
23. Liu S, Yang L, Zhang C. Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry* 2020 Apr;7(4):e17–18. [https://doi.org/10.1016/S2215-0366\(20\)30077-8](https://doi.org/10.1016/S2215-0366(20)30077-8) PMID:32085841
24. Giuntella O, Hyde K, Saccardo S, Sadoff S. Lifestyle and mental health disruptions during COVID-19. *Proc Natl Acad Sci U S A*. 2021 Mar 2;118(9):e2016632118. <https://doi.org/10.1073/pnas.2016632118> PMID:33571107
25. Smith G. Does gender influence online survey participation?: A record-linkage analysis of university faculty online survey response behavior. ERIC Document Reproduction Service No ED 501717. 2008. <https://files.eric.ed.gov/fulltext/ED501717.pdf>