

# Ecological study of breast cancer incidence among nationals and non-nationals in the Gulf Cooperation Council countries

Karima Chaabna,<sup>1</sup> Heta Ladumor<sup>2</sup> and Sohaila Cheema<sup>1</sup>

<sup>1</sup>Institute for Population Health, Weill Cornell Medicine – Qatar, Doha, Qatar (Correspondence to: Karima Chaabna: kac2047@qatar-med.cornell.edu).

<sup>2</sup>Alumni Affairs, Weill Cornell Medicine – Qatar, Doha, Qatar.

## Abstract

**Background:** Breast cancer incidence is increasing in the Gulf Cooperation Council (GCC) countries: Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, and United Arab Emirates.

**Aims:** This study analysed geographical patterns, time trends, and age distribution of female breast cancer incidence among nationals and non-nationals in GCC countries.

**Methods:** Available cancer registry data for 1979–2016 were retrieved for the GCC countries. Age-standardized rates (ASRs) per 100 000 women were calculated using the World standard population. Comparisons were made by calculating comparative incidence figures.

**Results:** From 1998 to 2012, incidence among nationals was highest in Bahrain (ASR 61.85), Kuwait (ASR 52.66), and Qatar (ASR 56.90) and lowest in Saudi Arabia (ASR 19.76), Oman (ASR 22.33), and United Arab Emirates (ASR 31.05). In the most recent period, data were available only in Qatar (2014–2016) and Saudi Arabia (2013–2015). Non-nationals and nationals in Qatar had higher incidence rates than in Saudi Arabia. Incidence among nationals in Qatar was at least twice that in Saudi Arabia (comparative incidence figure 2.32). Incidence among non-nationals in Qatar was 3 times higher than in Saudi Arabia. Among nationals in Kuwait, 10.8% of cases of breast cancer occurred in women aged < 40 years in 2008–2012, compared with 24.2% in non-nationals in Qatar in 2014–2016.

**Conclusion:** Breast cancer incidence has increased over time among women in most GCC countries, likely reflecting the improvements in healthcare access and screening programmes. Nationals and non-nationals developed breast cancer at a younger age than women in other high-income countries. Increased screening uptake is still required in the region. Evidence-based, locally-informed interventions should be implemented to address risk factors specific to the nationals and non-nationals in the GCC countries.

Keywords: breast cancer, cancer incidence, cancer screening, mammography, cancer prevention, Gulf Cooperation Council, nationals, non-nationals

Citation: Chaabna K; Ladumor H; Cheema S. Ecological study of breast cancer incidence among nationals and non-nationals in the Gulf Cooperation Council countries. *East Mediterr Health J.* 2023;29(1):40–48. <https://doi.org/10.26719/emhj.23.005>

Received: 19/10/21; accepted: 19/10/22

Copyright © Authors 2023; Licensee: World Health Organization. EMHJ is an open access journal. This paper is available under the Creative Commons Attribution Non-Commercial ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

## Introduction

Breast cancer is the second most common cancer and the leading cause of cancer-related mortality in women worldwide (1). Although there has been an improvement in the overall survival rate of breast cancer, there are differences by country and region. Limited screening, diagnosis, and therapy are likely contributing factors (2).

The Gulf Cooperation Council (GCC) is an alliance of 6 countries: Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, and United Arab Emirates (UAE). These countries have rich endowments of natural resources, generous government-supported subsidies and welfare policies, large populations of expatriate workers, and common cultural, religious, and social heritage (3, 4). To support economic development in the past 2 decades, the GCC countries have experienced substantial population growth due to mass migration of temporary workers (4). Intense non-national population growth has reached > 80% of the population size in Qatar and UAE. Young healthy migrants have contributed mainly to the non-national population growth, which has skewed the age

distribution towards the 15–49 years group (5). In parallel, there has been an increased adoption of westernized and sedentary lifestyles in the GCC countries (6) that has been linked to a variety of cancers including breast cancer (7). Breast cancer incidence is 3–5 times lower among nationals in GCC countries than in other high-income countries (8); however, incidence among GCC nationals increased by 40% between 1998 and 2009 (6). To the best of our knowledge, incidence has not been assessed separately for GCC nationals and non-nationals. As a result of the unique demographic profile of non-nationals in the region and the potential healthy migrant effect on trends (3,4), it is essential to analyse breast cancer incidence time trends, geographical patterns, and age distribution according to nationality.

This study aimed to characterize female breast cancer incidence in GCC countries. The study objectives were to: (1) analyse geographical patterns and time trends of female breast cancer incidence among GCC nationals and non-nationals separately; (2) explore incidence differences between nationals and non-nationals; and

(3) assess breast cancer age distribution among nationals and non-nationals.

## Methods

### Study design

We used an ecological study design to evaluate breast cancer incidence among nationals and non-nationals in GCC countries.

### Incidence data sources

We retrieved publicly available secondary cancer registry data from 4 sources (Table 1): International Agency for Research on Cancer (IARC); Gulf Centre for Cancer Control & Prevention (GCCCP); Qatar National Cancer Registry (QNCR); and Saudi National Cancer Registry (SNCR). IARC and GCCCP report cancer surveillance data collected by national cancer registries (Bahrain Cancer Registry, Kuwait Cancer Registry, Oman National Cancer Registry, QNCR, SNCR, and United Arab Emirates Cancer Registry). Cancer registry incidence data for Oman were also obtained from the report by Al-Lawati et al. (9). Data from GCCCP, QNCR, SNCR, and Al-Lawati et al. were primarily used to supplement missing information from IARC to allow for a comprehensive and up-to-date description of breast cancer incidence in GCC countries. No data estimated from modelling analysis (e.g. GLOBOCAN 2020) were used in our analysis, which used only real data.

GCCCP reported 15-year incidence data (1998–2012) for nationals of GCC countries (10). These data were used to analyse geographical patterns of incidence across the countries between 1998 and 2012. The IARC Cancer in Five Continents (CI5) database was searched for available data on breast cancer incidence for GCC nationals and non-nationals, without time restriction (Table 1). Data were not available in the IARC database for UAE and for non-nationals of Bahrain, Oman, Qatar, and Saudi Arabia. From the QNCR, we retrieved separate incidence data for nationals and non-nationals in Qatar for 2014–2016. From the SNCR, we retrieved separate incidence data for nationals and non-nationals in Saudi Arabia for 2013–2015. Al-Lawati et al. (9) reported 20-year incidence data (1996–2016) for nationals in Oman, which were used to supplement the minimal data available for Oman from IARC. The retrieved incidence data were used to further analyse geographical patterns, and to investigate time trends (1979–2016), age distribution, and national versus non-national incidence.

### Statistical analysis

To conduct a valid comparison between populations with different age structures, we calculated the age-standardized rate (ASR), which

Table 1 Breast cancer incidence data sources in Gulf Cooperation Council countries

Country	Bahrain		Kuwait		Oman		Qatar		Saudi Arabia		United Arab Emirates	
	Nationals	Non-nationals	Nationals	Non-nationals	Nationals	Non-nationals	Nationals	Non-nationals	Nationals	Non-nationals	Nationals	Non-nationals
Data sources												
Year of data collection												
Al-Lawati et al.	–	–	–	–	✓	–	–	–	–	–	–	–
GCCCP	✓	–	✓	–	✓	–	–	–	✓	–	–	–
SNCR	–	–	–	–	–	–	–	–	✓	–	–	–
QNCR	–	–	–	–	–	–	–	✓	–	–	–	–
IARC CI5 Volume XI <sup>a</sup>	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume X	✓	–	✓	–	–	–	–	–	–	–	–	–
IARC CI5 Volume IX	✓	–	✓	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VIII	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VII	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VII	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VII	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VII	–	–	–	–	–	–	–	–	–	–	–	–
IARC CI5 Volume VII	–	–	–	–	–	–	–	–	–	–	–	–

IARC = International Agency for Research on Cancer; CI5 = Cancer in Five Continents; GCCCP = Gulf Centre for Cancer Control & Prevention; QNCR = Qatar National Cancer Registry; SNCR = Saudi National Cancer Registry.

✓ data available for that population and time period.

– data unavailable for that population and time period.

<sup>a</sup>Eleven volumes of CI5 have been published by IARC. Volumes VII to IX are available on IARC websites. Each volume reports cancer registry incidence data for a specific period.

<sup>b</sup>For Oman, the time period was 1993–1997 in Volume VIII and 1998–2001 in Volume IX.

is the weighted average of age-specific incidence rates using the direct standardization method with the World Standard Population (11). Incidence rates were standardized per 100 000 women. Age-specific incidence rate was calculated as number of new breast cancer cases per 100 000 women for a specific period, country, nationality, and narrow age range. The number of cases was extracted by sex, nationality, and 5-year-age groups (0 to > 75 or > 85 years). The comparative incidence figure (CIF) and its 95% confidence interval (CI) were calculated to assess ASR differences according to time period, country, and nationality (11). CIF was a ratio of the ASR in a specific period, country, and nationality compared with the ASR in another period, country, and nationality. We considered the difference to be significant if the value of 1.0 was not included within the 95% CI of CIF.

The population at risk for female breast cancer included all individuals susceptible to breast cancer. Person-years at risk by nationality and age group were retrieved from the same cancer incidence data source, except for the data from QNCR (2014–2016). Person-years at risk were calculated by the sum of population size in women by age group and nationality for all 3 years retrieved from the Labor Force Survey conducted by the Ministry of Development Planning and Statistics (<https://www.psa.gov.qa/en/statistics1/Pages/default.aspx>).

## Results

### Geographic patterns of breast cancer incidence in GCC countries

Over the period 1998–2012, breast cancer incidence in nationals was highest in Bahrain (ASR = 61.85), while it was 3 times lower in Saudi Arabia (ASR = 19.76; CIF = 3.13; 95% CI: 3.12932–3.12933) and 2 times lower in UAE (ASR


= 31.05; CIF = 1.99; 95% CI: 1.9919–1.9920). Breast cancer incidence among nationals was 56.90, 52.66, and 22.33 per 100 000 in Qatar, Kuwait, and Oman, respectively.

The geographical pattern over shorter periods of time differed from that over the 15-year period (Figure 1). Until 2003–2007, nationals in Bahrain had the highest ASRs when compared with the nationals of other GCC countries (except for UAE where data were not available) and compared with non-nationals in Kuwait. Thereafter, incidence among nationals in Bahrain declined in 2008–2012 and was lower than among nationals in Kuwait (CIF = 0.87; 95% CI: 0.86594–0.86595) and Qatar (CIF = 0.93; 95% CI: 0.93607–0.93608) but still higher than among nationals in Saudi Arabia (CIF = 2.1; 95% CI: 2.051829–2.051833) and among non-nationals in Kuwait (CIF = 1.22; 95% CI: 1.221258–1.221261). In the most recent period, data were available only in Qatar (2014–2016) and Saudi Arabia (2013–2015). Non-nationals and nationals in Qatar had higher incidence than non-nationals and nationals in Saudi Arabia. Breast cancer incidence among nationals in Qatar was at least twice that of nationals in Saudi Arabia in 2003–2007 (CIF = 2.17; 95% CI: 2.16753–2.16754), 2008–2012 (CIF = 2.19; 95% CI: 2.19194–2.19195), and 2013–2016 (CIF = 2.32; 95% CI: 2.32574–2.32575). Incidence among non-nationals in Qatar was 3 times higher than among non-nationals in Saudi Arabia.

### Time trends of breast cancer incidence in GCC countries

Overall, breast cancer ASR increased in all GCC countries (Figure 1). However, a decline in the trends (among nationals in Qatar and Saudi Arabia) and even a decrease (among nationals in Bahrain and non-nationals in Kuwait) was observed in recent years. Breast cancer incidence among nationals in Bahrain increased between

**Figure 1** Breast cancer age-standardized rate per 100 000 women for Gulf Cooperation Council country nationals and non-nationals between 1979 and 2016.



1998–2002 and 2003–2007 (CIF = 1.2; 95% CI: 1.196026–1.196030) and decreased between 2003–2007 and 2008–2012 (CIF = 0.9; 95% CI: 0.899017–0.899019). Breast cancer incidence among nationals in Kuwait doubled between 1979–1982 and 1988–1993 (CIF = 2.01), remained the same until 1994–1997 (CIF = 1.0; 95% CI: 1.00030–1.00031), and increased between 1994–1997 and 1998–2002 (CIF = 1.26; 95% CI: 1.257689–1.257693), 1998–2002 and 2003–2007 (CIF = 1.21; 95% CI: 1.211430–1.211433, and 2003–2007 and 2008–2012 (CIF = 1.16; 95% CI: 1.162534–1.162536). Breast cancer incidence among non-nationals in Kuwait increased between 1979–1982 and 1988–1993, remained the same until 1994–1997 (CIF = 1.0; 95% CI: 0.998832–0.998837), increased between 1994–1997 and 2003–2007 (CIF = 1.3; 95% CI: 1.302703–1.302708), and decreased in 2008–2012 (CIF = 0.92; 95% CI: 0.922989–0.922992). Breast cancer incidence among nationals in Oman increased between 1993–1997 and 1998–2001 (CIF = 1.15; 95% CI: 1.146130–1.146132). The ASR (22.33) estimated for 1996–2015 using data reported by Al-Lawati et al. (9) was similar to that in 1998–2012 (ASR = 20.75). Data for breast cancer incidence in Qatar were not available before 2003. The incidence among nationals increased between 2003–2007 and 2008–2012 (CIF = 1.18; 95% CI: 1.175398–1.175403) and between 2008–2012 and 2014–2016 (CIF = 1.04; 95% CI: 1.039074–1.039079). Data for breast cancer incidence in Saudi Arabia were not available prior to 2003. The incidence among nationals increased between 2003–2007 and 2008–2012 (CIF = 1.16; 95% CI: 1.162481–1.162483) and slightly decreased thereafter (2014–2016 CIF = 0.98).

### Breast cancer incidence among nationals versus non-nationals

Breast cancer incidence in Kuwait was significantly lower among nationals than non-nationals and the gap between the 2 populations diminished until 1994–1997 (1988–1993: CIF = 0.96, 95% CI: 0.956013–0.956017; 1994–1997: CIF = 0.96, 95% CI: 0.957419–0.957424). Thereafter, breast cancer incidence was higher among nationals than non-nationals and the gap increased over time (CIF = 1.06;

95% CI: 1.056254–1.056257, in 1998–2002 and CIF = 1.41; 95% CI: 1.410320–1.410323 in 2008–2012). Breast cancer incidence in Qatar was 50% lower among nationals than non-nationals during 2014–2016 (CIF = 0.50; 95% CI: 0.502528–0.502530; ASR = 55.89 and 111.23, respectively). Incidence in Saudi Arabia was lower among nationals than non-nationals during 2013–2015 (CIF = 0.69; 95% CI: 0.687872–0.687873; ASR = 24.0 and 34.9, respectively).

### Breast cancer incidence by age group


Breast cancer incidence increased with age among nationals and non-nationals (Figure 2). No breast cancer was reported among nationals or non-nationals aged < 20 years, except among non-nationals in Qatar (age-specific incidence rate = 1.8). Among women aged < 40 years, breast cancer cases accounted for between 10.8% of all cases among nationals in Kuwait in 2008–2012, and 24.2% of all cases among non-nationals in Qatar in 2014–2016 (Figure 3).

## Discussion

Breast cancer incidence was highest among nationals and non-nationals in Qatar, followed by nationals in Kuwait and Bahrain, nationals and non-nationals in Saudi Arabia, and nationals in Oman and UAE. In recent years, the incidence of breast cancer increased in all GCC countries but appeared to have decreased among nationals in Qatar and Saudi Arabia, and even reversing among nationals in Bahrain and non-nationals in Kuwait. Breast cancer incidence data for non-nationals were available only for Kuwait, Qatar, and Saudi Arabia. Non-nationals in Qatar and Saudi Arabia were more likely to have breast cancer than nationals, whereas in recent years, nationals in Kuwait were more likely than non-nationals to have breast cancer. Nationals and non-nationals in GCC countries appeared to develop breast cancer at a younger age than women in other high-income countries.


Several factors likely explain why breast cancer incidence in GCC countries was lower than in other

**Figure 2 Breast cancer age-specific incidence rate per 100 000 women in Gulf Cooperation Council country nationals and non-nationals.**





**Figure 3** Percentage of all breast cancer cases occurring in Gulf Cooperation Council country national and non-national women aged < 40 years.



high-income countries. Estimates of cancer incidence are closely dependent on levels of screening and detection. There is limited information on effectiveness of national-level breast cancer detection in GCC countries (14). In 2008–2009, the proportion of women aged 40–75 years who had undergone mammography within the previous 2 years was low among nationals and non-nationals (<15%) (14). Therefore, incidence differences between GCC countries and other high-income countries are likely to diminish as the proportion of women undergoing mammography increases. Thus, while other high-income countries are beginning to consider reduced routine screening to avoid harm caused by excessive screening, in the GCC countries, increased participation in screening programmes may be critical for breast cancer detection and saving lives (17).

Few studies have explored the barriers and enhancers of screening programmes in the region. Cultural, practical, and personal barriers have been reported (18–20). In GCC countries, participation in screening programmes may be low because of fear of treatment, doctors/examiners, hospitals, and consequences (18,21); embarrassment during breast examination (18,19); lack of awareness and knowledge about screening programmes (19,22) and doctor's recommendations (18,19); and concerns over consulting a male or a non-Arabic-speaking doctor (19). Conversely, feelings of susceptibility and a supportive social environment were facilitators for screening in the region (20). Therefore, awareness campaigns aimed at enhancing knowledge and changing perceptions about breast cancer and the implementation of culturally sensitive screening programmes are necessary for tackling the breast cancer burden in the region.

Despite the likelihood that low screening rates may lead to underestimation of breast cancer incidence in GCC countries, the increase in incidence may reflect improved screening practices and better access to healthcare systems in recent years, compared with the period before the socioeconomic development in the region (23). Concomitantly, improvements in the healthcare system has resulted in more robust reporting of epidemiological statistics (24). Additional incidence data to assess the

recent trends, and evaluation of screening programmes are required.

Time trends of the breast cancer stage may clarify whether the observed increase in breast cancer incidence over time was a true increase (observed at all stages) or the result of earlier diagnosis (increase mainly in early stages). Unfortunately, there were limited data about the specific stage of breast cancer diagnosed over time for GCC countries (25,26).

Breast cancer incidence was compared between nationals and non-nationals in Kuwait, Qatar, and Saudi Arabia. GCC country nationals have free access to all healthcare services (28), but employers must pay annual fees that enable non-national workers to access healthcare services at a reduced cost (29). The exception is in Oman, where non-nationals must pay to access public and private health services other than emergency services (29). In Qatar, despite similar access to the healthcare system for nationals and non-nationals, we observed a higher breast cancer incidence among non-nationals – similar to that observed in other high-income countries – probably because of greater awareness of breast cancer symptoms, more willingness to seek medical advice, and higher mammography screening uptake by non-nationals than nationals. This hypothesis needs to be investigated in future research.

Nationals and non-nationals in GCC countries seemed to have developed breast cancer at a younger age than women in other high-income countries. Breast cancer incidence was higher among women aged < 40 years in GCC countries than other high-income countries (31). Less variability among GCC countries and between nationals and non-nationals was observed in this age group. Cancer screening guidelines developed in the region are similar to international guidelines (32). Therefore, the higher incidence of breast cancer in women aged < 40 years was less likely because of early detection through screening. There is a need to investigate the higher proportion of national and non-national women diagnosed during the premenopausal period in GCC countries than in other high-income countries.

A study strength was the use of cancer registry data over a long period of 38 years. In comparison, previous studies evaluated patterns and trends over 5–10-year periods and did not analyse available data for non-nationals (6,8,35). Albeshan et al. (8) used the IARC modelled GLOBOCAN incidence estimation for 2008 and 2012 instead of cancer registry incidence data. This conflicts with the recommendation of IARC that GLOBOCAN incidence estimations in 2008 and 2012 should not be compared because of methodological differences in calculating these estimates (12).

Limitations included the lack of data on breast cancer incidence among non-nationals. While the data on non-nationals in Kuwait spanned 24 years, those on non-nationals in Qatar and Saudi Arabia were available only for 3 years, precluding in-depth exploration. Therefore, there is a need for long-term data from GCC countries to investigate incidence differences between nationals and non-nationals. The IARC CI5 database contained no data relating to nationals in UAE, either because of poor quality or lack of reporting to IARC. Hence, the analysis conducted on UAE may have been of lower precision than that in the other GCC countries.

The quality of cancer-related data was limited in GCC countries than other high-income countries. The proportion of breast cancer cases that were microscopically verified and histologically confirmed cases was >97% in the region (36). Despite suggesting good data validity, such a high proportion of microscopically verified breast cancer cases may be a sign of registry incompleteness because of a lack of use of other case-finding procedures (37), and that the GCC registries are

over-reliant on pathology laboratories as the primary source of information (36). This seemed to be confirmed by the percentage of breast cancer cases identified during necropsy examination of women in whom breast cancer had not been diagnosed during their lifetime (< 1% in GCC countries). Consequently, cases detected by other procedures such as death certification were probably missing within the dataset (36). Comprehensive case-finding procedures including microscopic verification and death certificates is required for GCC cancer registries to improve their data quality (36).

## Conclusion

Female breast cancer incidence has increased over time among most GCC countries, likely reflecting improvements in healthcare access and screening programmes. However, in contrast to other high-income countries, increased participation in screening continues to be critical for the region. GCC nationals and non-nationals appeared to have developed breast cancer at a younger age than women in other high-income countries. Risk factors specific to nationals and non-nationals in GCC countries should be further investigated. Evidence-based, locally informed interventions addressing these factors must be implemented. Educational interventions should be implemented to increase awareness about breast cancer and eliminate associated barriers. Data collection related to breast cancer among nationals and non-nationals must be enhanced in the region to fully understand its epidemiology and improve national guidelines.

## Acknowledgement

We would like to thank QNCR, Ministry of Public Health – Qatar, and Qatar Cancer Information Center for providing us data on breast cancer in Qatar. We would also like to acknowledge Ms. Danielle Jones from the Writing Center and Dr. Ross MacDonald from the Library of Weill Cornell Medicine – Qatar, who helped with editing.

**Funding:** None.

**Competing interests:** None declared.

## Étude écolologique de l'incidence du cancer du sein chez les ressortissantes et les non-ressortissantes des pays membres du Conseil de coopération du Golfe

### Résumé

**Contexte :** L'incidence du cancer du sein est en augmentation dans les pays membres du Conseil de coopération du Golfe (CCG) : Arabie saoudite, Bahreïn, Émirats arabes unis, Koweït, Oman et Qatar.

**Objectifs :** La présente étude a permis d'analyser les schémas géographiques, les tendances dans le temps et la répartition selon l'âge de l'incidence du cancer du sein chez les ressortissantes et les non-ressortissantes des pays membres du CCG.

**Méthodes :** Les données disponibles issues des registres du cancer pour la période 1979-2016 ont été extraites pour les pays membres du CCG. Les taux standardisés selon l'âge (TSA) pour 100 000 femmes ont été calculés sur la base de la population mondiale standard. Les comparaisons ont été faites en calculant les taux d'incidence comparatifs.

**Résultats :** Entre 1998 et 2012, l'incidence parmi les ressortissantes était la plus élevée à Bahreïn (TSA : 61,85), au Koweït (TSA : 52,66) et au Qatar (TSA : 56,90) et la plus faible en Arabie saoudite (TSA : 19,76), à Oman (TSA : 22,33) et aux Émirats arabes unis (TSA : 31,05). Au cours de la période la plus récente, seules des données concernant le

Qatar (2014-2016) et l'Arabie saoudite (2013-2015) étaient disponibles. Les ressortissantes et les non-ressortissantes du Qatar présentaient des taux d'incidence plus élevés que celles de l'Arabie saoudite. L'incidence parmi les ressortissantes du Qatar était au moins deux fois plus élevée qu'en Arabie saoudite (voir la figure 2.32 sur l'incidence comparative), et celle parmi les non-ressortissantes était trois fois plus élevée qu'en Arabie saoudite. Parmi les ressortissantes du Koweït, 10,8 % des cas de cancer du sein sont survenus chez des femmes âgées de moins de 40 ans entre 2008 et 2012, tandis que ce taux était de 24,2 % chez les non-ressortissantes du Qatar pour la période 2014-2016.

**Conclusion :** L'incidence du cancer du sein a augmenté au fil du temps chez les femmes dans la plupart des pays membres du CCG, reflétant probablement les progrès réalisés en matière d'accès aux soins de santé et de programmes de dépistage. Les ressortissantes et les non-ressortissantes ont contracté le cancer du sein à un âge plus précoce que les femmes dans d'autres pays à revenu élevé. Une augmentation du taux de dépistage est encore nécessaire dans la Région. Des interventions fondées sur des données probantes et adaptées au contexte local devraient être mises en œuvre pour s'attaquer aux facteurs de risque spécifiques aux ressortissantes et aux non-ressortissantes des pays membres du CCG.

## دراسة إيكولوجية لحالات الإصابة بسرطان الثدي بين المواطنين وغير المواطنين في دول مجلس التعاون الخليجي

كريمة شعبانه، هيتا لادمور، سهيلة شيبا

### الخلاصة

الخلفية: تزايد حالات الإصابة بسرطان الثدي في دول مجلس التعاون الخليجي: البحرين، والكويت، وعمان، وقطر، والمملكة العربية السعودية، والإمارات العربية المتحدة.

الأهداف: هدفت هذه الدراسة الى تحليل الأنماط الجغرافية والاتجاهات الزمنية والتوزيع العمري للإصابة بسرطان الثدي لدى الإناث بين المواطنين وغير المواطنين في دول مجلس التعاون الخليجي.

طرق البحث: جرى استخلاص بيانات سجل السرطان المتاحة عن المدة بين عامي 1979-2016 لدول مجلس التعاون الخليجي. وحُسبت المعدلات الموحدة حسب السن لكل 100000 امرأة باستخدام المعيار العالمي للسكان. وأجريت مقارنات بحساب أرقام معدل الإصابة المقارنة.

النتائج: بالنسبة إلى الفترة الزمنية ما بين عامي 1998-2012، تبين أن أعلى معدل للإصابة بين المواطنين في دول مجلس التعاون الخليجي كان في كل من البحرين (معدل موحّد حسب السن 61.85)، والكويت (معدل موحّد حسب السن 52.66)، وقطر (معدل موحّد حسب السن 56.90) وتبين أن أدنى معدل للإصابة كان في المملكة العربية السعودية (معدل موحّد حسب السن 19.76)، وعمان (معدل موحّد حسب السن 22.33)، والإمارات العربية المتحدة (معدل موحّد حسب السن 31.05). ولكن في الفترة الأخيرة، لم تتوفر سوى البيانات الخاصة بقطر (2014-2016) والمملكة العربية السعودية (2013-2015). وسجلت المواطنين وغير المواطنين في قطر معدلات إصابة أعلى منها في المملكة العربية السعودية. وبلغ معدل الإصابة بين المواطنين في قطر ضعف مثيله على الأقل في المملكة العربية السعودية (الشكل 32.2 لمعدل الإصابة المقارنة). أما معدل الإصابة بين غير المواطنين في قطر، فجاء أعلى 3 مرات مقارنة بالمعدل في المملكة العربية السعودية. وحدثت 10.8% من حالات الإصابة بسرطان الثدي بين المواطنين في الكويت بين النساء اللاتي تقل أعمارهن عن 40 عامًا في المدة بين عامي 2008-2012، مقارنة بنسبة 24.2% بين غير المواطنين في قطر في المدة بين عامي 2014-2016.

الاستنتاجات: ازدادت معدلات الإصابة بسرطان الثدي مع مرور الوقت بين النساء في معظم دول مجلس التعاون الخليجي، الأمر الذي يعكس على الأرجح تحسّسات في الوصول إلى الرعاية الصحية وبرامج الفحص. وتصاب المواطنين وغير المواطنين بسرطان الثدي في سن أصغر من سن النساء في البلدان الأخرى ذات الدخل المرتفع. ولا يزال من المطلوب زيادة الإقبال على الفحص في المنطقة. لذا، ينبغي تنفيذ تدخلات مُسندة بالبيّنات ومستنيرة محليًا للتصدي لعوامل الخطر الخاصة تحديداً بالمواطنين وغير المواطنين في دول مجلس التعاون الخليجي.

## References

1. Fitzmaurice C, Abate D, Abbasi N, Abbastabar H, Abd-Allah F, Abdel-Rahman O, et al. Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 29 cancer groups, 1990 to 2017: a systematic analysis for the Global Burden of Disease Study. *JAMA Oncology*. 2019 Dec;5(12):1749–68. <https://doi.org/10.1001/jamaoncol.2019.2996> PMID:31560378
2. Lin L, Yan L, Liu Y, Yuan F, Li H, Ni J. Incidence and death in 29 cancer groups in 2017 and trend analysis from 1990 to 2017 from the Global Burden of Disease Study. *J Hematol Oncol*. 2019 Sep 12;12(1):96. <https://doi.org/10.1186/s13045-019-0783-9> PMID:31511035
3. Chaabna K, Cheema S, Abraham A, Alrouh H, Mamtani R. Adult mortality trends in Qatar, 1989-2015: National population versus migrants. *PLoS One*. 2018 Sep 25;13(9):e0203996. <https://doi.org/10.1371/journal.pone.0203996> PMID:30252887
4. Chaabna K, Cheema S, Mamtani R. Migrants, healthy worker effect, and mortality trends in the Gulf Cooperation Council countries. *PLoS One*. 2017 Jun 20;12(6):e0179711. <https://doi.org/10.1371/journal.pone.0179711> PMID:28632794

5. Total population – both sexes. De facto population in a country, area or region as of 1 July of the year indicated. Figures are presented in thousands. World population prospects: the 2015 revision. United Nations; 2016.
6. Al-Othman S, Haoudi A, Alhomoud S, Alkhenizan A, Khoja T, Al-Zahrani A. Tackling cancer control in the Gulf Cooperation Council Countries. *Lancet Oncol.* 2015 May;16(5):e246–57. [https://doi.org/10.1016/S1470-2045\(15\)70034-3](https://doi.org/10.1016/S1470-2045(15)70034-3) PMID:25943069
7. Moore SC, Lee IM, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA internal medicine.* 2016 Jun 1;176(6):816–25. <https://doi.org/10.1001/jamainternmed.2016.1548> PMID:27183032
8. Albeshan SM, Mackey MG, Hossain SZ, Alfuraih AA, Brennan PC. Breast Cancer Epidemiology in Gulf Cooperation Council countries: a regional and international comparison. *Clin Breast Cancer.* 2018 Jun;18(3):e381–92. <https://doi.org/10.1016/j.clbc.2017.07.006> PMID:28781021
9. Al-Lawati NA, Al-Bahrani BJ, Al-Raisi SS, Al-Lawati JA. Twenty-year trends of cancer incidence in Omanis, 1996–2015. *Oman Med J.* 2019 Jul;34(4):361–87. PMID:31360330
10. 15-year cancer incidence among nationals of GCC states 1998–2012. Gulf Center for Cancer Control & Prevention; 2015.
11. Boniol M, Heanue M. Age-standardisation and denominators. In: Curado MP, Edwards B, Shin HR, Storm H, Ferlay J, Heanue M, et al., editors. *Cancer incidence in five continents, volume IX.* Lyon: International Agency for Research on Cancer; 2007:99–101.
12. So VHT, Channon AA, Ali MM, Merdad L, Al Sabahi S, Al Suwaidi H, et al. Uptake of breast and cervical cancer screening in four Gulf Cooperation Council countries. *Eur J Cancer Prev.* 2019 Sep;28(5):451–6. <https://doi.org/10.1097/CEJ.0000000000000466> PMID:30260812
13. Crunching numbers: what cancer screening statistics really tell us [website]. National Cancer Institute; 2018 (<https://www.cancer.gov/about-cancer/screening/research/what-screening-statistics-mean>, accessed 8 December 2022).
14. Al-Azri M, Al-Rubaie K, Al-Ghafri S, Al-Hinai M, Murthi Panchatcharam S. Barriers and attitudes toward breast cancer screening among Omani women. *Asian Pac J Cancer Prev.* 2020 May 1;21(5):1339–47. <https://doi.org/10.31557/APJCP.2020.21.5.1339> PMID:32458642
15. Donnelly TT, Al Khater AH, Al-Bader SB, Al Kuwari MG, Al-Meer N, Malik M, et al. Beliefs and attitudes about breast cancer and screening practices among Arab women living in Qatar: a cross-sectional study. *BMC Women's Health.* 2013 Dec 13;13:49. <https://doi.org/10.1186/1472-6874-13-49> PMID:24330708
16. Bener A, Honein G, Carter AO, Da'ar Z, Miller C, Dunn EV. The determinants of breast cancer screening behavior: a focus group study of women in the United Arab Emirates. *Oncol Nurs Forum.* 2002 Oct;29(9):E91–8. <https://doi.org/10.1188/02.ONF.E91-E98> PMID:12370705
17. Abdel-Aziz SB, Amin TT, Al-Gadeeb MB, Alhassar AI, Al-Ramadan A, Al-Helal M, et al. Perceived barriers to breast cancer screening among Saudi women at primary care setting. *J Prev Med Hyg.* 2018 Mar 20;59(1):E20–9. <https://doi.org/10.15167/2421-4248/jpmh2018.59.1.689> PMID:29938236
18. Renganathan L, Ramasubramaniam S, Al-Touby S, Seshan V, Al-Balushi A, Al-Amri W, et al. What do Omani women know about breast cancer symptoms? *Oman Med J.* 2014 Nov;29(6):408–13. <https://doi.org/10.5001/omj.2014.110> PMID:25584157
19. Goodman A. The development of the Qatar healthcare system: a review of the literature. *Int J Clin Med.* 2015 Mar;6(3):177–85. <https://doi.org/10.4236/ijcm.2015.63023>
20. Wong MCS, Lao XQ, Ho KF, Goggins WB, Tse SLA. Incidence and mortality of lung cancer: global trends and association with socioeconomic status. *Sci Rep.* 2017 Oct 30;7(1):14300. <https://doi.org/10.1038/s41598-017-14513-7> PMID:29085026
21. Cancer Incidence Report Saudi Arabia 2015. Saudi Arabia: Saudi Health Council, National Health Information Center, Saudi Cancer Registry; 2015.
22. Qatar National Cancer Registry. Breast cancer incidence in females in Qatar. Ministry of Public Health; 2015.
23. Gulf Labour Markets and Migration (GLMM). Demography, migration and labour market in Qatar. GLMM–EN–No. 8/2014. San Domenico di Fiesole, Italy: European University Institute and Gulf Research Center; 2015 ([https://gulfmigration.grc.net/media/pubs/exno/GLMM\\_EN\\_2017\\_03.pdf](https://gulfmigration.grc.net/media/pubs/exno/GLMM_EN_2017_03.pdf), accessed 8 December 2022).
24. Comparison of health care coverage for migrant workers in the GCC [website]. migrant-rights.org; 2020 (<https://www.migrant-rights.org/wp-content/uploads/2020/03/Comparison-Of-Health-Care-Coverage-For-Migrant-Workers-In-The-GCC-.pdf>, accessed 8 December 2022).
25. Ziegler RG, Hoover RN, Pike MC, Hildesheim A, Nomura AMY, West DW, et al. Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst.* 1993 Nov 17;85(22):1819–27. <https://doi.org/10.1093/jnci/85.22.1819> PMID:8230262
26. Tanner LTA, Cheung KL. Correlation between breast cancer and lifestyle within the Gulf Cooperation Council countries: a systematic review. *World J Clin Oncol.* 2020 Apr 24;11(4):217–42. <https://doi.org/10.5306/wjco.v11.i4.217> PMID:32355643
26. American Cancer Society guidelines for the early detection of cancer [website]. American Cancer Society; 2021 (<https://www.cancer.org/healthy/find-cancer-early/american-cancer-society-guidelines-for-the-early-detection-of-cancer.html>, accessed 8 December 2022).
27. Ten-year cancer incidence among nationals of the GCC States 1998–2007. King Faisal Specialist Hospital and Research Center, Saudi Arabia: Gulf Center for Cancer Control and Prevention; 2011 (<https://www.moh.gov.bh/Content/Files/Publications/GCC%20Cancer%20Incidence%202011.pdf>, accessed 8 December 2022).



28. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018 Nov;68(6):394–424. <https://doi.org/10.3322/caac.21492> PMID:30207593
29. Bray f, Ferlay J. Data comparability and quality. Lyon: International Agency for Research on Cancer; 2020 ([https://publications.iarc.fr/\\_publications/media/download/3751/8b06cadf6dc94192856d21665c8fb29f649ca77a.pdf](https://publications.iarc.fr/_publications/media/download/3751/8b06cadf6dc94192856d21665c8fb29f649ca77a.pdf), accessed 8 December 2022).
30. Chaabna K, Boniol M, de Vuyst H, Vanhems P, Antônio de Ávila Vitoria M, Curado MP. Geographical patterns of Kaposi's sarcoma, nonHodgkin lymphomas, and cervical cancer associated with HIV infection in five African populations. *Eur J Cancer Prev.* 2012 Jan;21(1):1–9. <https://doi.org/10.1097/CEJ.0b013e32834a802a> PMID:21955799