

Cigarette use and exposure to second-hand smoke and tobacco advertising among Tunisian adolescents

Yosr Ayedi¹, Chahida Hariz¹, Afef Skhiri¹ and Radhouane Fakhfakh¹

¹Department of Epidemiology and Biostatistics, Abderrahmane Mami Hospital, Ariana, Tunisia. (Correspondence to Yosr Ayedi: yosr.ayeedi.93@gmail.com).

Abstract

Background: The Global Youth Tobacco Survey was conducted in Tunisia in 2001, 2007, 2010 and 2017. It is the only national survey that examined exposure to second-hand smoke and tobacco advertising among young people, and there has been no description of the trend.

Aims: To describe cigarette use trends among Tunisian adolescents and their exposure to second-hand smoke and tobacco advertising from 2001 to 2017.

Methods: The Global Youth Tobacco Survey is a school-based cross-sectional survey conducted by the World Health Organization. It uses a two-stage cluster sampling design to obtain a representative sample of students aged 13–15 years. A standardized questionnaire is used for data collection. We compared the prevalence and 95% confidence intervals (CI) of ever and current cigarette use, exposure to second-hand smoke in and outside the home, and exposure to tobacco advertising, over 4 years (2001, 2007, 2010 and 2017).

Results: Current cigarette use decreased from 11.1% (95% CI: 10.0–12.3%) in 2001 to 7.7% (95% CI: 6.5–9.0%) in 2017 ($P < 0.001$). Exposure to second-hand smoke at home decreased from 62.5% (95% CI: 60.7–64.2%) to 46.7% (95% CI: 44.5–49.0%) over the same period ($P < 0.001$), but exposure outside the home increased from 65.4% (95% CI: 63.7–67.1%) in 2001 to 73.3% (95% CI: 71.2–75.3%) in 2017 ($P < 0.001$). Exposure to anti-tobacco messages in the media decreased from 87.8% (95% CI: 86.3–89.1%) in 2001 to 64.4% (95% CI: 62.2–66.5%) in 2017 ($P < 0.001$).

Conclusion: While the prevalence of cigarette use and second-hand smoke exposure at home decreased, exposure outside the home increased. Efforts are needed to ensure compliance with smoke-free laws to decrease the prevalence of exposure to second-hand smoke.

Keywords: tobacco, tobacco advertising, smoking, second-hand smoke, pollution, adolescent, Global Youth Tobacco Survey, Tunisia.

Citation: Ayedi Y, Hariz C, Skhiri A, Fakhfakh R. Cigarette use and exposure to second-hand smoke and tobacco advertising among Tunisian adolescents. *East Mediterr Health J.* 2023;29(9):708–715. <https://doi.org/10.26719/emhj.23.075>

Received: 05/09/22; Accepted: 01/04/23

Copyright © Authors 2023; Licensee: World Health Organization. EMHJ is an open access journal. This paper is available under the Creative Commons Attribution Non-Commercial ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Introduction

Tobacco was the main cause of death among males in 2019 worldwide, responsible for 20% of deaths among males. For women, tobacco was the sixth leading cause of death worldwide, responsible for 15.4% of all deaths among women (1). Worldwide, in 2019, about 1.14 billion people aged 15 years and older smoked cigarettes (2). Generally, regular adult smokers began smoking during adolescence and one-third started at 14 years (3). People who smoke their first cigarette before the age of 18 years are more likely to become heavy smokers and nicotine dependent in the future, and are less likely to quit, which puts them at higher risk of lung cancer or other tobacco-induced diseases (4). Considerable effort has been made globally to control tobacco use by helping smokers to quit and preventing smoking initiation.

In 2004, a total of 603 000 deaths were estimated to be related to second-hand smoke; 28% of these deaths occurred among children (6). The increase in people's knowledge of the effects of tobacco use and second-hand smoke as a result of the media and anti-tobacco messages

has helped tobacco control efforts (7). WHO launched the Framework Convention on Tobacco Control (FCTC) in 2003, which was the first international treaty on tobacco control (8). In line with the FCTC, WHO introduced the WHO MPOWER measures: M for monitoring tobacco use and prevention policies, P for protecting people from tobacco smoke; O for offering help to quit tobacco use; W for warning about the dangers of tobacco; E for enforcing bans on tobacco advertising, promotion and sponsorship; and R for raising taxes on tobacco.

Tunisia started a national strategic plan to curb the epidemic of tobacco use among adults and young people in 1998. The strategy was further enforced by Tunisia's ratification of the FCTC in 2010 (10). In Tunisia, the prevalence of smoking among adult males was reported to be 48.3% (95% confidence interval (CI): 46.3–50.3%) by the Tunisian Health Examination Survey in 2016 (9). The Global Youth Tobacco Survey (GYTS) is a main component of the MPOWER plan of action. It is a multinational survey conducted by WHO (11) in more than 185 countries to monitor tobacco use among young people aged 13–15 years (12). In Tunisia, this survey has been conducted 4

times: in 2001, 2007, 2010 and 2017. To our knowledge, the GYTS is the only national survey that examined exposure to second-hand smoke and to the media and advertising among young people.

The aim of our study was to identify the trends in cigarette use among Tunisian adolescents from 2001 to 2017 and to describe their exposure to second-hand smoke and the media and advertising related to tobacco.

Methods

Study design

The GYTS is a cross-sectional, descriptive and school-based survey conducted by WHO. It uses a two-stage cluster sample design to obtain representative samples of students aged 13–15 years. In Tunisia, the age 13–15 years old matches students in the seventh, eighth and ninth school grades. In the GYTS, the complete list of all public schools is sent to the tobacco centre at the United States Centers for Disease Control and Prevention (CDC) where schools are chosen randomly in proportion to the number of students enrolled in the specified grade. Then, classes are randomly chosen according to the city population and size (one or two classes per school).

The GYTS in Tunisia is carried out in April and May of each survey year. Physicians and nurses of medical schools are responsible for data collection, under the direction of the medical school and university. The surveys are funded by WHO. Each student in the age range 13–15 years (seventh, eighth and ninth grades) in the selected classes who is present in the class on the day of survey is eligible to participate in the study.

Questionnaire

The GYTS survey uses a standard methodology and the questionnaire is validated by CDC and WHO experts (13). It contains core questions about the main tobacco concerns focusing on:

- prevalence of all smoked tobacco products and conventional cigarettes
- smokers' access to tobacco products
- smokers' behaviours related to stopping smoking
- exposure to the media and advertising
- exposure to second-hand smoke.

The questionnaire has been translated into Arabic and then re-translated into English and sent back to CDC for further checks to ensure accuracy and reliability. It was first pretested with a focus group of adolescents to ensure the translation was pertinent and precise. The questionnaire contained 69 questions in 2001, 63 questions in 2007, 70 questions in 2010 and 63 in 2017: 27 questions are common to all 4 surveys.

We focused on trends in the prevalence of conventional cigarette smoking and exposure to second-hand smoke and to tobacco advertising.

Measures

Ever cigarette smoker was defined as someone who had ever smoked cigarettes, even if they had only taken 1 or 2 puffs in their lives. Current cigarette user was defined as someone who had smoked cigarettes anytime during the past 30 days, that is, had given any answer other than 0 days to the question: "In the past 30 days, how many days did you smoke cigarettes?"

Participants were considered to have been exposed to second-hand smoke inside the home if they gave any answer other than 0 days to the question: "In the past 7 days, how many days have people smoked in your home, in your presence?" Similarly, they were considered exposed outside the home if they gave any answer other than 0 days to the question and "In the past 7 days, how many days have people smoked in your presence in places other than in your home?"

Consent

Oral consent of parents of the students is usually taken the day before the survey.

Data analysis

Anonymized data were available at the official US CDC site (https://nccd.cdc.gov/GTSS/rdPage.aspx?rdReport=OSH_GTSS.loreByLocation&rdRequestForwarding=Form). We analysed the data using R version 4.2.0 and R studio version 2022.07.01 software. In each survey, adjusted and weighting factors were applied to each student record to adjust for the probability of selection and non-response (by school, class and student).

The weighting factor was: $W = W_1 \times W_2 \times F_1 \times F_2 \times F_3 \times F_4$, where: W_1 = the reverse of probability of selection of the school; W_2 = the reverse of probability of selection of the class within the school; F_1 = adjustment factor of non-response of schools according to size (large, medium, small); F_2 = adjustment factor of class calculated by school; F_3 = adjustment factor of student non-response calculated within this class; and F_4 = adjustment factor post-stratification calculated by sex and grade.

The weighting factor was applied through the survey package of R Studio. Unweighted numbers of students were documented in tables. Indicators were described using weighted percentages reflecting the population estimates. We calculated the 95% confidence intervals (CI) for each proportion. The association between 2 qualitative variables was assessed with the chi-squared test. Trends were assessed using the Cochran Armitage trend test. A two-sided 5% significance level was used for all calculations.

Results

From 2001 to 2017, the number of schools included in the survey increased from 50 to 67. The overall response rate varied from 94.1% (2942/3127) in 2001 to 92.9% (1863/2005) in 2017 (Table 1).

Conventional cigarettes

The male to female ratio was about the same in the 4 years: 0.97 in 2001 and 0.93 in 2017. In 2001, about 23.0% (95% CI: 21.5–24.5%) of the respondents had tried to smoke a cigarette, even if only one or two puffs: 35.4% (95% CI: 32.9–37.9%) of boys and 11.4% (95% CI: 9.9–13.1%) of girls. This proportion increased to 25.0% (95% CI: 23.1–27.1%) in 2017, with the increase greater among boys: 38.8% (95% CI: 35.6–42.0%) of boys and 11.6% (95% CI: 9.6–13.8%) of girls. However, these increases were not significant ($P > 0.05$).

As for current cigarette use, the prevalence decreased significantly from 11.1% (95% CI: 10.0–12.3%) in 2001 to 7.7% (95% CI: 6.5–9.0%) in 2017 ($P < 0.001$). Among boys over the same period, the prevalence of smoking decreased from 19.1% (95% CI: 17.1–21.2%) to 14.2% (95% CI: 12.1–16.7%; $P < 0.001$). Among girls, the prevalence decreased from 3.6% (95% CI: 2.8–4.7%) to 1.4% (95% CI: 0.8–2.4%; $P < 0.001$) (Table 2).

Exposure to second-hand smoke

Between 2001 and 2017, exposure to second-hand smoke at home in the 7 days before the survey decreased significantly from 62.5% (95% CI: 60.7–64.2%) to 46.7% (95% CI: 44.5–49.0%; $P < 0.001$). This reduction was significant for both boys and girls ($P < 0.001$) (Table 3).

Exposure to second-hand smoke outside the home increased significantly between 2001 and 2017, from 65.4% (95% CI: 63.7–67.1%) to 73.3% (95% CI: 71.2–75.3%; $P < 0.001$). This increase was significant for both boys and girls ($P < 0.001$) (Table 3).

Most respondents were in favour of implementing smoke-free public places by law, although this support decreased significantly from 87.0% (95% CI: 85.7–88.2%) in 2001 to 81.8% (95% CI: 80.0–83.5%) in 2017 ($P < 0.001$), and decreased for both boys and girls ($P < 0.001$) (Table 3).

Exposure to tobacco advertising

Exposure to anti-tobacco messages in the media decreased from 87.8% (95% CI: 86.3–89.1%) in 2001 to 64.4% (95% CI: 62.2–66.5%) in 2017 ($P < 0.001$). This decrease was significant for both boys and girls ($P < 0.001$) (Table 4). However, exposure to anti-tobacco messages at sports and cultural events increased significantly, from 34.2% (95% CI: 32.5–35.9%) in 2001 to 72.2% (95% CI: 70.1–74.2%) in 2017 ($P < 0.001$). This increase was significant for

both boys and girls ($P < 0.001$) (Table 4). Two-thirds of students (67.3%; 95% CI: 64.7–69.8%) had seen advertising for tobacco use in 2010. This proportion decreased significantly to 43.7% (95% CI: 41.2–46.2%) in 2017 ($P < 0.001$). This decrease was significant significantly for both boys and girls ($P < 0.001$) (Table 4). The proportion of respondents who had received free promotional cigarettes was small and did not change significantly over the years (Table 4).

Discussion

The GYTS is one of the most important tobacco monitoring tools and helps countries implement the MPOWER package. The questions are in line with the MPOWER package and focus on important aspects of tobacco use and tobacco control. Monitoring the prevalence of tobacco use over time is essential to identify changes and link the national tobacco control strategy to the current situation.

Conventional cigarettes

One in 4 students had ever tried to smoke a cigarette: 1 boy out of 3 and 1 girl out of 10. In the United States, data from the National Youth Tobacco Survey from 2014 to 2016 showed that 21% of adolescents had ever tried to smoke a cigarette (14). The GYTS in the United Arab of Emirates in 2013 focused on expatriate adolescents only and reported that 32% of boys had tried to smoke a cigarette, at a mean age of 12–13 years (15). A previous Tunisian national survey, which included 4172 adolescents aged 12–20 years from public and private schools, reported that among students aged 12–14 years, 26.9% had tried to smoke a cigarette (16). In the Sfax Region in the south of Tunisia, ever cigarette smoking was reported in 16.7% of school students (32.6% of boys and 5.9% of girls) (17). Our findings are similar to these studies and indicate a high prevalence of cigarette experimentation among boys and girls in Tunisia.

Our findings show that the prevalence of current cigarette use among adolescent Tunisians has decreased over time, overall and for boys and girls. According to the last Youth Risk Behavior Survey in the United States conducted in 2019, a significant decrease in current cigarette use had occurred among students in the ninth grade (14–15 years), from 13.5% in 2009 (18) to 3.8% in 2019 (19). In a 45-country analysis of GYTS data in 2013 and

Table 1 Schools, classes and students included in the Global Youth Tobacco Survey, by year, Tunisia

Samples	Year			
	2001	2007	2010	2017
Schools, no.	NA	50	50	67
Classes, no.	NA	76	71	100
All students who participated, no.	4282	2155	1747	2448
Students aged 13–15 years who participated, no.	2942	1499	1294	1863
Response rate, %	94.1 (2942/3127)	92.4 (1499/1622)	94.0 (1294/1377)	92.9 (1863/2005)

NA: data not available.

Table 2 Ever and current cigarette smoking, by sex and year, Tunisia Global Youth Tobacco Survey

Cigarette smoker	% (95% confidence intervals)				P
	2001 (n = 2942)	2007 (n = 1499)	2010 (n = 1294)	2017 (n = 1863)	
Ever smoked					
Overall	23.0 (21.5–24.5)	24.6 (22.5–26.8)	18.6 (16.5–20.8)	25.0 (23.1–27.1)	0.39
Males	35.4 (32.9–37.9)	39.8 (36.2–41.4)	30.6 (27.0–34.4)	38.8 (35.6–42.0)	0.27
Females	11.4 (9.9–13.1)	9.5 (7.6–11.9)	8.0 (6.2–10.4)	11.6 (9.6–13.8)	0.78
Current smoker					
Overall	11.1 (10.0–12.3)	8.3 (7.0–9.8)	6.6 (5.3–8.1)	7.7 (6.5–9.0)	< 0.001
Males	19.1 (17.1–21.2)	15.1 (12.6–17.9)	12.4 (9.9–15.3)	14.2 (12.1–16.7)	< 0.001
Females	3.6 (2.8–4.7)	1.6 (0.7–2.9)	1.6 (0.9–2.9)	1.4 (0.8–2.4)	< 0.001

Table 3 Exposure to second-hand smoke and perceptions about smoke-free public places mandate, by year, Tunisia Global Youth Tobacco Survey

Exposure to second-hand smoke and perception	% (95% confidence intervals)				P
	2001 (n = 2942)	2007 (n = 1499)	2010 (n = 1294)	2017 (n = 1863)	
At home					
Overall	62.5 (60.7–64.2)	52.2 (49.7–54.7)	50.6 (47.9–53.3)	46.7 (44.5–49.0)	< 0.001
Males	62.3 (59.8–64.8)	53.3 (49.7–56.8)	49.3 (45.4–53.3)	46.3 (43.1–49.5)	< 0.001
Females	62.6 (60.2–65.0)	51.0 (47.4–54.6)	54.4 (47.7–55.2)	47.1 (43.9–50.3)	< 0.001
Outside the home					
Overall	65.4 (63.7–67.1)	65.6 (63.2–68)	61.6 (59–64)	73.3 (71.2–75.3)	< 0.001
Males	69.7 (67.2–72)	70.0 (66.6–73.2)	64.3 (60.4–68)	75.3 (72.4–78)	< 0.001
Females	61.4 (58.9–63.8)	61.4 (57.8–64.8)	59.2 (55.4–62.8)	71.3 (68.3–74.1)	< 0.001
In favour of smoke-free places					
Overall	87.0 (85.7–88.2)	85.3 (83.4–87)	78.3 (76–80.3)	81.8 (80–83.5)	< 0.001
Males	86.0 (84–87.7)	83.9 (81.1–86.4)	77.4 (73.8–80.5)	82.0 (79.3–84.3)	< 0.001
Females	88.1 (86.3–89.6)	86.8 (84.2–89.1)	79.3 (76–82.2)	81.6 (79–84)	< 0.001

2014, the median global prevalence of current cigarette use across all countries was 6.8% (9.7% in boys and 3.5% in girls), which is lower than the prevalence in our 4 surveys overall and for boys, but higher than current cigarette use we found for girls. Given the findings of the 2017 GYTS in Tunisia, the country has the fourth highest prevalence of adolescent cigarette use in the Middle East and North African Region, after Jordan (2014 GYTS), Lebanon (2013 GYTS) and Qatar (2013 GYTS) (20). Other studies in North African countries showed that a greater proportion of Tunisian boys smoked than Egyptian, Libyan, Moroccan and Sudanese boys (21). In Malaysia, the prevalence of current cigarette use decreased from 19.9% in 2003 to 14.8% in 2016, which is almost double the prevalence in our study (22). In Morocco, the current cigarette use among 13–15-year-old schoolchildren increased from 3.0% in 2006 to 5.2% in 2010, but both were lower than the prevalence among Tunisian schoolchildren (23). In the city of Sousse in Tunisia, the results of a cross-sectional survey in 2013–2014 in 16 public schools showed that 4.5% of participants were cigarette users, which is lower than the national prevalence in the GYTS surveys (24). Even though cigarette consumption among Tunisian

schoolchildren decreased between 2001 and 2017 it is however still high and needs to be tackled to reduce the prevalence further.

Exposure to second-hand smoke

Second-hand smoke outside the home among adolescents increased between 2001 and 2017. A study in 131 countries found that exposure to second-hand smoke outside the home was 57.6% in 2018 and it had not decreased from 1999 – it remained the same in 46 of 131 countries (35.1%) and increased in 40 (30.5%). This increase was found in almost all WHO regions (exposure was 59.4% for exposure at least one day a week in the Middle East and North Africa Region) and in countries that did not ratify the FCTC (25). The overall exposure to second-hand smoke in public places among non-smoking adolescents was 44.2% across 168 countries between 2001 and 2017. The exposure was higher among boys than girls. Exposure ranged from 39.8% in the Middle East and North Africa Region to 73.7% in the European Region (26). In Africa, from 2006 to 2011, exposure to second-hand smoke among adolescents was 39.0%; it ranged from 24.9% in Cape Verde to 80.4% in Mali, with no differences between the sexes (27). These

Table 4 Prevalence of exposure to tobacco advertising, by year, Tunisia, Global Youth Tobacco Survey

Exposed to:	% (95% confidence intervals)				P
	2001 (n = 2942)	2007 (n = 1499)	2010 (n = 1294)	2017 (n = 1863)	
Anti-tobacco messages in media					
Overall	87.8 (86.3–89.1)	71.8 (69.4–74.0)	85.4 (83.4–87.2)	64.4 (62.2–66.5)	< 0.001
Male	87.3 (85.2–89.1)	71.1 (67.7–74.2)	84.0 (80.9–86.7)	59.4 (56.1–62.5)	< 0.001
Female	88.2 (86.1–90.0)	72.6 (69.2–75.7)	86.6 (83.8–88.9)	69.1 (66.1–72.0)	< 0.001
Anti-tobacco messages at sports and cultural events					
Overall	34.2 (32.5–35.9)	26.4 (24.2–28.7)	19.3 (17.2–21.5)	72.2 (70.1–74.2)	< 0.001
Male	29.9 (27.6–32.4)	22.8 (19.9–25.9)	19.3 (16.4–22.7)	67.1 (64–70.1)	< 0.001
Female	38.2 (35.7–40.6)	29.4 (26.3–32.8)	19.0 (16.2–22.2)	77.3 (74.5–79.9)	< 0.001
Publicity at points of sale					
Overall	NA	NA	67.3 (64.7–69.8)	43.7 (41.2–46.2)	< 0.001
Male	NA	NA	67.1 (63.3–70.8)	47.4 (43.9–51.0)	< 0.001
Female	NA	NA	67.3 (63.6–70.7)	39.8 (36.3–43.4)	< 0.001
Promotional offers of free cigarettes					
Overall	5.5 (4.7–6.4)	4.8 (3.8–6.0)	4.3 (3.3–5.5)	5.6 (4.6–6.7)	0.89
Male	8.1 (6.8–9.7)	5.7 (4.2–7.6)	5.6 (4.0–7.7)	8.0 (6.4–9.9)	0.70
Female	3.1 (2.3–4.1)	3.9 (2.7–5.6)	3.0 (2.0–4.6)	3.3 (2.3–4.6)	0.82

NA: data not available as this question was not included in the questionnaire at that time.

results show that exposure to second-hand smoke in Tunisia is among the highest in the world.

The proportion of students in favour of laws that establish smoke-free public places decreased for both sexes. Tunisia established its first tobacco law (no. 98-17) in February 1998 which aimed to protect people from tobacco harm. Article 10 of this law prohibits smoking in public places (28). This law was enforced by the decree of November 1998 (29), decree of September 2009 (30) and ratification of the FCTC in 2010. Article 8 of the FCTC calls on countries to adopt and implement effective national legislations to protect people from exposure to tobacco in indoor and outdoor public places (8). However, the compliance of Tunisians and respect for these laws seem to be weak given the high rates of exposure to second-hand smoke outside the home (31). In the most recent report of MPOWER in the Middle East and North Africa Region, Tunisia had a score of 1 out of 3 for smoke-free places, which means only up to 2 public places were completely smoke free (32). A longitudinal study found evidence that, in addition to positive impact on exposure to second-hand smoke, laws on smoke-free places led to a possible decrease in smoking prevalence (33).

Exposure to tobacco advertising

In the 2001 Tunisian GYTS, a greater proportion of respondents were exposed to anti-tobacco messages in the media (internet, magazines, television) than respondents in the 2017 GYTS. This result is similar to findings in Greece (34), Italy (35) and Myanmar (36). A longitudinal study in the United States found a positive effect of anti-tobacco messages on teenagers' susceptibility to tobacco smoke. In fact, this exposure decreased the susceptibility

to tobacco smoke by 2 or 3 years (37). From 2010 to 2017, the proportions of students exposed to cigarette advertising at points of sale decreased. A systematic review in 2009 concluded that exposure to promotion of cigarette use at points of sale increased the odds of ever smoking, frequent smoking or occasional smoking (38). This explains why the tobacco industry spends around 80% of their advertising budget on promotions at points of sale (39). Article 13 of the FCTC calls on countries to ban every kind of tobacco promotions, advertisings and sponsorships. In Tunisia, law no. 98-17 forbids all types of promotion of tobacco products in public places, but it does not include a ban on promotion at points of sale (28).

Strengths and limitations

A strength of our study is that the GYTS is the only standardized worldwide survey on tobacco use and attitudes among adolescents aged 13–15 years. The GYTS is a national survey conducted in all governorates and cities in the country. The sample size of students who answered the questionnaire was large and the response rates were always more than 92%.

Our study has some limitations. Smoking behaviour and exposure to second-hand smoke were self-reported and no quantitative method was used to confirm the students' responses, which may introduce biases. Only students in public schools were included, thus students in private schools or adolescents who were not in school were not represented. Students in private schools and out-of-school teenagers represent about 10% of Tunisian adolescents (40).

Conclusions

WHO recommends that countries implement a monitoring survey every 5 years. It has been 5 years since the last GYTS in Tunisia and a new GYTS survey is needed. Efforts to ensure complete compliance with

smoke-free public places law are needed to decrease the prevalence of exposure to second-hand smoke. A complete ban on point of sales promotions is strongly recommended to decrease the exposure of vulnerable young people to tobacco advertising.

Acknowledgement

This study was a collaborative project of WHO, US CDC and the Ministry of Health of Tunisia. We thank the study participants and research assistants.

Funding: None.

Competing interests: None declared.

Consommation de cigarettes et exposition à la fumée secondaire et à la publicité en faveur du tabac chez les adolescents tunisiens

Résumé

Contexte : L'enquête mondiale sur le tabagisme chez les jeunes a été menée en Tunisie en 2001, 2007, 2010 et 2017. Il s'agit de la seule enquête nationale portant sur l'exposition à la fumée secondaire et à la publicité en faveur du tabac chez les jeunes. Aucune description de la tendance n'a été réalisée.

Objectif : Décrire les tendances de la consommation de cigarettes chez les adolescents tunisiens et leur exposition à la fumée secondaire et à la publicité en faveur du tabac, entre 2001 et 2017.

Méthodes : L'enquête mondiale sur le tabagisme chez les jeunes est une enquête transversale en milieu scolaire menée par l'Organisation mondiale de la Santé. L'enquête repose sur un échantillonnage en grappes à deux degrés afin d'obtenir un échantillon représentatif d'élèves âgés de 13 à 15 ans. Un questionnaire standardisé est utilisé pour la collecte de données. Nous avons comparé, sur la période des quatre années (2001, 2007, 2010 et 2017), la prévalence et les intervalles de confiance (IC) à 95 % de la consommation de cigarettes antérieure et au moment des enquêtes, l'exposition à la fumée secondaire à l'intérieur et à l'extérieur du domicile, ainsi que l'exposition à la publicité en faveur du tabac.

Résultats : La consommation de cigarettes au fil des enquêtes a diminué, passant de 11,1 % (IC à 95 % : 10,0-12,3 %) en 2001 à 7,7 % (IC 95 % : 6,5-9,0 %) en 2017 ($p < 0,001$). L'exposition à la fumée secondaire à domicile a diminué, passant de 62,5 % (IC à 95 % : 60,7-64,2 %) à 46,7 % (IC à 95 % : 44,5-49,0 %) au cours de la même période ($p < 0,001$), mais l'exposition en dehors du domicile a augmenté, passant de 65,4 % (IC à 95 % : 63,7-67,1 %) en 2001 à 73,3 % (IC 95 % : 71,2-75,3 %) en 2017 ($p < 0,001$). L'exposition aux messages antitabac dans les médias a diminué, passant de 87,8 % (IC à 95 % : 86,3-89,1 %) en 2001 à 64,4 % (IC 95 % : 62,2-66,5 %) en 2017 ($p < 0,001$).

Conclusion : Si la prévalence de la consommation de cigarettes et de l'exposition à la fumée secondaire à domicile a diminué, l'exposition en dehors du domicile a augmenté. Des efforts sont nécessaires pour garantir le respect des lois antitabac afin de réduire la prévalence de l'exposition à la fumée secondaire.

تدخين السجائر والتعرض للتدخين غير المباشر والترويج للتدخين بين صفوف المراهقين التونسيين

يسر العيادي، شهيدة الحريزي، عفاف السخيري، رضوان الفخفاخ

الخلاصة

الخلفية: أجري "المسح العالمي للتبغ بين الشباب" في تونس في الأعوام 2001 و 2007 و 2010 و 2017، وهو المسح الوطني الوحيد الذي فُحص التعرض للتدخين غير المباشر والترويج للتبغ بين الشباب، ولم يرد فيه وصف للتوجه.

الأهداف: وصف توجهات تدخين السجائر بين صفوف المراهقين التونسيين وتعرضهم للتدخين غير المباشر والترويج للتبغ بينهم في الفترة من 2001 إلى 2017.

طرق البحث: المسح العالمي للتبغ بين الشباب هو مسح مقطعي أجرته منظمة الصحة العالمية في المدارس. ويستخدم المسح تصميمًا لأخذ العينات عنقوديًا على مرحلتين، بغرض الحصول على عينة ممثلة للطلاب الذين تتراوح أعمارهم بين 13 و 15 عامًا. واستُخدم استبيان موحد لأغراض جمع البيانات. وقارنًا معدل الانتشار وفواصل ثقة قدرها 95٪ بين تدخين السجائر في المطلق والوقت الحالي، والتعرض للتدخين غير المباشر داخل المنزل وخارجه، والتعرض للترويج للتبغ، على مدى 4 سنوات.

النتائج: انخفض التدخين الحالي للسجائر من 11.1٪ في 2001 (فاصل الثقة 95٪: 10.0–12.3٪) إلى 7.7٪ في عام 2017 (فاصل الثقة 95٪: 6.5–9.0٪) (القيمة الاحتمالية > 0.001). وانخفض التعرض للتدخين غير المباشر في المنزل من 62.5٪ (فاصل الثقة 95٪: 60.7–64.2٪) إلى 46.7٪ (فاصل الثقة 95٪: 44.5–49.0٪) خلال الفترة نفسها (القيمة الاحتمالية > 0.001)، ولكن التعرض خارج المنزل زاد من 65.4٪ في 2001 (فاصل ثقة 95٪: 63.7–67.1٪) إلى 73.3٪ في عام 2017 (فاصل الثقة 95٪: 71.2–75.3٪) (القيمة الاحتمالية > 0.001). وانخفض التعرض للرسائل المناهضة للتبغ في وسائل الإعلام من 87.8٪ في 2001 (فاصل الثقة 95٪: 86.3–89.1٪) إلى 64.4٪ في 2017 (فاصل الثقة 95٪: 66.5–62.2٪) (القيمة الاحتمالية > 0.001).

الاستنتاجات: في حين انخفض معدل انتشار تدخين السجائر والتعرض لدخان التبغ غير المباشر في المنزل، زاد التعرض خارج المنزل. ويلزم بذل الجهود لضمان الامتثال لقوانين حظر التدخين بغرض خفض معدل انتشار التعرض لدخان التبغ غير المباشر.

References

1. Abbafati C, Abbas KM, Abbasi-Kangevari M, Abd-Allah F, Abdelalim A, Abdollahi M, et al. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396(10258):1223–49. [https://doi.org/10.1016/S0140-6736\(20\)30752-2](https://doi.org/10.1016/S0140-6736(20)30752-2)
2. Reitsma MB, Kendrick PJ, Ababneh E, Abbafati C, Abbasi-Kangevari M, Abdoli A, et al. Spatial, temporal, and demographic patterns in prevalence of smoking tobacco use and attributable disease burden in 204 countries and territories, 1990–2019: a systematic analysis from the Global Burden of Disease Study 2019. *Lancet*. 2021;397(10292):2337–60. [https://doi.org/10.1016/S0140-6736\(21\)01169-7](https://doi.org/10.1016/S0140-6736(21)01169-7)
3. The epidemiology of tobacco use among young people in the United States and worldwide. In: Preventing tobacco use among youth and young adults: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2012.
4. Cantrell J, Bennett M, Mowery P, Xiao H, Rath J, Hair E, et al. Patterns in first and daily cigarette initiation among youth and young adults from 2002 to 2015. *PLoS One*. 2018;13(8):1–20. <https://doi.org/10.1371/journal.pone.0200827>
5. Hughes J, Kabir Z, Bennett K, Hotchkiss JW, Kee F, Leyland AH, et al. Modelling future coronary heart disease mortality to 2030 in the British Isles. *PLoS One*. 2015;10(9):1–12. <https://doi.org/10.1371/journal.pone.0138044>
6. Öberg M, Jaakkola MS, Woodward A, Peruga A, Prüss-Ustün A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet*. 2011;377(9760):139–46. [https://doi.org/10.1016/S0140-6736\(10\)61388-8](https://doi.org/10.1016/S0140-6736(10)61388-8)
7. Blake KD, Viswanath K, Blendon RJ, Vallone D. The role of tobacco-specific media exposure, knowledge, and smoking status on selected attitudes toward tobacco control. *Nicotine Tob Res*. 2010;12(2):117–26. <https://doi.org/10.1093/ntr/ntp184>
8. WHO Framework Convention on Tobacco Control [Internet]. Geneva: World Health Organization; 2005 (<https://fctc.who.int/>, accessed 20 January 2022).
9. Tunisian Health Examination Survey – 2016. Tunis: République Tunisienne, Ministère de la Santé, Institut National de la Santé; 2019 (<http://www.santetunisie.rns.tn/images/rapport-final-enquete2020.pdf>, accessed 20 January 2022).
10. Harizi C, El-Awa F, Ghedira H, Audera-Lopez C, Fakhfakh R. Implementation of the WHO Framework Convention on Tobacco Control in Tunisia: Progress and challenges. *Tob Prev Cessat*. 2020;6:1–8. <https://doi.org/10.18332/tpc/130476>
11. Global Youth Tobacco Survey. Tunisia [Internet]. Geneva: World Health Organization (https://extranet.who.int/ncdsmicrodata/index.php/catalog/GYTS#_r=&collection=&country=217&dtype=&from=1999&page=1&ps=&sid=&sk=&sort_by=nation&sort_order=&to=2019&topic=&view=s&vk=, accessed 9 September 2020).
12. Global youth tobacco survey [Internet]. Geneva: World Health Organization (<https://www.who.int/tobacco/surveillance/gyts/en/>, accessed 31 August 2020).
13. Global Youth Tobacco Survey Collaborative Group. Global youth tobacco survey (GYTS): core questionnaire with optional questions. Version 1.2. Atlanta, GA: Centers for Disease Control and Prevention; 2014.
14. Sharapova S, Reyes-Guzman C, Singh T, Phillips E, Marynak KL, Agaku I. Age of tobacco use initiation and association with current use and nicotine dependence among US middle and high school students, 2014–2016. *Tob Control*. 2020;29(1):49–54. <https://doi.org/10.1136/tobaccocontrol-2018-054593>
15. Asfour LW, Stanley ZD, Weitzman M, Sherman SE. Uncovering risky behaviors of expatriate teenagers in the United Arab Emirates: a survey of tobacco use, nutrition and physical activity habits. *BMC Public Health*. 2015;15(1):944. <https://doi.org/10.1186/s12889-015-2261-9>
16. Fakhfakh R, Jaidane I, Hsairi M, Ben Hamida AM. Les facteurs de risque et de protection de l'initiation à la cigarette chez les adolescents tunisiens [Cigarette smoking initiation among Tunisian adolescents: Risk and protective factors]. *Rev Epidemiol Sante Publique*. 2015;63(6):369–79. <http://dx.doi.org/10.1016/j.respe.2015.09.005>
17. Ben Ayed H, Yaich S, Ben Hmida M, Ben Jemaa M, Trigui M, Karray R, et al. Prevalence and factors associated with smoking among Tunisian secondary school-adolescents. *Int J Adolesc Med Health*. 2021;33(6):379–87. <https://doi.org/10.1515/ijamh-2019-0088>

18. Centers for Disease Control and Prevention. Youth risk behavior surveillance—United States, 2009. Surveillance summaries, June 2010. *MMWR Surveill Summ.* 2010;59(5):1–142.
19. Creamer MLR, Everett Jones S, Gentzke AS, Jamal A, King BA. Tobacco product use among high school students – youth risk behavior survey, United States, 2019. *MMWR Suppl.* 2020;69(1):56–63. <https://doi.org/10.15585/mmwr.su6901a7>
20. D'Angelo D, Ahluwalia IB, Pun E, Yin S, Palipudi K, Mbulo L. Current cigarette smoking, access, and purchases from retail outlets among students aged 13–15 years—Global Youth Tobacco Survey, 45 countries, 2013 and 2014. *MMWR Morb Mortal Wkly Rep.* 2016;65(34):898–901. <https://doi.org/10.15585/mmwr.mm6534a3>
21. Madkour AS, Ledford EC, Andersen L, Johnson CC. Tobacco advertising/promotions and adolescents' smoking risk in Northern Africa. *Tob Control.* 2014;23(3):244–52. <https://doi.org/10.1136/tobaccocontrol-2012-050593>
22. Lim K, Ghazali S, Lim H, Kee C, Cheah Y, Pradhaman Singh B, et al. Tobacco use and other aspects related to smoking among school-going adolescents aged 13–15 years in Malaysia: analysis of three cross-sectional nationally representative surveys in 2003, 2009 and 2016. *Tob Induc Dis.* 2020;18(September):1–10. <https://doi.org/10.18332/tid/127231>
23. Shaikh MA. Prevalence, correlates, and changes in tobacco use between 2006 and 2010 among 13–15 year Moroccan school attending adolescents. *J Pak Med Assoc.* 2014;64(11):1306–9.
24. Nawel Z, Jihen M, Rim G, Sana B, Hassen G. Tobacco use: the main predictor of illicit substances use among young adolescents in Sousse, Tunisia. *Int J Adolesc Med Health.* 2018;32(5). <https://doi.org/10.1515/ijamh-2017-0213>
25. Ma C, Heiland EG, Li Z, Zhao M, Liang Y, Xi B. Global trends in the prevalence of secondhand smoke exposure among adolescents aged 12–16 years from 1999 to 2018: an analysis of repeated cross-sectional surveys. *Lancet Glob Health.* 2021;9(12):e1667–78. [https://doi.org/10.1016/S2214-109X\(21\)00365-X](https://doi.org/10.1016/S2214-109X(21)00365-X)
26. Ma C, Heiland EG, Li Z, Zhao M, Liang Y, et al. Secondhand smoke exposure among never-smoking youth in 168 countries. *J Adolesc Health.* 2015;56(2):167–73. [https://doi.org/10.1016/S2214-109X\(21\)00365-X](https://doi.org/10.1016/S2214-109X(21)00365-X)
27. Owusu D, Mamudu HM, John RM, Ibrahim A, Ouma AEO, Veeranki SP. Never-smoking adolescents' exposure to secondhand smoke in Africa. *Am J Prev Med.* 2016;51(6):983–98. <https://doi.org/10.1016/j.amepre.2016.08.040>
28. [Law no. 98-17 of February 23 1998, relative to prevention of the harmful effects of smoking]. *Journal Officiel de la République Tunisienne*; 1998 27 février:399–400.
29. Decree no. 98–2248, November, 16 1998. Fixant les lieux affectés à l'usage collectif dans lesquels il est interdit de fumer [Identifying smoke-free public places]. *Tunis: Government of Tunisia*; 1998.
30. Décret no. 2009-2611 du 14 septembre 2009, complétant le décret n° 98-2248 du 16 novembre 1998 fixant les lieux affectés à l'usage collectif dans lesquels il est interdit de fumer [Completing decree no. 98-2248 of November, 16 1998]. *Tunis: Government of Tunisia*; 2009.
31. Ben Amar W, Chakroun A, Zribi M, Khemekhem Z, Ben Jemaa F, Maatoug S. Dispositif législatif de lutte anti-tabagique en Tunisie : entre insuffisances et défaut d'application [Anti tobacco legislation and regulation in Tunisia: between shortcomings and lack of application]. *JIM Sfax.* 2017;17:21–6.
32. Heydari G, Zaatari G, Al-Lawati JA, El-Awa F, Fouad H. MPOWER, needs and challenges: trends in the implementation of the WHO FCTC in the Eastern Mediterranean Region. *East Mediterr Health J.* 2018;24(01):63–71. <https://doi.org/10.26719/2018.24.1.63>
33. Becker CM, Lee JGL, Hudson S, Hoover J, Civils D. A 14-year longitudinal study of the impact of clean indoor air legislation on state smoking prevalence, USA, 1997–2010. *Prev Med (Baltim).* 2017;99:63–6. <https://doi.org/10.1016/j.ypmed.2017.01.016>
34. Kyrlesi A, Soteriades ES, Warren CW, Kremastinou J, Papastergiou P, Jones NR, et al. Tobacco use among students aged 13–15 years in Greece: the GYTS project. *BMC Public Health.* 2007;7(1):3. <https://doi.org/10.1186/1471-2458-7-3>
35. Gorini G, Gallus S, Carreras G, Cortini B, Vannacci V, Charrier L, et al. A long way to go: 20-year trends from multiple surveillance systems show a still huge use of tobacco in minors in Italy. *Eur J Public Health.* 2019;29(1):164–9. <https://doi.org/10.1093/EURPUB/CKY132>
36. Tun N, Chittin T, Agarwal N, New M, Thauang Y, Phyo P. Tobacco use among young adolescents in Myanmar: findings from global youth tobacco survey. *Indian J Public Health.* 2017;61(5):54. https://doi.org/10.4103/ijph.IJPH_236_17
37. Weiss JW, Cen S, Schuster D, Unger J, Johnson CA, Mouttapa M, et al. Longitudinal effects of pro-tobacco and anti-tobacco messages on adolescent smoking susceptibility. *Nicotine Tob Res.* 2006;8(3):455–65. <https://doi.org/10.1080/14622200600670454>
38. Paynter J, Edwards R. The impact of tobacco promotion at the point of sale: a systematic review. *Nicotine Tob Res.* 2009;11(1):25–35. <https://doi.org/10.1093/ntr/ntn002>
39. Ma H, Reimold AE, Ribisl KM. Trends in cigarette marketing expenditures, 1975–2019: an analysis of federal trade commission cigarette reports. *Nicotine Tob Res.* 2022;24(6):919–23. <https://doi.org/10.1093/ntr/ntab272>
40. Middle East and North Africa out-of-school children initiative. Summary, Tunisia: country report on out-of-school children. *Amman: UNICEF MENA Regional Office*; 2015 (https://www.unicef.org/mena/media/6661/file/Tunisia%20Country%20Report%20on%20OOSC%20Summary_EN.pdf%20.pdf, accessed 20 January 2022).