

# Incidence of acute myocardial infarction in Islamic Republic of Iran: a study using national registry data in 2012

A. Ahmadi,<sup>1</sup> H. Soori,<sup>2</sup> Y. Mehrabi,<sup>2</sup> K. Etemad<sup>2</sup>, T. Samavat<sup>3</sup> and A. Khaledifar<sup>4</sup>

معدّل وقوع احتشاء عضل القلب الحاد في جمهورية إيران الإسلامية: دراسة باستخدام بيانات السجل الوطني في عام 2012

علي أحمددي، حامد سوري، يادولا مهراي، كوروش اعتقاد، طاهرة ساوات، أرسلان خالديفار

الخلاصة: لم يتم إعداد تقارير عن بيانات سكانية حول معدلات احتشاء عضل القلب في جمهورية إيران الإسلامية على النطاق الوطني أو على نطاق المحافظات. ولذلك قام الباحثون - في دراسة مستعرضة - بجمع بيانات عن 20 750 حالة جديدة من حالات احتشاء عضل القلب (التصنيف الدولي للأمراض 10 الرواميز 121-22) أدخلت إلى المستشفيات وسُجّلت من قِبَل مكتب التسجيل الإيراني لاحتشاء عضل القلب في عام 2012. وتم حساب معدّل الوقوع الخام والمصحّح حسب السن لكل 100 000 شخص - بشكل مباشر - في كل من المحافظات الـ 31 وفي البلد ككل باستخدام معيار منظمة الصحة العالمية لعدد السكان. وقد شكّل الذكور بوجه عام 72.4% من الحالات، وكان العمر الوسطي لديهم عند الوقوع أقل بكثير من النساء [59.6 (SD 13.3) سنة مقابل 65.4 (SD 12.6) سنة]. وكانت نسبة وقوعات الذكور إلى الإناث: 2.63. وكان معدّل وقوع احتشاء عضل القلب المعيّر حسب السن 73.3 لكل 100 000 في البلد ككل (95% CI: 72.3-74.3%)، وتباين بين المحافظات الإحدى والثلاثين بشكل كبير من 24.5 إلى 152.5 لكل 100 000. وهذه الدراسة تقدم بيانات أولية لرصد وإدارة الأمراض القلبية الوعائية في البلد.

ABSTRACT Population-based data on myocardial infarction rates in the Islamic Republic of Iran have not been reported on a national or provincial scale. In a cross-sectional study, data were collected on 20 750 new cases of myocardial infarction (ICD10 codes I21-22) admitted to hospitals and registered by the Iranian Myocardial Infarction Registry in 2012. The crude and age-adjusted incidence for the 31 provinces and the whole country were directly calculated per 100 000 people using the WHO standard population. Overall, males comprised 72.4% of cases and had a significantly lower mean age at incidence than women [59.6 (SD 13.3) years versus 65.4 (SD 12.6) years]. The male:female incidence ratio was 2.63. The age-standardized myocardial infarction incidence rate was 73.3 per 100 000 in the whole country (95% CI: 72.3%-74.3%) and varied significantly from 24.5 to 152.5 per 100 000 across the 31 provinces. The study provides baseline data for monitoring and managing cardiovascular diseases in the country.

Incidence de l'infarctus aigu du myocarde en République islamique d'Iran : étude à partir des données du registre national en 2012

RÉSUMÉ Les données en population sur les taux d'infarctus du myocarde en République islamique d'Iran n'ont pas été rapportées à l'échelle régionale ou nationale. Dans une étude transversale, des données ont été recueillies sur 20 750 nouveaux cas d'infarctus du myocarde (CIM-10 codes I21-22) admis dans des établissements hospitaliers et enregistrés dans le Registre iranien des infarctus du myocarde en 2012. L'incidence brute et l'incidence ajustée pour l'âge dans les 31 provinces et l'ensemble du pays ont été directement calculées pour 100 000 personnes à l'aide de la population type de l'OMS. Globalement, les hommes représentaient 72,4 % des cas et étaient nettement moins âgés en moyenne à l'incidence que les femmes (59,6 ans [ET 13,3 ans] contre 65,4 ans [ET 12,6 ans]). Le rapport d'incidence homme-femme était de 2,63. Le taux d'incidence standardisé en fonction de l'âge pour l'infarctus du myocarde était de 73,3 pour 100 000 dans l'ensemble du pays (IC à 95 % : 72,3 % - 74,3 %) et variait significativement, de 24,5 à 152,5 pour 100 000, entre les 31 provinces. L'étude fournit des données initiales pour le suivi et la prise en charge des maladies cardio-vasculaires dans le pays.

<sup>1</sup>Department of Epidemiology & Biostatistics, School of Public Health; <sup>2</sup>Safety Promotion & Injury Prevention Research Center, Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Islamic Republic of Iran (Correspondence to H. Soori: hsoori@yahoo.com). <sup>3</sup>Ministry of Health & Medical Education, Center for Non-Communicable Disease Control, Cardiovascular Office, Tehran, Islamic Republic of Iran. <sup>4</sup>Department of Cardiology, School of Medicine & Hajar Hospital, Shahrekord University of Medical Sciences, Shahrekord, Islamic Republic of Iran.

Received: 22/03/14; accepted: 08/09/14

## Introduction

Despite decreases in mortality rates due to cardiovascular diseases in most developed countries, these diseases have become the most important health problem and cause of mortality in many developing countries worldwide, including the Islamic Republic of Iran (1,2). Although cardiovascular diseases were previously of greatest concern to high-income countries (3), the age-adjusted incidence of these diseases has recently declined in these countries. Coronary heart disease is also on the rise in low- and moderate income countries (4) due to increases in life expectancy and changes in urbanization and lifestyle that have increased the prevalence of cardiovascular disease risk factors (5–7).

Cardiovascular diseases are generally manifested by myocardial infarction, and the World Health Organization (WHO) has suggested that myocardial infarction rates can be used as a proxy for cardiovascular disease rates in epidemiological studies (8). Comparison of rates of myocardial infarction in different geographical areas and communities within a country reveal important information that can be applied in evidence-based decision-making, research, prioritization and health systems planning in order to better identify myocardial infarction etiology and risk factors and to assess cardiovascular disease prevention approaches (9–11).

Reliable data to determine myocardial infarction rates are rare in developing countries. In the Islamic Republic of Iran, for example, the data required to assess myocardial infarction incidence have been mainly based on a limited number of hospitals and limited studies on small populations (12–14). According to a report by the Iranian Ministry of Health and Medical Education in 2012, cardiovascular diseases overall comprised 39% of the causes of death, and the mortality rate

due to myocardial infarction was 85 per 100 000 (6). To date, no adequate estimate of myocardial infarction incidence has been reported for the whole country or in all the regions. In view of this, and reports of the varying distribution of cardiovascular disease risk factors across the different provinces (15,16), the present study was conducted to determine the incidence rate of myocardial infarction at the national and provincial level in the Islamic Republic of Iran.

## Methods

### Study setting

The Iranian Myocardial Infarction Registry was set up in 2009 by the Office of Cardiovascular Diseases of the Iranian Ministry of Health and Medical Education. At the end of 2011, the coverage of the Registry was complete and was usable across the whole country (6,17). The Registry covers all hospitals that are equipped with a coronary care unit in 31 provinces of the country.

### Data collection

In a cross-sectional study, data were collected on 20 750 new cases of myocardial infarction recorded by the Registry in 2012. Inclusion criteria were based on the WHO and World Heart Federation definition of myocardial infarction diagnosis as per the *International Classification of Diseases (ICD10)* codes I21 and I22 (8). Patients with myocardial infarction history or no definite diagnosis by the cardiologist were excluded from the study. The data collected included patients' age, sex and province of residence. The ethics committee of Shahid Beheshti University of Medical Sciences approved this study.

### Data analysis

Crude incidence rates were calculated for each province and the whole country per 100 000 people for different age

groups. To estimate the population denominator for the incidence, the data of the population census from the Statistical Center of Iran in 2011 were used. To adjust the rates, we used the direct standardization method and the WHO standard population (18). All continuous values were reported as means and standard deviation (SD) and categorical variables as percentages. The *t*-test and analysis of variance were employed to compare differences in means of continuous variables between 2 groups and among more than 2 groups. The data were analysed by *Stata* software, version 12. *P*-values < 0.05 were considered as significant.

## Results

Of 20 750 new cases of myocardial infarction, 15 033 (72.4%) were in males and 5717 (27.6%) in females. The male:female incidence ratio was 2.63.

### Age at myocardial infarction

The mean age at myocardial infarction was 61.2 (SD 13.4) years in the whole population. The mean age in men [59.6 (SD 13.3) years] was significantly lower compared with women [65.4 (SD 12.6) years] ( $P = 0.001$ ).

The mean age in different provinces by sex is shown in Table 1. The mean age at myocardial infarction varied significantly across different provinces ( $P = 0.001$ ). The lowest mean age was recorded in patients residing in Semnan province [59.1 (SD 21.9) years], followed by Tehran [60.4 (SD 13.5) years] and Lorestan [60.1 (SD 14.5) years], and these mean ages were significantly different from those in Ardabil [63.2 (SD 15.1) years] ( $P = 0.042$ ). The highest mean age was in Zanjan province [64.4 (SD 12.8) years] and this was higher than in Sistan and Baluchistan [60.3 (SD 13.9) years] ( $P = 0.019$ ). The mean age of men was not significantly different across the provinces. For women, the

**Table 1 Mean age at incidence of myocardial infarction by sex in provinces of the Islamic Republic of Iran, 2012**

Province	Mean (SD) age (years)			P-value
	Total	Males	Females	
Alborz	60.4 (13.3)	58.6 (13.1)	65.6 (12.5)	0.001
Ardabil	63.2 (15.1)	60.8 (14.6)	68.4 (15.0)	0.001
West Azerbaijan	61.5 (13.6)	60.4 (13.5)	64.3 (13.3)	0.001
East Azerbaijan	61.0 (12.7)	59.0 (12.6)	64.9 (11.8)	0.001
Bushehr	61.3 (13.4)	60.9 (12.9)	62.5 (14.6)	0.462
Chaharmahal & Bakhtiari	62.4 (13.3)	60.2 (13.3)	68.9 (11.0)	0.001
Ilam	62.4 (11.9)	59.2 (11.8)	67.4 (10.3)	0.001
Esfahan	61.7 (13.6)	59.7 (13.5)	66.9 (12.7)	0.001
Fars	61.0 (13.6)	59.4 (13.7)	65.3 (12.3)	0.001
Ghazvin	61.4 (12.8)	59.6 (12.6)	66.4 (11.9)	0.001
Gilan	60.6 (13.2)	58.9 (13.3)	65.7 (11.7)	0.001
Golestan	60.4 (12.5)	58.7 (12.2)	64.4 (12.1)	0.001
Hamadan	60.0 (13.8)	59.3 (14.0)	62.1 (12.9)	0.068
Hormozgan	62.1 (13.8)	60.0 (14.0)	67.9 (11.4)	0.001
Kerman	61.4 (12.9)	59.5 (12.7)	66.5 (12.0)	0.001
Kermnanshah	62.0 (13.7)	60.1 (13.3)	66.2 (13.8)	0.001
South Khorasan	61.6 (13.5)	60.1 (13.9)	65.3 (11.8)	0.009
Razavi Khorasan	61.8 (12.9)	60.6 (13.0)	65.3 (12.1)	0.001
North Khorasan	61.7 (13.0)	59.6 (13.1)	66.3 (11.5)	0.001
Khuzestan	60.8 (13.2)	59.4 (13.3)	64.6 (12.1)	0.001
Kohgiluyeh & Boyer-Ahmad	61.0 (11.8)	58.5 (11.1)	66.8 (11.5)	0.001
Kurdistan	62.6 (13.6)	60.4 (14.2)	67.7 (10.8)	0.001
Lorestan	60.1 (14.5)	59.0 (14.0)	63.0 (15.4)	0.001
Markazi	61.0 (13.2)	59.7 (13.5)	64.9 (11.5)	0.001
Mazandaran	61.9 (13.2)	60.0 (13.3)	66.6 (11.6)	0.001
Qom	58.6 (12.7)	57.2 (12.0)	66.6 (13.7)	0.008
Semnan	59.1 (12.9)	58.1 (12.7)	62.0 (13.0)	0.012
Sistan & Baluchestan	60.3 (13.9)	58.0 (13.0)	66.0 (14.5)	0.001
Tehran	60.4 (13.5)	59.0 (13.3)	64.2 (13.3)	0.001
Yazd	60.8 (13.3)	59.7 (13.4)	63.9 (12.6)	0.001
Zanjan	64.4 (12.8)	63.6 (12.7)	66.8 (12.9)	0.18
Total	61.2 (13.4)	59.6 (13.3)	65.4 (12.6)	0.001

SD = standard deviation.

highest mean age was in Ardabil [68.4 (SD 15.0) years] and this was significantly different from the lowest age, in Hamadan [62.1 (SD 12.9) years] ( $P = 0.001$ ).

### Myocardial infarction incidence rate

Crude and age-standardized myocardial infarction incidence rates in the country's provinces are shown in Table 2. The crude myocardial infarction

incidence rate was 64.9 per 100 000 in the whole country. The highest crude myocardial infarction incidence rate was in North Khorasan at 126.7 per 100 000 people, followed by Yazd, Semnan, Kerman, West Azarbaijan, and Khouzeestan (121.7, 119.5, 118, 108.2 and 108.1 per 100 000 respectively).

After adjustment, the crude myocardial infarction incidence rate in the whole country increased to 73.3 per

100 000 (95% CI: 72.3%–74.3%). After standardization, the highest myocardial infarction incidence was in North Khorasan (152.5 per 100 000), followed by Kerman, Khouzeestan, Yazd, Semnan and West Azarbaijan (149.2, 143.7, 141.3, 132.5 and 130.8 per 100 000 respectively), and the lowest incidence was in Qom [24.5 per 100 000 (95% CI: 19.6%–29.9%)].

Figure 1 illustrates the geographical distribution of myocardial infarction

**Table 2 Crude and adjusted myocardial infarction incidence rates per 100 000 population in provinces of the Islamic Republic of Iran, 2012**

Province	No. of cases	Incidence (per 100 000 population)		
		Crude rate	Adjusted rate	95% CI
Alborz	852	73.2	97.8	91.0–104.5
Ardabil	506	91.0	108.4	98.8–117.9
West Azerbaijan	1 455	108.2	130.8	124.0–137.7
East Azerbaijan	636	35.4	39.9	36.7–43.0
Bushehr	173	39.9	58.6	49.6–67.7
Chaharmahal & Bakhtiari	321	85.5	105.0	93.3–116.7
Chaharmahal & Bakhtiari	103	43.3	61.6	49.3–73.9
Esfahan	1 079	45.5	52.3	49.1–55.5
Fars	1 019	49.0	58.7	55.0–62.3
Ghazvin	432	78.8	97.8	88.5–107.2
Gilan	1 226	92.0	96.6	91.2–102.1
Golestan	526	69.1	89.2	81.4–97.0
Hamadan	419	51.5	57.6	52.0–63.2
Hormozgan	203	34.5	49.3	42.2–56.5
Kerman	1 396	118.5	149.2	141.2–157.2
Kermnanshah	604	67.7	79.4	72.9–85.9
South Khorasan	212	76.5	81.3	70.1–92.6
Razavi Khorasan	1 161	44.9	53.5	50.3–56.5
North Khorasan	452	126.7	152.5	138.2–166.7
Khuzestan	1 957	108.1	143.7	137.2–150.3
Kohgiluyeh & Boyer-Ahmad	125	61.5	62.9	51.5–74.3
Kurdistan	259	39.1	47.2	41.4–53.1
Lorestan	670	87.9	106.4	98.1–114.7
Markazi	522	76.7	86.5	79.0–94.1
Mazandaran	1 333	85.9	97.3	92.0–102.6
Qom	99	19.8	24.5	19.6–29.5
Semnan	347	119.0	132.5	118.4–146.6
Sistan & Baluchestan	253	32.7	44.1	38.5–49.7
Tehran	1 684	49.9	51.4	47.6–55.1
Yazd	571	121.7	141.3	129.5–153.0
Zanjan	155	34.0	40.1	33.7–46.5
Total	20 750	64.9	73.3	72.3–74.3

CI = confidence interval.


incidence in the different provinces of the country.

## Discussion

This is the first report on the hospital-based age-standardized incidence of myocardial infarction in the Islamic Republic of Iran. Our study showed that the incidence rate of myocardial

infarction was higher than the mean national rate in more than half of the provinces. These provincial disparities in myocardial infarction incidence rates mirror a national study on noncommunicable diseases risk factors in the Islamic Republic of Iran that mapped the differences in the prevalence of cardiovascular disease risk factors across the country's provinces (19,20). The wide difference in myocardial

infarction incidence rates—ranging from 24.5 per 100 000 in Qom to 152.5 per 100 000 in North Khorasan—could be due to different genetic or environmental factors, disparities in disease risk factors and different socioeconomic status distribution across the provinces as well as to varying coverage in local myocardial infarction registries (8,12,20). This needs further investigation in future studies. The especially



**Figure 1** Incidence-adjusted rates of myocardial infarction in provinces of the Islamic Republic of Iran, 2012

low incidence in Qom could also be explained by inadequate registration of myocardial infarctions, and this too should be addressed in subsequent studies to obtain a more accurate estimate of myocardial infarction in this province.

In literature searches we found no studies using similar data on myocardial infarction incidence rates from the Islamic Republic of Iran. The mortality rate due to myocardial infarction has been reported as 85 per 100 000 in the Islamic Republic of Iran (6). In our study, the age-adjusted incidence of recorded cases of myocardial infarction was lower at 73.3 per 100 000. Potential reasons for this inconsistency are differences in measures, defective registration of the causes of death and cardiac arrest and/or registration of myocardial infarction instead of the correct cause of death (21,22). It should also be noted that myocardial infarction cases who died

at home or outside hospital were not included in the records of myocardial infarction in our study. In the Islamic Republic of Iran the mortality rate due to acute myocardial infarction before arrival at hospital has been reported to be 20% (23). About 20% of first myocardial infarctions were detected only on death certificates (24) and, in view of this, the estimated incidence rate of myocardial infarction in our study can be revised up to 87.9 per 100 000 people. In other words, approximately 4150 myocardial infarction cases were estimated to die before arriving in hospital. This indicates that, despite the failure to register some cases of myocardial infarction outside hospital, a low number of deaths due to myocardial infarction are estimated to be registered in the Islamic Republic of Iran and therefore our findings based on myocardial infarction cases registered in hospital seem reasonable.

The standardized incidence rate of myocardial infarction in this study is high compared with a 30-year study in Japan that reported myocardial infarction incidence rates of 7.4 and 27 per 100 000 in 1979 and 2008 respectively (25). Our results are also higher compared with age-adjusted, standardized mortality rates due to myocardial infarction in Korea in 2003 and 2007 at 30.86 and 28.5 per 100 000 respectively (26). However, the incidence rate was lower compared with a study in Finland reporting myocardial infarction incidence rates in men and women of 738.8 and 292.9 per 100 000 individuals respectively in a population of 35- to 84-year-olds (9). The rate was also lower compared with other parts of the WHO Eastern Mediterranean Region; in Kuwait the myocardial infarction incidence rate was reported to be 130.1 per 100 000 patients admitted to hospitals (27). The Gulf Registry of Acute Coronary Events reported that the highest acute coronary syndrome incidence rate in Arab countries was in Oman, where the crude and standardized incidences of myocardial infarction with ST-elevation myocardial infarction were 92.8 and 184.9 per 100 000 respectively (28). The rate of myocardial infarction we found in the Islamic Republic of Iran is lower than in Oman. Since a population-based myocardial infarction incidence rate has not been yet reported from Arab countries, and in view of the mortality rate due to acute coronary syndrome in Yemen (10%), Oman (9.6%) and other Arab countries (3.3%), the myocardial infarction incidence rate seems to be higher in the Islamic Republic of Iran than in Arab countries (except for Oman, Kuwait and Yemen) (27–29).

The incidence rate of coronary heart disease in developed countries has been reported to range from 200–500 per 100 000 in men and 60–150 per 100 000 in women (30). In one study in the United Kingdom

the incidence rate of acute myocardial infarction was 0.48 per 100 000 and the rate of all coronary events was 91 per 100 000. Our rate of myocardial infarction was higher compared with that study, but lower in comparison with all coronary events (30).

The age of male and female patients (59.6 and 65.4 years respectively) and the total population (61.2 years) in our study in the Islamic Republic of Iran was lower compared with studies in Japan (25), Puerto Rico (31) and The Netherlands (10), which reported the mean ages of myocardial infarction patients as 66, 67 and 75 years respectively. On the other hand, the mean age of incidence in our study was higher compared with a study in India (57.5 years) (7). In Kuwait, 87% of myocardial infarction patients were male and the mean age was reported to be 55 years (27), which is lower than in our study. According to registries of acute coronary events in 65 hospitals in 6 Arab countries (Yemen, Oman, Bahrain, Kuwait, Qatar and the United Arab Emirates), the mean age at incidence was 55 years (29). Our study indicates that mean age at incidence of myocardial infarction is higher in the Islamic Republic of Iran than in these Arab countries (except for Yemen, with a mean age at incidence of 61 years). The difference in mean age across different studies could be attributed to different population age distributions, differences in life expectancies and lifestyles and variations in distribution of and coping with cardiovascular disease risk factors.

In the present study, the majority of the patients were male and the male:female incidence ratio was 2.63. This ratio is similar to studies in other countries (9,27–29,32); for example, in Finland the male:female ratio was reported to be 2.52 (9). The sex ratio in the Islamic Republic of Iran was similar to that reported by the Gulf

Registry of Acute Coronary Events, in which 76% of the patients were male (29).

Although the quality of hospital registry of myocardial infarction data in the Islamic Republic of Iran has been reported as acceptable (33), failure to register myocardial infarction cases who die outside hospital and/or at home was one of the limitations of the present study. We recommend that this is addressed in future research, accompanied by periodical assessments of the myocardial infarctions registry in order to determine the accuracy of the registered cases until the registry of data of myocardial infarctions becomes consistent in the country. Among the strengths of this study were that it was hospital-based, it included all the provinces of the country (based on formal provincial borders), it estimated the incidence using population denominators from the Statistical Center of Iran, as per WHO standards, it used a definite diagnosis by a cardiologist, and it was the first study on the incidence of myocardial infarction in the whole country.

## Conclusion

Reports of age-adjusted incidence rates of myocardial infarction in developing countries, particularly by province or state, are rare. Without such an index, planners of health systems inevitably use alternative indices such as mortality. We report for the first time the myocardial infarction incidence rates of hospitalized patients at provincial level in the Islamic Republic of Iran. The standardized incidence rate of myocardial infarction in the whole country (73.3 per 100 000) was higher than in some Arab countries of the Eastern Mediterranean Region (except for Oman, Kuwait and Yemen). The mean age

of myocardial infarction in men was significantly lower than in women. As myocardial infarction is the principal cause of death in the community, estimates of myocardial infarction incidence rate by province are valuable in health policy-making, resource allocation and prioritization and prevention and control of cardiovascular diseases. This study has paved the way for planning for and reducing disparities among the provinces, and monitoring and managing myocardial infarctions at the micro- and macro level in the Islamic Republic of Iran. The myocardial infarction cases who died at home or outside hospital were not included in our estimates of myocardial infarction incidence. It is recommended that the Iranian Myocardial Infarction Registry officials put in place a system for registering outside-hospital deaths in order to derive the actual myocardial infarction incidence rate in our country.

## Acknowledgements

We gratefully thank the personnel of the Cardiology Office (Dr Hozhabri and Mr Babrian), the personnel of the treatment deputies of the universities of medical sciences, the nurses and cardiologists in hospital coronary care units across the Islamic Republic of Iran and the officials and advisors of the Iranian Myocardial Infarction Registry Programme, particularly Dr Seyyed-Saeed Hashemi-Nazari.

**Funding:** This work was approved by the Management Centre of Non-Communicable Diseases and Office of Cardiovascular Diseases Prevention of the Iranian Ministry of Health and Medical Education (approval no. 305/837) and was part of a PhD thesis in Epidemiology by Ali Ahmadi at Shahid Beheshti University of Medical Sciences.

**Competing interests:** None declared.

## References

- Onen CL. Epidemiology of ischaemic heart disease in sub-Saharan Africa. *Cardiovasc J Afr.* 2013 Mar;24(2):34-42. PMID:23612951
- Khosravi A, Taylor R, Naghavi M, Lopez AD. Mortality in the Islamic Republic of Iran, 1964-2004. *Bull World Health Organ.* 2007 Aug;85(8):607-14. PMID:17768519
- Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al.; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Executive summary: heart disease and stroke statistics-2014 update: a report from the American Heart Association. *Circulation.* 2014 Jan 21;129(3):399-410. PMID:24446411
- Gaziano TA, Bitton A, Anand S, Abrahams-Gessel S, Murphy A. Growing epidemic of coronary heart disease in low- and middle-income countries. *Curr Probl Cardiol.* 2010 Feb;35(2):72-115. PMID:20109979
- Xavier D, Pais P, Devereaux PJ, Xie C, Prabhakaran D, Reddy KS, et al.; CREATE registry investigators. Treatment and outcomes of acute coronary syndromes in India (CREATE): a prospective analysis of registry data. *Lancet.* 2008 Apr 26;371(9622):1435-42. PMID:18440425
- Samavat T, Hojatzadeh E. Programs for prevention and control of cardiovascular diseases. Tehran: Ministry of Health and Medical Education, Health Deputy Press; 2012.
- Jeemon P, Reddy KS. Social determinants of cardiovascular disease outcomes in Indians. *Indian J Med Res.* 2010 Nov;132:617-22. PMID:21150014
- Mendis S, Thygesen K, Kuulasmaa K, Giampaoli S, Mähönen M, Ngu Blackett K, et al.; Writing group on behalf of the participating experts of the WHO consultation for revision of WHO definition of myocardial infarction. World Health Organization definition of myocardial infarction: 2008-09 revision. *Int J Epidemiol.* 2011 Feb;40(1):139-46. PMID:20926369
- Havulinna AS, Pääkkönen R, Karvonen M, Salomaa V. Geographic patterns of incidence of ischemic stroke and acute myocardial infarction in Finland during 1991-2003. *Ann Epidemiol.* 2008 Mar;18(3):206-13. PMID:18280920
- Jolobe OM. Incidence of recognized and unrecognized myocardial infarction in men and women aged 55 and older: the Rotterdam Study. *Eur Heart J.* 2006 Jun;27(11):1383-4. PMID:16603576
- Pedigo A, Aldrich T, Odoi A. Neighborhood disparities in stroke and myocardial infarction mortality: a GIS and spatial scan statistics approach. *BMC Public Health.* 2011;11:644. PMID:21838897
- Shafiee ENI, Asadi M, Haery Nejad M, Veisizadeh B, Kosary A. Myocardial infarction event rate in Bushehr Port: WHO MONICA Project. *ISMJ.* 2004;6(2):144-50 [Persian].
- Fakhrzadeh H, Bandarian F, Adibi H, Samavat T, Malekafzali H, Hodjatzadeh E, et al. Coronary heart disease and associated risk factors in Qazvin: a population-based study. *East Mediterr Health J.* 2008 Jan-Feb;14(1):33-41. PMID:18557450
- Talaei M, Sarrafzadegan N, Sadeghi M, Oveisgharan S, Marshall T, Thomas GN, et al. Incidence of cardiovascular diseases in an Iranian population: the Isfahan Cohort Study. *Arch Iran Med.* 2013 Mar;16(3):138-44. PMID:23432164
- Yarahmadi Sh, Etamad K, Hazaveh AM, Azhang N. Urbanization and non-communicable risk factors in the capital city of 6 big provinces of Iran. *Iran J Public Health.* 2013;42(1) Suppl1:113-8. PMID:23865027
- Moradi G, Mohammad K, Majdzadeh R, Ardakani HM, Naieni KH. Socioeconomic inequality of non-communicable risk factors among people living in Kurdistan Province, Islamic Republic of Iran. *Int J Prev Med.* 2013;4(6):671-83.
- Myocardial infarction registry [Persian]. Tehran: Cardiovascular Office, Ministry of Health and Medical Education; 2009. (<http://ehr2.behdasht.gov.ir/miregistry/>, accessed 23 October 2014).
- Ahmad OB, Lopez AD, Murray CJ, Lozano R, Inoue M. Age standardization of rates: a new WHO standard. GPE Discussion Paper Series No. 31. Geneva: World Health Organization; 2001 (<http://www.who.int/healthinfo/paper31.pdf>, accessed 23 October 2014).
- Ahmadi A, Mobasheri M, Soori H. Prevalence of major coronary heart disease risk factors in Iran. *Int J Epidemiol Res.* Forthcoming 2014.
- Esteghamati A, Etamad K, Koohpayehzadeh J, Abbasi M, Meysamie A, Noshad S, et al. Trends in the prevalence of diabetes and impaired fasting glucose in association with obesity in Iran: 2005-2011. *Diabetes Res Clin Pract.* 2014 Feb;103(2):319-27. PMID:24447808
- Yavari P, Abadi AR, Mehrabi Y. Mortality and changing epidemiological trends in Iran during 1979-2001. *Hakim Research Journal.* 2004;6(3):14-7 [Persian].
- Riska O. Misclassification of deaths in Iran. *Tidsskr Nor Laegeforen.* 2008 9;128(19):2230.
- Talebizadeh N, Haghdoost A, Mirzazadeh A. An epidemiological model (Markov Chain) of cardiovascular disease in Iran. *Payesh. Journal of The Iranian Institute For Health Sciences Research.* 2010;8(2):163-70 [Persian].
- Greenlee RT, Naleway AL, Vidaillet H. Incidence of myocardial infarction in a general population: the Marshfield Epidemiologic Study Area. *WMJ.* 2002;101(7):46-52. PMID:12426920
- Takii T, Yasuda S, Takahashi J, Ito K, Shiba N, Shirato K, et al.; MIYAGI-AMI Study Investigators. Trends in acute myocardial infarction incidence and mortality over 30 years in Japan: report from the MIYAGI-AMI Registry Study. *Circ J.* 2010 Jan;74(1):93-100. PMID:19942783
- Jhun HJ, Kim H, Cho SI. Time trend and age-period-cohort effects on acute myocardial infarction mortality in Korean adults from 1988 to 2007. *J Korean Med Sci.* 2011 May;26(5):637-41. PMID:21532854
- Al-Owaish RA. Descriptive epidemiology of acute myocardial infarction in Kuwait, 1978. *Bull World Health Organ.* 1983;61(3):509-16. PMID:6603924
- Al-Lawati J, Sulaiman K, Panduranga P. The epidemiology of acute coronary syndrome in Oman: results from the Oman-RACE study. *Sultan Qaboos Univ Med J.* 2013 Feb;13(1):43-50. PMID:23573381
- Thalib L, Zubaid M, Rashed W, Almahmeed W, Al-Lawati J, Sulaiman K, et al. Regional variability in hospital mortality in patients hospitalized with ST-segment elevation myocardial infarction: findings from the Gulf Registry of Acute Coronary Events. *Med Princ Pract.* 2011;20(3):225-30. PMID:21454991
- Rothwell PM, Coull AJ, Silver LE, Fairhead JF, Giles MF, Love-lock CE, et al.; Oxford Vascular Study. Population-based study of event-rate, incidence, case fatality, and mortality for all acute vascular events in all arterial territories (Oxford Vascular Study). *Lancet.* 2005 Nov 19;366(9499):1773-83. PMID:16298214
- Zevallos Juan Carlos YJ, Banchs Héctor, Juan A, Sánchez González, Mattei Hernando. Gender disparities in Puerto Ricans hospitalized with an initial acute myocardial infarction.

- tion: a population-based Perspective. *P R Health Sci J.* 2012;31(4):192–8. PMID:23844466
32. Zubaid M, Rashed WA, Husain M, Mohammad BA, Ridha M, Basharithulla M, et al. A registry of acute myocardial infarction in Kuwait: Patient characteristics and practice patterns. *Can J Cardiol.* 2004 Jun;20(8):783–7. PMID:15229759
33. Shahraz S, Barzanjeh A, Bahari A, Nadery Y, Farzadfar F, Beyranvand MR, et al. The completeness of medical records to assess quality of hospital care: the case of acute myocardial infarction in a district-level general hospital in Iran. *Arch Iran Med.* 2012 Oct;15(10):592–5. PMID:23020532