

Low-risk planned caesarean versus planned vaginal delivery at term: early and late infantile outcomes

Najmeh Maharlouei,¹ Parisa Mansouri,² Mozhgan Zahmatkeshan³ and Kamran B. Lankarani⁴

¹Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. ²Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. ³Gastroenterohepatology Research Center, Nemazee Teaching Hospital, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. ⁴Health policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. (Correspondence to: Najmeh Maharlouei: najmeh.maharlouei@gmail.com).

Abstract

Background: Understanding the impact of delivery mode on neonatal morbidity becomes essential in the context of rising Caesarian delivery rates.

Aims: We aimed to compare the selected outcomes in neonates born by low-risk planned Caesarian delivery versus planned normal vaginal delivery (NVD).

Methods: This prospective cohort study examined early, and late neonatal complications among 1071 neonates born through low-risk planned Caesarian delivery and 1367 neonates born through planned NVD, in Fars, Islamic Republic of Iran, during 2012–2014.

Results: Gestational age of neonates born through Caesarian delivery was significantly lower than their counterparts in NVD group. Accordingly, babies' birth weights were 3166 (± 442.4) grams in Caesarian delivery group and 3213 (± 454.8) grams in NVD group. Normal skin colour at birth was more prevalent in the Caesarian delivery group compared to the NVD group (85% vs. 81.3%, $P = 0.04$). No significant differences were detected between the two groups regarding birth trauma, birth height and head circumference, and developing infection, icterus and convulsion during neonatal period. Also, height and weight at two years of age did not significantly differ in both groups.

Conclusion: The results of this study show that neonates born by Caesarian delivery and NVD had the same early and late outcomes.

Keywords: vaginal delivery, caesarean delivery, outcome, cohort study, Iran

Citation: Maharlouei N; Mansouri P; Zahmatkeshan M; Lankarani K. B. Low-risk planned caesarean versus planned vaginal delivery at term: early and late infantile outcomes. *East Mediterr Health J.* 2018;25(7):503–513. <https://doi.org/10.26719/emhj.18.066>

Received: 16/03/17; accepted: 16/11/17

Copyright © World Health Organization (WHO) 2019. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license (<https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Introduction

The rate of caesarean delivery has increased worldwide over the past decades (1–3). There has been a rise in Caesarian delivery rate in the United States of America from 1996 to 2004 when 1.2 million women (29.1% of all births) had Caesarian delivery (4). In many high-income countries it has exceeded 30% over the past decades. In Germany from 1990 to 2010 it doubled from 15.7 to 31.9% of all births (5). Moreover; worldwide estimates of Caesarian delivery at maternal request are 6–8% in Northern Europe, 11.2% in the United States, 17.3% in Australia and 70% in Brazil (6). A significant rise in Caesarian delivery rate from 14.3% in 1979 to 85.3% in 2009 in the Islamic Republic of Iran has also been reported, although this was a hospital-based report (7).

Caesarian delivery has been considered as a global burden for many years (8) and has been the most common surgical procedure performed on American women (9). A higher risk of infant morbidity including breastfeeding complications (10), infections and respiratory distresses and even maternal bleeding has been reported to be associated with Caesarian delivery (11). Moreover, compared to planned vaginal delivery, planned CD was related to higher rate of severe maternal morbidity (12).

However, according to studies, Caesarian delivery is selected to avoid certain medical conditions such as fetal distress, fear from urinary sequelae, and anal incontinence (13,14). A main concern is whether the rise in the Caesarian delivery rate is the reason for the fall in perinatal mortality during the past decades (3). Although Caesarian delivery is more expensive than normal vaginal delivery (NVD) (15), it is more common in the private sector (16). Also, the rate of Caesarian delivery went beyond the rate recommended by the World Health Organization (WHO); i.e., 15% of all deliveries, over the past decades.

The decision to choose Caesarian delivery or NVD based only on the emotional status of a pregnant woman is due to evidence indicating a lack of difference in complication rates between Caesarian delivery and NVD patients (17). Within these controversies, understanding the impact of delivery mode on neonatal morbidity and mortality becomes essential in the context of rising Caesarian delivery rates. Therefore, the present study aims to compare the neonates born by low-risk planned Caesarian delivery and those born by planned NVD regarding selected neonatal outcomes.

Methods

Study design and population

This prospective cohort study is a part of the cohort study which has been started since 2012 in Fars, the fifth most populated province in the Islamic Republic of Iran (18). Aiming to compare neonatal outcome of low-risk planned Caesarian delivery and planned NVD, we included mothers whose infant was alive and delivery date was after 37th weeks, and who had no history of maternal or fetal complications. We also excluded those whose date of delivery was influenced by maternal or fetal complications, including premature labour pain, ruptured membrane, fetal intra-uterine growth retardation or meconium staining. The study was approved by the Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.REC.1397.464) (19).

Data collection

In this article we used data collected during fetal period, 2, 6 and 24 months after birth. Data were collected during pregnancy included demographic, medical history and obstetric and gynaecological history of mothers. The checklist used 2 months after delivery consisted of two main parts: the first part asked questions regarding mode of delivery and whether it was planned and if the mode of delivery was NVD. The next question was concerning type of NVD; routine NVD, physiologic NVD, NVD in water, or painless NVD in which intrathecal analgesic is used routinely. Those mothers who experienced Caesarian delivery were asked about the reason of Caesarian delivery.

In the next part of the study, we asked mothers to answer questions regarding neonatal outcomes based on the infants' health card, vaccination card as well as the diary we had given the mothers in the first phase of the study (during pregnancy). In this part, we asked about infant's weight, height and head circumference at birth as well as presence of congenital anomalies (if any). Since APGAR is not recorded in either the health card or the vaccination card, we had to ask mothers about the neonate's skin colour, crying, and limbs' movement after birth and the place the neonate was kept immediately after birth; i.e. next to mother or in a neonatal intensive care unit (NICU)/neonatal ward. We also asked about birth trauma including bone fracture(s) and/or dislocation(s) and scalp hematoma, if the child had developed icterus, infection, and convulsion in the neonatal period. The history was considered positive if the diagnosis was confirmed by a paediatrician, or the infant was admitted to hospital due to the mentioned problems. We also asked if breastfeeding was started within two hours after delivery, and the duration of exclusive breastfeeding.

After the infants passed their 6 and 24 months, the mothers were contacted and asked about any ailments/disease(s) their children had developed that necessitated a visit to a doctor. Also details of infants' feeding including duration of exclusive breastfeeding and weaning were requested, as well as information about

any disease(s) confirmed by a physician, any hospital admissions, reason(s), and frequency and duration of each hospitalization (if any).

Statistical analysis

Statistical analyses were performed using SPSS statistical software, ver. 18.0 (SPSS Inc., Chicago, IL: USA). All the participants were categorized into two groups; those who had planned NVD and those who had planned CD. All the comparisons were made between these two groups. Independent *t*-test was used to compare the quantitative variables, while chi-square test or Fisher's exact test was employed to compare the qualitative ones. All differences with *P*-values less than 0.05 were considered as statistically significant. The data were reported as mean \pm standard deviation (SD) and frequency (percentage) as indicated.

Results

Two months after delivery, we interviewed 4577 out of the 6922 mothers (66.1%) who had participated in the first phase of the cohort study, which have been started in 2012 (17). We found that 72 respondents (1.8%) had experienced either still birth (32; 0.7%) or neonatal mortality (40; 0.8%). Among the rest of the respondents (4505 mothers) whose pregnancies resulted in live births, 1250 mothers (27.7%) had emergency delivery – either NVD or Caesarian delivery. Hence, of the 3255 mothers who had experienced planned deliveries, 1457 had planned NVDs (44.8%) and 1798 had planned Caesarian deliveries (55.2%). Considering that we aimed to compare neonatal outcome of low-risk pregnancies, we excluded all planned deliveries in which mother and/or the fetus was considered high risk by the maternity care provider. Therefore, data of 1367 low-risk planned NVD and 1071 low-risk planned Caesarian delivery were analyzed. Different indications of Caesarian delivery have been illustrated in Figure 1.

Comparison of maternal and neonatal characteristics between the two groups – low-risk planned NVD versus low-risk planned Caesarian delivery – is presented in Table 1, which indicates that the mean age of mothers in NVD group was slightly lower than that of the Caesarian delivery group (26.5 vs. 28.4 years; $P < 0.001$), which was not clinically significant. Besides, the mothers who had Caesarian delivery were more likely to have a higher education level and significantly more likely to be employed (12.1%) compared to their counterpart in NVD group (5%; $P < 0.001$). A remarkably higher proportion of mothers who gave birth through Caesarian delivery had received maternity care at private clinics (46.4% vs. 20.3%, $P < 0.001$). Also, they are more likely to have a history of infertility (9.2% vs. 4.01%; $P < 0.001$), abortion/stillbirths (23.1% vs. 16.5%; $P < 0.001$), and children with physical and/or mental disabilities (2.4% vs. 0.9%; $P = 0.003$). Yet, health insurance coverage was similar in both groups ($P = 0.4$). In Caesarian delivery group, 9 (0.8%) twin and 1 (0.1%) triplet deliveries were reported, while in NVD group just 1 (0.1%) twin pregnancy was mentioned. No significant

Table 1: Comparing maternal and neonatal characteristics the group who had planned NVD versus those who had C-section

Characteristics	Planned NVD N = 1367	Planned C-section N = 1071	P-value
Maternal			
Age (years)	26.5 (± 4.9)	28.4 (± 4.7)	< 0.001*
No. of previous children	2 (0-6)	2 (0-5)	0.04 [†]
Up to secondary school	704 (51.5)	331 (31)	< 0.001 ^{§§}
High school diploma	470 (34.4)	405 (37.7)	
Undergraduate degree	186 (13.6)	317 (29.6)	
Postgraduate degree	7 (0.5)	18 (1.7)	
Employment			
Employed	69 (5)	130 (12.1)	< 0.001 [§]
Unemployed (housewife/student)	1298 (95)	942 (87.9)	
Place of received maternity care			
Private clinics	277 (20.3)	497 (46.4)	< 0.001 [§]
Governmental Clinics	1090 (79.7)	574 (53.6)	
Having insurance	1270 (92.9)	992 (92.6)	0.4 [§]
Positive history of infertility	56 (4.1)	99 (9.2)	< 0.001 [§]
Positive history of abortion/stillbirth	225 (16.5)	247 (23.1)	< 0.001 [§]
Having children with physical and/or mental problem	12 (0.9)	26 (2.4)	0.003 [§]
Place of delivery			
Hospital	1165 (85.3)	1026 (95.9)	< 0.001 ^{§§§}
Maternity hospital	188 (13.6)	45 (4.1)	
Facility centre	1 (0.1)	0	
Gynaecologist clinic	8 (0.6)	0	
At home	6 (0.4)	0	
Sex			
Male	675 (49.4)	543 (50.7)	0.55 ^{§§}
female	691 (50.5)	532 (49.3)	
Ambiguous	1 (0.1)	0	
Number of tones			
Single tone	1366 (99.9)	1061 (99.1)	0.007 ^{§§}
Twin	1 (0.1)	9 (0.8)	
Triplet	0	1 (0.1)	
Congenital anomalies	11 (0.8)	7 (0.7)	0.8 [§]

*Independent T test

†Two-independent samples tests

§Chi-square test

§§Fisher's exact test

P values less than 0.05 is considered significant.

Data are reported as mean (± standard deviation), Median (min - max), and frequency (%); NVD = normal vaginal delivery

difference was observed between the NVD and Caesarian delivery groups regarding the infants' gender.

Comparison of the immediate neonatal outcomes between planned NVD and low-risk planned Caesarian delivery groups is presented in Table 2. Accordingly, gestational age (39.2 vs. 38.4 weeks; $P < 0.001$) and birth weight (3213 vs. 3166 grams; $P < 0.001$) were significantly higher in the NVD group compared to the Caesarian delivery group. However, the differences were not

clinically significant. No statistical differences were found between the two groups in neonates' head circumference and height at birth. The newborns delivered through Caesarian delivery were more likely to have normal skin colour compared to those delivered via NVD (85% vs. 81.3%; $P = 0.04$). However, no significant differences were detected between the neonates of two groups regarding grade of crying, limbs movement vigor, respiration, skull hematoma, birth trauma, and place of care after birth (next to mother or in NICU/neonatal

Figure 1: Flowchart of participants' recruitment




Table 2: Comparing immediate neonatal outcomes in planned NVD versus planned C-Section

Neonatal outcomes	Planned NVD N = 1367	Planned C-section N = 1071	P-value
Gestational age at birth (weeks)	39.2 (± 1.2)	38.4 (± 1.2)	< 0.001*
Birth weight (grams)	3213 (± 454.8)	3166 (± 442.4)	0.01*
Height at birth (centimetres)	50.2 (± 2.8)	50 (± 3.0)	0.1*
Head circumference	34.8 (± 1.6)	35 (± 1.5)	0.2*
Place of care after birth			
Next to mother	1283 (93.9)	997 (92.9)	0.4§
NICU or neonatal ward	84 (6.1)	77 (7.1)	
Crying			
Vigorous	1302 (95.2)	1034 (96.5)	0.2§
Weak	54 (4.0)	32 (3.0)	
No crying	11 (0.8)	5 (0.5)	
Skin colour			
Pink/normal	1112 (81.3)	910 (85.0)	0.04§
Reddish	221 (16.2)	144 (13.4)	
Cyanotic/pale	34 (2.5)	17 (1.6)	
Movement			
Vigorous	1349 (98.7)	1060 (99)	0.5§§
Weak	14 (1.0)	10 (0.9)	
Poor/none	4 (0.3)	1 (0.1)	
Respiration			
Normal	1331 (97.4)	1043 (97.4)	0.3§
Nasal oxygen	27 (2.1)	25 (2.3)	
Intubation	9 (0.6)	3 (0.3)	
Birth trauma			
Skull hematoma	5 (0.4)	2 (0.2)	0.3§§
Skull deformity	1 (0.1)	0	0.6§§
Clavicle	5 (0.4)	3 (0.3)	0.5§§
Femur	0	2 (0.2)	0.2§§

*Independent t test

§Chi-square test

§§Fisher's exact test

P values less than 0.05 is considered significant.

Data are reported as mean (\pm standard deviation), or frequency (%); NVD = normal vaginal delivery**Table 3: Comparing early neonatal outcomes in both groups; planned NVD versus planned C-Section**

Neonatal outcomes	Planned NVD N = 1367	Planned C-section N = 1071	P-value
Icterus			
Developing icterus	469 (34.3)	381 (43.9)	0.52§
Start day	3.8 (\pm 2.3)	3.6 (\pm 2.5)	0.6*
Duration of icterus	1.6 (\pm 0.6)	1.7 (\pm 0.5)	0.07*
Blood exchange	3.0 (0.6)	0	0.58§
Phototherapy	323 (69.3)	270 (70.9)	
Medication	14 (3.0)	9 (2.4)	
Home remedy &/ breastfeeding	221 (16.2)	144 (13.4)	
Photo and breast feeding cessation and medication	124 (26.4)	100 (26.2)	

Table 3: Comparing early neonatal outcomes in both groups; planned NVD versus planned C-Section (concluded)

Neonatal outcomes	Planned NVD N = 1367	Planned C-section N = 1071	P-value
Infection			
Developing infection	75 (5.5)	70 (6.5)	0.3 [§]
Start day	5.1 (± 4.7)	6.8 (± 8.1)	0.09*
Hospitalization due to infection ⁶	16 (21.3)	7 (10.0)	0.2 [§]
Duration of admission	2.1 (± 1.4)	2 (± 1.4)	0.8*
Convulsion			
Developing convulsion	13 (1.0)	12 (1.1)	0.8 ^{§§}
Start day	7.8 (± 2.6)	9.9 (± 2.5)	0.7*
Hospitalization due to convulsion ⁶⁶	4 (30.8)	3 (25.0)	0.1 [§]

*Independent T test

§Chi-square test

§§Fisher's exact test

⁶The proportion of neonates admitted due to infection⁶⁶The proportion of neonates admitted due to convulsion

P values less than 0.05 is considered significant.

Data are reported as mean (± standard deviation), or frequency (%); NVD = normal vaginal delivery

Table 4: Comparing outcomes during the first two years of life in children born through planned NVD versus planned C-section

Late outcomes	Planned NVD N = 1367	Planned C-Section N = 1071	P-value
Height at 2 years old (centimetres)	84.6 (± 4.0)	85.7 (± 3.9)	0.1*
Weight at 2 years old (kilogrammes)	11.6 (± 3.6)	11.8 (± 1.5)	0.3*
Hospital admission/paediatrician visit			
The child visited by a doctor (for reasons other than check-up)	471 (34.5)	332 (31.0)	0.07 [§]
History of hospital admission	202 (14.8)	183 (17.1)	0.1*
Duration of total hospitalization ³	5 (1-105)	4 (1- 63)	0.9**
Congenital anomalies/diseases			
Presence of at least one congenital anomaly/diseases	113 (8.3)	91 (8.5)	0.4 [§]
n (n/113*100) n (n/91*100)			
G6PD deficiency	79 (69.9)	73 (80.2)	
Thalassaemia minor	16 (14.1)	5 (5.5)	
Musculoskeletal (club foot, polydactyly)	4 (3.5)	2 (2.2)	
Cardiac problems	2 (1.8)	7 (7.7)	
Oro-pharyngeal problems	3 (2.6)	3 (3.3)	
Urogenital problems	3 (2.6)	1 (1.1)	
Down Syndrome	2 (1.7)	1 (1.1)	
Ophthalmic problems	1 (0.8)	2 (2.2)	
Other problems	7 (6.2)	3 (3.3)	
Developing diseases during the first two years			
Dermatologic problems	43 (3.2)	34 (3.1)	0.9 [§]
Respiratory problems including asthma	19 (1.4)	13 (1.2)	0.6 ^{§§}
Nephrology problems	4 (0.3)	2 (0.2)	0.7 ^{§§}
Neurologic problems	4 (0.3)	1 (0.1)	0.5 ^{§§}
Musculoskeletal problems	1 (0.1)	1 (0.1)	0.9 ^{§§}

*Independent T test

**Mann-Whitney Test

³Duration of total hospitalization= summation of number of hospital admission* duration of each admission

§Chi-square test

§§Fisher's exact test

P values less than 0.05 is considered significant.

Data are reported as mean (± standard deviation), or frequency (%); NVD = normal vaginal delivery

Table 5: Comparing breast feeding pattern in children born through planned NVD versus planned C-Section

Breast feeding	Planned NVD N = 1367	Planned C-section N = 1071	P-value
Initiation of breast feeding			
Was not started at all	1 (0.1)	3 (0.2)	< 0.001 [§]
Within 2 hours after delivery	1316 (96.3)	1001 (93.5)	
Within 3 three days after delivery	50 (3.6)	67 (6.3)	
Duration of breast feeding (months)	19.9 (± 6)	18.8 (± 8.1)	< 0.001*
Less than 7 days after birth	20 (1.5)	19 (1.7)	< 0.001 ^{§§}
The first month of life	16 (1.2)	37(3.4)	
Up to 2 months old	15 (1.1)	20 (1.8)	
Up to 4 months old	35 (2.4)	62 (5.9)	
Up to 6 months old	46 (3.2)	56 (5.2)	
Up to 1 year old	54 (4.0)	67 (6.3)	
Up to 1.5 years old	149 (11)	92 (8.6)	
Up to 2 years old	1032 (75.5)	716 (66.9)	
For more than 2 years	0	2 (0.2)	
Duration of exclusive breast feeding (weeks) 20.9 (± 1.6)		19.8 (± 2.2)	< 0.001*
Never	15 (1.1)	16 (1.5)	0.01 ^{§§}
Less than one month	29 (2.1)	45 (4.2)	
Less than 4 months	101 (7.4)	107 (10)	
4 months	212 (15.5)	195 (18.2)	
6 months	1010 (73.9)	708 (66.1)	

*Independent T test

§Chi-square test

§§Fisher's exact test

P values less than 0.05 is considered significant.

Data are reported as mean (± standard deviation), or frequency (%); NVD = normal vaginal delivery

ward). Neonatal outcomes are described in Table 3. Neonatal complications did not statistically differ in those born through Caesarian delivery and those in NVD group; i.e., icterus ($P = 0.52$), infection ($P = 0.3$) and convulsion ($P = 0.8$).

Certain aspects of health status during the first two years of life were also compared between the two groups and the results are presented in Table 4. Accordingly, no significant difference was found between the two groups regarding height ($P = 0.1$) and weight ($P = 0.3$) at two years of age. Nevertheless, compared to the children born through Caesarian delivery, those born through NVD had more frequently developed ailment/disease(s) for which they had to be visited by a doctor during the first two years of life (34.5% vs. 31%; $P = 0.07$). However, those born through Caesarian delivery had been more frequently hospitalized (17.1% vs. 14.8%; $P = 0.1$). Yet, the two groups were similar regarding presence of congenital anomalies/diseases ($P = 0.4$) and developing nephrological ($P = 0.7$), dermatological ($P = 0.9$), neurological ($P = 0.5$), musculoskeletal ($P = 0.9$), and respiratory problems including asthma ($P = 0.6$).

According to what had been recorded by mothers in their diaries, breastfeeding had never been started for

three infants (0.2%) in Caesarian delivery group and one infant (0.1%) in NVD group. The reason was multiple congenital anomalies in oropharynx of the infants. Breastfeeding was more likely to be started within two hours after delivery in the neonates born through NVD compared to neonates in Caesarian delivery group (96.3% vs. 93.5%). For the remainder of the neonates, breastfeeding was started within three days; 3.6% in NVD group comparing to 6.3% in Caesarian delivery group. Duration of breastfeeding was also longer in the NVD group compared to the Caesarian deliverygroup (19.9 vs. 18.8 months, $P < 0.001$). Also, the duration of exclusive breastfeeding was significantly ($P < 0.001$) higher in NVD group compared to Caesarian delivery infants; 20.9 weeks (± 1.6) vs. 19.8 weeks (± 2.2), respectively.

Discussion

Appropriateness of Caesarian delivery and NVD for pregnant women has been widely debated. In this cohort study, we compared maternal and neonatal characteristics as well as immediate, early, and late neonatal complications among the women who delivered their babies by NVD and Caesarian delivery. The mean age of the mothers was lower in the NVD group than in the Caesarian

delivery group. In addition, the mothers in the CD group were more likely to be more educated and be employed. Besides, a significantly higher proportion of Caesarian deliveries were performed at private settings. Similar results were also obtained in other studies (15,16).

The results of our study demonstrated a higher birth weight in the infants delivered by NVD. However, birth weight was reported to be higher in the Caesarian delivery neonates than in the NVD ones (20). The longer gestational age in the NVD group in our study could be a reason for fetal weight gain. Also, the infants born through Caesarian delivery were more likely to have normal pink skin at birth. However, no significant differences were detected between the two groups regarding birth height, place of care after birth (next to mother or in neonatal ward), type of crying, movement, respiration, skull hematoma, and birth trauma. Some studies have reported that mode of delivery could not be related to birth trauma (21), but it has also been reported that Caesarian delivery is protective against birth trauma (22). Other studies have reported an association between Caesarian delivery and newborn's serious respiratory morbidity (23,24), respiratory distress (25), and transient tachypnea (26). In addition, delivery by Caesarian delivery has been identified as a risk factor for child asthma (27), respiratory morbidity, and longer NICU stay. Conversely, delivery by Caesarian delivery was reported not to be associated with the subsequent development of asthma, allergic rhinitis, or atopic dermatitis in Korean children (28).

Evaluation of early neonatal outcomes in the groups under study did not show any differences between the two groups regarding development of icterus, infection and other early neonatal complications. This was in agreement with the results of another study indicating no significant difference between NVD and Caesarian delivery neonates concerning neonatal complications (29). In a Chinese study, short-term maternal outcomes were similar in NVD and Caesarian delivery mothers, and Caesarian delivery women had even better neonatal benefits (30). In contrast, a Swedish study reported that Caesarian delivery either without medical indication or in emergency situations were associated with higher risks for maternal and neonatal morbidity (11).

The study results also revealed no significant difference between the two groups with respect to height and weight at the end of the second year of life. In contrast, Caesarian delivery has been reported as a risk factor for child obesity (27). Moreover, a systematic review reported a strong association between Caesarian delivery and increased offspring Body Mass Index (BMI), overweight, and obesity in adulthood (31). Compared to vaginally born

infants, Caesarian delivery infants have different timing, composition, and acquirement of intestinal flora, which may contribute to intestinal microbial composition in the first year of life, causing obesity and other health outcomes (27).

There was no significant difference between the NVD and Caesarian delivery group regarding the rate and length of hospital admission. This was in contrast to the results of other studies, showing higher newborn hospitalization for vaginally born neonates (20). Nevertheless, similar to the present research, that study demonstrated no significant difference between Caesarian delivery and NVD born infants with regard to duration of hospital stay. It has been shown that in preterm births, Caesarian delivery significantly increases the risk of longer neonatal length of hospital stay compared to NVD (32). Our study results also indicated no significant differences between the two groups regarding dermatological, nephrological, spinal cord, and musculoskeletal problems.

In this study, a slightly higher but significant proportion of the Caesarian delivery neonates did not start breastfeeding at all. Duration of breastfeeding was higher in the NVD group than in the Caesarian delivery group. Compared to the Caesarian delivery infants, a significantly higher proportion of the NVD infants had six-month exclusive breastfeeding. These findings were consistent with those of other studies (29,32,33).

The main strength of this study is that all data have been collected prospectively in a cohort study by trained research assistants and we did not use retrospectively collected hospital data. The limitation of the study is that we had to rely on mothers' claims, although they were trained to write all important events regarding their baby as well as themselves in the notebook given to them during their.

Conclusion

The results of this study show that neonates born by Caesarian delivery and normal vaginal delivery had the same early and late outcomes. Also Caesarian delivery neonates were more likely to have normal pink skin at birth. However, breastfeeding habits were better among NVD neonates. Considering the controversies reported in studies on Caesarian delivery and NVD outcomes, more research is needed to evaluate the short-term and long-term effects of Caesarian delivery.

Acknowledgements

We would like to express our special gratitude to the Vice-chancellor for Research of Shiraz University of Medical Sciences for their financial support. It should be noted that this article is extracted from Parisa Mansouri's thesis (No: 87/1006).

Funding: Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran.

Competing interests: None declared.

Césarienne planifiée à faible risque par rapport à l'accouchement vaginal planifié à terme : résultats précoces et tardifs pour l'enfant

Résumé

Contexte : Comprendre l'impact du mode d'accouchement sur la morbidité néonatale devient essentiel dans le contexte de la hausse des taux de césariennes.

Objectifs : Nous voulions comparer certains résultats chez les nouveau-nés par césarienne planifiée à faible risque par rapport à l'accouchement vaginal normal planifié.

Méthodes : L'étude de cohorte prospective a examiné les complications néonatales précoces et tardives chez 1071 nouveau-nés suite à une césarienne planifiée à faible risque et 1367 nouveau-nés suite à un accouchement vaginal planifié, dans la province de Fars, République islamique d'Iran, entre 2012 et 2014.

Résultats : L'âge gestationnel des nouveau-nés par césarienne était significativement inférieur à celui de leurs homologues du groupe des accouchements vaginaux normaux. Par conséquent, le poids à la naissance des bébés était de 3166 ($\pm 442,4$) grammes dans le groupe des césariennes et de 3213 ($\pm 454,8$) grammes dans le groupe des accouchements vaginaux normaux. La couleur normale de la peau à la naissance était plus fréquente dans le groupe des césariennes que dans l'autre groupe (85 % contre 81,3%, $p = 0,04$). Aucune différence significative n'a été détectée entre les deux groupes en ce qui concerne les traumatismes à la naissance, la taille de naissance et le périmètre crânien, ainsi que le fait de contracter une infection, un ictère et les convulsions durant la période néonatale. De plus, la taille et le poids à l'âge de deux ans ne différaient pas de façon significative dans les deux groupes.

Conclusion : Les résultats de cette étude montrent que les nouveau-nés par césariennes et par accouchement vaginal normal présentent les mêmes résultats précoces et tardifs.

الولادة القيصرية منخفضة المخاطر المخطط لها مقابل الولادة المهبلية في موعدها المخطط لها: مخرجات الولادة المبكرة والمتأخرة بالنسبة للرضع

نجمة مهارلوي، بريس منصور، موزكان زحمتكشان، كامران باقري لنكراني

الخلاصة

الخلفية: أصبح فهم تأثير نوع الولادة على المراضة بين الأطفال حديثي الولادة ضرورياً في سياق معدلات الولادة القيصرية المتزايدة.

الأهداف: هدفتنا إلى مقارنة المخرجات المحددة بين الأطفال حديثي الولادة الذين وُلدوا من خلال الولادة القيصرية منخفضة المخاطر المخطط لها مقابل الولادة المهبلية الطبيعية المخطط لها.

طرق البحث: بحثت هذه الدراسة الأترابية الاستباقية مضاعفات الولادة المبكرة والمتأخرة بين ١٠٧١ طفلاً حديث الولادة وُلدوا من خلال الولادة القيصرية منخفضة المخاطر المخطط لها و ١٣٦٧ طفلاً حديث الولادة وُلدوا من خلال الولادة المهبلية الطبيعية المخطط لها، في مدينة فارس، جمهورية إيران الإسلامية، في الفترة بين عامي ٢٠١٢ و ٢٠١٤.

النتائج: كان العمر الحملي للأطفال حديثي الولادة الذين وُلدوا من خلال الولادة القيصرية أقل بكثير من العمر الحملي للأطفال حديثي الولادة الذين وُلدوا من خلال الولادة المهبلية الطبيعية. لذلك، كان الوزن عند الولادة ٣١٦٦ (± ٤٤٢) جراماً في فئة الأطفال الذين وُلدوا من خلال الولادة القيصرية، و ٣٢١٣ (± ٤٥٤) جراماً في فئة الأطفال الذين وُلدوا من خلال الولادة المهبلية الطبيعية. وكان لون البشرة الطبيعي عند الولادة أكثر انتشاراً بين فئة الأطفال الذين وُلدوا من خلال الولادة القيصرية مقارنة بفئة الأطفال الذين وُلدوا من خلال الولادة المهبلية الطبيعية (٨٥٪ مقابل ٨١,٣٪، $P = ٠,٠٤$). ولم يُلاحظ أي اختلافات جوهرية بين الفئتين فيما يتعلق برضح الولادة، والطول عند الولادة، ومحيط الرأس، والإصابة بالعدوى، واليرقان والاختلاجات في فترة الولادة الحديثة. ولم يختلف الطول والوزن عند عمر الستين اختلافاً كبيراً بين الفئتين.

الاستنتاجات: توضح نتائج هذه الدراسة أن الأطفال حديثي الولادة الذين وُلدوا من خلال الولادة القيصرية منخفضة المخاطر والولادة المهبلية الطبيعية لديهم نفس حصائل الولادة المبكرة والمتأخرة.

References

1. Ba'aqeel HS. Cesarean delivery rates in Saudi Arabia: a ten-year review. Ann Saudi Med. 2009;29(3):179–83. <http://dx.doi.org/10.5144/0256-4947.51773>
2. Gould JB, Danielsen B, Korst LM, Phibbs R, Chance K, Main E, et al. Cesarean delivery rates and neonatal morbidity in a low-risk population. Obstet Gynecol. 2004;104(1):11–9. <http://dx.doi.org/10.1097/01.AOG.0000127035.64602.97>
3. De M, Biswas S, Ganguly RP, Begum SA, Patel S, Haldar S, et al. Impact of increased rate of caesarean section on perinatal outcome: sociolegal evaluation. Journal of the Indian Medical Association. 2011;109(5):312–4, 29.

4. Menacker F, Declercq E, Macdorman MF. Cesarean delivery: background, trends, and epidemiology. *Semin Perinatol*. 2006;30(5):235–41. <http://dx.doi.org/10.1053/j.semperi.2006.07.002>
5. Kottwitz A. Mode of birth and social inequalities in health: the effect of maternal education and access to hospital care on cesarean delivery. *Health Place*. 2014;27:9–21. <http://dx.doi.org/10.1016/j.healthplace.2014.01.005>
6. Sharpe AN, Waring GJ, Rees J, McGarry K, Hinshaw K. Cesarean section at maternal request—the differing views of patients and healthcare professionals: a questionnaire based study. *Eur J Obstet Gynecol Reprod Biol*. 2015;192:54–60. <http://dx.doi.org/10.1016/j.ejogrb.2015.06.014>
7. Badakhsh MH, Seifoddin M, Khodakarami N, Gholami R, Moghimi S. Rise in cesarean section rate over a 30-year period in a public hospital in Tehran, Iran. *Arch Iran Med*. 2012;15(1):4–7.
8. Hendaus MA, Alhammadi AH, Khalifa MS, Muneer E. Does cesarean section pose a risk of respiratory syncytial virus bronchiolitis in infants and children? *Asian Pac J Trop Med*. 2014;7 Supplement 1:S134–6. [http://dx.doi.org/10.1016/S1995-7645\(14\)60219-4](http://dx.doi.org/10.1016/S1995-7645(14)60219-4)
9. Kamath BD, Todd JK, Glazner JE, Lezotte D, Lynch AM. Neonatal outcomes after elective cesarean delivery. *Obstet Gynecol*. 2009;113(6):1231–8. <http://dx.doi.org/10.1097/AOG.0b013e3181a66d57>
10. Maharlouei N, Pourhaghighi A, Shahraki HR, Zohoori D, Lankarani KB. Factors Affecting Exclusive Breastfeeding, Using Adaptive LASSO Regression. *Int J Community Based Nurs Midwifery*. 2018 Jul; 6(3): 260–271.
11. Karlström A, Lindgren H, Hildingsson I. Maternal and infant outcome after caesarean section without recorded medical indication: findings from a Swedish case-control study. *BJOG*. 2013;120(4):479–86, discussion 86. <http://dx.doi.org/10.1111/1471-0528.12129>
12. Liu S, Liston RM, Joseph KS, Heaman M, Sauve R, Kramer MS. Maternal Mortality and Severe Morbidity Associated With Low-Risk Planned Cesarean Delivery Versus Planned Vaginal Delivery at Term. *CMAJ*. 2007;176(4):455–60. <http://dx.doi.org/10.1503/cmaj.060870>
13. Bergenhenegouwen LA, Meertens LJ, Schaaf J, Nijhuis JG, Mol BW, Kok M, et al. Vaginal delivery versus caesarean section in preterm breech delivery: a systematic review. *Eur J Obstet Gynecol Reprod Biol*. 2014;172:1–6. <http://dx.doi.org/10.1016/j.ejogrb.2013.10.017>
14. Jankova I, Dimitrov A. Method of delivery and condition of newborn at 25-30 weeks of gestation. *Akusherstvo i ginekologiya*. 2010;49(6):8–13.
15. Begum B, Zaman RU, Rahman A, Rahaman MS, Uddin MK, Hafiz R, et al. Comparison of risks and benefits of normal vaginal and caesarian section deliveries in a public tertiary hospital in Bangladesh. *Mymensingh medical journal*. MMJ. 2009;18(1) Suppl:S12–4.
16. Fesseha N, Getachew A, Hiluf M, Gebrehiwot Y, Bailey P. A national review of cesarean delivery in Ethiopia. *Int J Gynaecol Obstet*. 2011;115(1):106–11. <http://dx.doi.org/10.1016/j.ijgo.2011.07.011>
17. Pomorski M, Woytoń R, Woytoń P, Kozłowska J, Zimmer M. [Cesarean section versus vaginal delivery--state-of-the-art paper]. *Ginekol Pol*. 2010;81(5):347–51.
18. Maharlouei N, Rezaianzadeh A, Hesami E, Moradi F, Mazloomi E, Joulaei H, et al. The preference of Iranian women to have normal vaginal or cesarean deliveries. *Journal of Research in Medical Sciences : the official journal of Isfahan University of Medical Sciences*. 2013;18(11):943–50.
19. Maharlouei N, Atefi S, Lankarani K. A Study Protocol to Follow a Birth Cohort: Fars Birth Cohort, Shiraz E-Med J. Online ahead of Print ; 20(3):e84308.
20. Arian I, Barut A, Harma M, Harma IM, Gezer S, Ulubasoglu H. Cesarean section with relative indications versus spontaneous vaginal delivery: short-term outcomes of maternofetal health. *Clin Exp Obstet Gynecol*. 2012;39(3):288–92.
21. Haji J, Hamilton JK, Ye C, Swaminathan B, Hanley AJ, Sermer M, et al. Delivery by Caesarean section and infant cardiometabolic status at one year of age. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada*. JOGC. 2014;36(10):864–9.
22. Liston FA, Allen VM, O'Connell CM, Jangaard KA. Neonatal outcomes with caesarean delivery at term. *Arch Dis Child Fetal Neonatal Ed*. 2008;93(3):F176–82. <http://dx.doi.org/10.1136/adc.2006.112565>
23. Hansen AK, Wisborg K, Ulbjerg N, Henriksen TB. Elective caesarean section and respiratory morbidity in the term and near-term neonate. *Acta Obstet Gynecol Scand*. 2007;86(4):389–94. <http://dx.doi.org/10.1080/00016340601159256>
24. Hansen AK, Wisborg K, Ulbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ*. 2008;336(7635):85–7. <http://dx.doi.org/10.1136/bmj.39405.539282.BE>
25. Gerten KA, Coonrod DV, Bay RC, Chambliss LR. Cesarean delivery and respiratory distress syndrome: does labor make a difference? *Am J Obstet Gynecol*. 2005;193(3 Pt 2):1061–4. <http://dx.doi.org/10.1016/j.ajog.2005.05.038>
26. Fogelson NS, Menard MK, Hulsey T, Ebeling M. Neonatal impact of elective repeat cesarean delivery at term: a comment on patient choice cesarean delivery. *Am J Obstet Gynecol*. 2005;192(5):1433–6. <http://dx.doi.org/10.1016/j.ajog.2005.01.010>
27. Huh SY, Rifas-Shiman SL, Zera CA, Edwards JW, Oken E, Weiss ST, et al. Delivery by caesarean section and risk of obesity in pre-school age children: a prospective cohort study. *Arch Dis Child*. 2012;97(7):610–6. <http://dx.doi.org/10.1136/archdischild-2011-301141>
28. Park YH, Kim KW, Choi BS, Jee HM, Sohn MH, Kim KE. Relationship between mode of delivery in childbirth and prevalence of allergic diseases in Korean children. *Allergy Asthma Immunol Res*. 2010;2(1):28–33. <http://dx.doi.org/10.4168/aair.2010.2.1.28>

29. Bodner K, Wierrani F, Grunberger W, Bodner-Adler B. Influence of the mode of delivery on maternal and neonatal outcomes: a comparison between elective cesarean section and planned vaginal delivery in a low-risk obstetric population. *Arch Gynecol Obstet.* 2011;283(6):1193–8. <http://dx.doi.org/10.1007/s00404-010-1525-y>
30. Liu X, Landon MB, Cheng W, Chen Y. Cesarean delivery on maternal request in China: what are the risks and benefits? *Am J Obstet Gynecol.* 2015;212(6):817.e1–9. <http://dx.doi.org/10.1016/j.ajog.2015.01.043>
31. Darmasseelane K, Hyde MJ, Santhakumaran S, Gale C, Modi N. Mode of delivery and offspring body mass index, overweight and obesity in adult life: a systematic review and meta-analysis. *PLoS One.* 2014;9(2):e87896. <http://dx.doi.org/10.1371/journal.pone.0087896>
32. Ahmeti F, Azizi I, Hoxha S, Kulik-Rechberger B, Rechberger T. Mode of delivery and mortality among preterm newborns. *Ginek Pol.* 2010;81(3):203–7.
33. Liu X, Zhang J, Liu Y, Li Y, Li Z. The association between cesarean delivery on maternal request and method of newborn feeding in China. *PLoS One.* 2012;7(5):e37336. <http://dx.doi.org/10.1371/journal.pone.0037336>
34. Ceriani Cernadas JM, Mariani G, Pardo A, Aguirre A, Perez C, Brener P, et al. [Cesarean delivery at term in low risk pregnancies: effects on neonatal morbidity]. *Arch Argent Pediatr.* 2010;108(1):17–2