

Comparison between Australasian triage scale and emergency severity index

Zaynab M. Elsayed^a, Asmaa B. El-Zeny^b, Mohamed S. Moustafa^a, Hany A. Ellouly^a

^aDepartment of Emergency Medicine, Faculty of Medicine, Suez Canal University, Ismailia,

^bDepartment of Emergency Medicine, Port Said General Hospital, Port Said, Egypt

Correspondence to Zaynab M. Elsayed, MD, Emergency Department, Suez Canal University, Ismailia, 1234, Egypt. Tel: +20 122 582 0996; e-mail: zaynab.oof@yahoo.com

Received: 1 January 2020

Accepted: 23 January 2020

Published: 27 April 2020

The Egyptian Journal of Surgery 2020, 39:455–460

Background

The aim of triage system is to ensure that the treatment of patients in the Emergency departments will be according to their clinical urgency. Also, it decreases waiting time and helps in proper direction of the patients to the most suitable area for better assessment and treatment.

Aim of the Work

The purpose of this study is to compare the efficiency of Australasian Triage Scale versus the Emergency Severity Index triage systems.

Methods

Full medical history of 167 Patients arrived at the ED from January till March 2016 were recorded, chief complain, examination, resources number also were recorded and triaged by both systems.

Results

Our results revealed that the most frequent age group was between 20 to 40 years old, while the least frequent age group was ≥ 80 years old. The frequency of male patients was higher than the female patients. ATS and ESI were agreed in urgency level 1. Urgency level 2 and 5 was higher in ESI system while, Urgency level 3 and 4 was higher in ATS. The majority of the patients were discharged, while only 2.4 percent were dead. There were insignificant differences between ATS system and final outcome regarding degree of urgency. While, there was significant over-triage in urgency level 2, while there was significant under-triage in urgency levels 3–5 in comparison to final outcome.

Conclusion

We found that both systems are applicable, but ESI showed some limitation. So we recommended using ATS since it was easier to use and did not show any limitation.

Keywords:

Australasian triage scale, emergency department, emergency severity index, triage

Egyptian J Surgery 39:455–460

© 2020 The Egyptian Journal of Surgery

1110-1121

Introduction

The word triage came from the French verb (trier), it means to sort or pick up [1]. Triage has a basic role in emergency department (ED)s, where many patients are presented at the same time. The triage system aims to ensure that the treatment of the patient will be according to their clinical urgency state [2].

Triage systems in EDs play an important role in patients' categorization regarding their disease severity; it also decides the location and priority of the treatment given. Minimization of the in-hospital mortality and decreasing the time needed to treatment, length of stay, and the resources used are the main targets of any triage system [3].

There are many triage systems, but only five are considered the most effective; they are the systems of choice [4]. Four various five-level triage systems are accepted internationally, which are Australasian triage scale (ATS), Manchester triage system

(MTS), Canadian triage and acuity scale (CTAS), and emergency severity index (ESI) [3,5].

The ATS was developed in Australia. ATS system categorizes the patients into five categories: category 1 (immediately life-threatening conditions) to category 5 (less urgent). The ATS is an effective way for estimating the severity of incoming patients by medical crew in the ED [5,6]. The ATS has an appropriate level of overall reliability in the ED, but it needs more work to reach a perfect agreement [7]. Using ATS, a great variability in the time of triage decisions was detected. Moreover, analysis of variance suggested presence of a difference between triage duration and staff, patients, and environmental variables [8].

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

The development of ESI system occurred in Boston, Massachusetts. It uses only one algorithm ratings from level 1 (the most acutely ill patients) to level 5 (the least resource-intensive patients). The triage staff evaluates the amount of resources required to discharge the patient from the ED in those not meeting criteria of ESI level 1 or 2. The ESI system can be used for both children and adults. The ESI integrates acuity and evaluates resource consumption to decide the priority of treatment [9–11].

Patients and methods

Patients

This prospective observational comparative study was done in the ED of Port Said General Hospital. The study included 167 patients arrived at the ED from January till March 2016. The study was approved by the ethics committee of faculty of medicine at Suez Canal University.

Inclusion and exclusion criteria

All the patients aged above 12 years old were eligible for inclusion in the study. Exclusion of those who aged below 12 years old or left the ED without being examined by a physician.


Methods

Full medical history was recorded with the following patients' data: patient name, mode of arrival, cause of admission, triage ratings by the triage team, urgency classification by the ED physicians, ED resources used, hospital admission rates (including death), and length of stay (LOS) in ED.

The documentation of the triage assessment included the following essential details:

- (1) Date and time of assessment.
- (2) Main presenting problem(s).
- (3) Relevant medical and surgical history.
- (4) Relevant assessment findings.
- (5) Initial triage category allocated.

Figure 1



Age (years) distribution among the studied triage patients.

- (6) Assessment and treatment area allocated.
- (7) Any diagnostic, first aid, or treatment measures initiated.

Statistical analysis

All statistical data were analyzed using SPSS version 20.0 (IBM Corp., Released 2011, IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp.). Differences were compared by χ^2 or Fisher exact tests or *t*-test. *P* value of less than 0.05 was considered significant. Analysis of variance test was followed by logistic regression analysis model of the dependent. Differences in distribution of urgency levels was tested by means of the Friedman's test.


Results

The study included 167 patients arrived at the ED of Port Said General Hospital.

Regarding the age and sex distribution of the patients in our study, the most frequent age group was between 20 and less than 40 years, whereas the least frequent age group was the patients aged greater than or equal to 80 years (Fig. 1). The frequency of male patients was higher than the female patients (Fig. 2).

Table 1 shows the triage characteristics of the studied patients. The mean±SD time of arrival of the studied patients was 15.65±5.73 o'clock, with a range of 1.10 a.m. to 23.45 p.m. Most patients (83.2%) arrived by a private vehicle without an ambulance. The median delay time was 15 min with a range of 0–110 min. Emergency procedures were taking place at resuscitation room in 9.0% of the patients, at trauma section in 13.2% of the patients, and at the general emergency room in 77.8% of the patients.

Figure 2



Sex distribution among the studied triage patients.

Table 1 Triage characteristics of the studied patients

Variables	All triaged patients (n=167) [n (%)]
Time of arrival	
Mean±SD	15.65±5.73
Range	1.10–23.45
Mode of arrival	
Ambulance	28 (16.8)
Private vehicle	139 (83.2)
Delay or waiting time (min)	
Median	15.0
Range	0–110
Triage place	
Resuscitation room	15 (9.0)
Trauma room	22 (13.2)
General emergency room	130 (77.8)
Treatment place	
Resuscitation room	22 (13.2)
Trauma room	8 (4.8)
General emergency room	137 (82.0)

Table 2 Provisional diagnosis of the studied patients

Variables	All triaged patients (n=167) [n (%)]
Disturbed level of consciousness/coma	10 (6.0)
Fall from height/multiple trauma	3 (1.8)
Allergy	1 (0.6)
Chest pain	20 (12.0)
Abdominal pain/vomiting	29 (17.4)
Severe headache	34 (20.4)
Dyspnea	8 (4.8)
Stab wound	1 (0.6)
Minor trauma	25 (15.0)
Immunization	6 (3.6)
History of previous illness	11 (6.6)
Upper respiratory tract infection	10 (6.0)
Eye trauma	4 (2.4)
Suffocation	5 (3.0)

In our study, most patients (20.4%) experienced severe headache, whereas allergy and stab wound were the least frequent, as shown in Table 2.

Table 3 shows the history of co-morbid conditions and risk factors (age >65, cardiac risk factor, and mechanism of injury) among the studied patients. More than half of the patients (53.9%) did not have any history of comorbid chronic conditions. Most studied patients (73.1%) did not have any risk factors. Only 10.2% of the studied patients had three risk factors, 12.0% had two risk factors, and 4.8% had one risk factor.

The number of resources used in our study are listed in Table 4. Approximately 42% of the studied patients did not consume any resources.

Table 5 shows the degree of urgency in ATS and ESI triage systems in the studied patients. ATS and ESI

Table 3 History of comorbid conditions and risk factors among the studied patients

Variables	All triaged patients (n=167) [n (%)]
Comorbid conditions	
No history of comorbid conditions	90 (53.9)
History of comorbid conditions	77 (46.1)
Number of risk factors	
No risk factors	122 (73.1)
One risk factor	8 (4.8)
Two risk factors	20 (12.0)
Three risk factors	17 (10.2)

Table 4 Number of resources used for the studied patients


Variables	All triaged patients (n=167) [n (%)]
Number of resources	
0	70 (41.9)
1	48 (28.7)
2	20 (12.0)
3	21 (12.6)
4	7 (4.2)
5	1 (0.6)

Table 5 Degree of urgency in Australasian triage scale and emergency severity index triage systems in the studied patients (n=167)

Variables	ATS [n (%)]	ESI [n (%)]	χ^2 test	P value
Degree of urgency				
Urgency 1	6 (3.6)	6 (3.6)	0.00	1.00
Urgency 2	35 (21.0)	41 (24.6)	0.61	0.43
Urgency 3	13 (7.8)	12 (7.2)	0.04	0.84
Urgency 4	51 (30.5)	43 (25.7)	0.95	0.33
Urgency 5	62 (37.1)	65 (38.9)	0.11	0.74

ATS, Australasian triage scale; ESI, emergency severity index. *P<0.05, significant.

Figure 3



triage systems were agreed in urgency level 1. Urgency levels 2 and 5 were higher in ESI system in comparison with ATS system. Urgency levels 3 and 4 were higher in ATS system in comparison with ESI system.

Table 6 Degree of urgency according Australasian triage scale system in comparison to final outcome of the studied patients (n=167)

Variables	ATS [n (%)]	Final outcome [n (%)]	χ^2 test	P value
Degree of urgency				
Urgency 1	6 (3.6)	4 (2.4)	0.41	0.52
Urgency 2	35 (21.0)	25 (15.0)	2.03	0.15
Urgency 3–5	126 (75.4)	138 (82.6)	2.60	0.11

ATS, Australasian triage scale. * $P < 0.05$, significant.

Table 7 Degree of urgency according emergency severity index system in comparison to final outcome of the studied patients (n=167)

Variables	ESI [n (%)]	Final outcome [n (%)]	χ^2 test	P value
Degree of urgency				
Urgency 1	6 (3.6)	4 (2.4)	0.41	0.52
Urgency 2	41 (24.6)	25 (15.0)	4.83	0.028*
Urgency 3–5	120 (71.8)	138 (82.6)	5.52	0.019*

ESI, emergency severity index. * $P < 0.05$, significant.

The final outcome of the studied patients is shown in Fig. 3. Overall, 82.6% had good prognosis, whereas 17.4% of them had poor prognosis (ICU admission or died).

Table 6 shows the degree of urgency according to ATS system in comparison with final outcome of the studied patients. There were insignificant differences between ATS system and final outcome regarding degree of urgency.

Table 7 shows the degree of urgency according to ESI system in comparison with final outcome of the studied patients. There was significant overtriage according to ESI system in urgency level 2, whereas there was significant undertriage in urgency levels 3–5 in comparison with the final outcome.

Discussion

There is pervasive and persistent need to develop an effective triage system with today's overcrowding in EDs, rising patient expectations, increasing financial pressure on emergency services, and limitation on staff number [1].

In this study, the mean age of studied patients was 42.86 ± 18.27 years. The most frequent age group was between 20 years and less than 40 years, whereas the least frequent age group was the patients aged greater than or equal to 80 years. The mean age of patients with poor prognosis was significantly higher than the mean age of patients with good.

Our results agreed with similar studies that evaluated a new prognostic tool for in-hospital mortality in nonsurgical ED patients and assessed the predictors of mortality among emergency medical admissions.

These studies showed that the higher the patient's age, the higher the risk of death within 30 days of hospital care after arrival at the ED. The results showed a 5% increase in mortality per year [12,13].

Moreover, Han *et al.* [14] stated that patients in old age groups (above 75 years of age) had a higher risk of death within 30 days after the ED arrival in comparison with younger patients having the same symptoms.

Furthermore, Platts-Mills *et al.* [15] recorded that ESI is not good in predicting which elderly emergency patients will need immediate life-saving intervention, showing a sensitivity and specificity of 42.3 and 99.2%, respectively.

Regarding sex distribution, the frequency of male patients was higher than the female patients (61.1 vs 38.9%, respectively). However, the female patients had higher prevalence of poor prognosis than male patients (58.6 vs 41.4%, respectively).

In similar prospect, Arslanian-Engoren [16] and Jneid *et al.* [17] had detected less improvement in outcomes in female sex in comparison with male sex. This may have occurred owing to underassessment, underdiagnosis, and undertreatment of women in EDs.

Kuhn *et al.* [18] have detected pervasive sex-related influences working against women from triage through to early ED treatment. This may be owing to differences in symptoms reported by women [19–21].

Our data revealed that there were insignificant differences between ATS system and final outcome

regarding degree of urgency ($P>0.05$). There was significant overtriage according to ESI system in urgency level 2, whereas there was significant undertriage in urgency levels 3–5 in comparison with final outcome ($P<0.05$).

In the same direction, Tanabe *et al.* [22] found that when triage urgency levels were compared with the 'real-outcome' urgency level, the ESI had an undertriage rate of 9% and an overtriage rate of 11%.

Using ESI system, most patients deemed by the expert panel to belong in urgency level 1 or 2 were undertriaged. Our result agreed with Storm-Versloot *et al.* [23] who recorded that ESI had the highest percentage of undertriage as compared with MTS and ISS. This variation between the reference standard and the formally structured systems may be owing to the fact that the expert panel of ED physicians knew what happened to the patient. Thus, they may have retrospectively evaluated such patients as being less (or more) urgent than they would have done on using a formally structured system and before knowing the outcome. Some studies found that the ATS improved triage practice. Physicians stated that their assessments of patients were more accurate, and consistent throughout the department. They reported better communication between medical staff and patients and their relatives and high triage accuracy. The overall agreement between ATS graduates and the chart auditor/expert within one level was 99.7% [24,25].

In relation to ATS triage decisions, Considine *et al.* [26] found that 61% of triage decisions were 'expected triage,' with 18% 'overtriage' and 21% 'undertriage decisions.'

In general, it is difficult to predict hospital admission. The predictions of hospital admission, resource use, and length of stay per urgency level in the triage systems in our study were consistent with other research studies [27–29].

Conclusion

- (1) There were insignificant differences between ATS system and final outcome regarding the degree of urgency.
- (2) There was significant overtriage according to ESI system in urgency level 2, whereas there was significant undertriage in urgency levels 3–5 in comparison with final outcome ($P<0.05$).

Recommendations

We found that both systems are applicable, but ESI has some limitation, so we recommended adapting ATS in the hospital ER, as it is easy to use and did not show any limitation as ESI. Moreover, we recommend training hospital staff on that system to improve outcome of ER and prevent further overcrowding.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- 1 Robertson-Steel I. Evolution of triage systems. *Emerg Med J* 2006; 23:154–155.
- 2 Forero R, Nugus P. Australasian College for Emergency Medicine (ACEM) Literature Review on the Australasian Triage Scale (ATS). University of New South Wales – Simpson Centre for Health Services Research. Sydney: South Western Sydney Clinical School Centre for Clinical Governance Research; 2012.
- 3 Weyrich P, Christ M, Celebi N, Riessen R. Triage systems in the emergency department. *Med Klin Intensivmed Notfmed* 2012; 107:67–78.
- 4 Travers DA, Waller AE, Bowling JM, Flowers D, Tintinalli J. Five-level triage system more effective than three-level in tertiary emergency department. *J Emerg Nurs* 2002; 28:395–400.
- 5 Christ M, Grossmann F, Winter D, Bingisser R, Platz E. Modern triage in the emergency department. *Dtsch Arztebl Int* 2010; 107:892–898.
- 6 Yousif K, Bebbington J, Foley B. Impact on patients triage distribution utilizing the Australasian Triage Scale compared with its predecessor the National Triage Scale. *Emerg Med Australas* 2005; 17:429–433.
- 7 Ebrahimi M, Heydari A, Mazlom R, Mirhaghi A. The reliability of the Australasian Triage Scale: a meta-analysis. *World J Emerg Med* 2015; 6:94–99.
- 8 Gerdtz MF, Bucknall TK. Triage nurses' clinical decision making. An observational study of urgency assessment *J Adv Nurs* 2001; 35:550–561.
- 9 Storm-Versloot MN, Ubbink DT, a Choi VC, Luitse JS. Observer agreement of the Manchester Triage System and the Emergency Severity Index: a simulation study. *Emerg Med J* 2009; 26:556–560.
- 10 van Veen M, Moll HA. Reliability and validity of triage systems in pediatric emergency care. *Scand J Trauma Resusc Emerg Med* 2009; 17:38.
- 11 Gilboy N, Tanabe P, Travers D, Rosenau AM. Emergency Severity Index (ESI): a triage tool for emergency department care, version 4. Implementation Handbook. 2012; 2012:12-0014.
- 12 Olsson T, Terent A, Lind L. Rapid Emergency Medicine score: a new prognostic tool for in-hospital mortality in nonsurgical emergency department patients. *J Intern Med* 2004; 255:579–587.
- 13 Goodacre S, Turner J, Nicholl J. Prediction of mortality among emergency medical admissions. *Emerg Med J* 2006; 23:372–375.
- 14 Han JH, Lindsell CJ, Hornung RW, Lewis T, Storrow AB, Hoekstra JW, *et al.* The elder patient with suspected acute coronary syndromes in the emergency department. *Acad Emerg Med* 2007; 14:732–739.
- 15 Platts-Mills TF, Travers D, Biese K, McCall B, Kizer S, LaMantia M, *et al.* Accuracy of the Emergency Severity Index triage instrument for identifying elder emergency department patients receiving an immediate life-saving intervention. *Acad Emerg Med* 2010; 17:238–243.
- 16 Arslanian-Engoren C. Gender and age bias in triage decisions. *J Emerg Nurs* 2000; 26:117–124.
- 17 Jneid H, Fonarow GC, Cannon CP, Hernandez AF, Palacios IF, Maree AO, *et al.* Sex differences in medical care and early death after acute myocardial infarction. *Circulation* 2008; 118:2803–2810.
- 18 Kuhn L, Page K, Davidson PM. Triage women with acute coronary syndrome: a review of the literature. *J Cardiovas Nurs* 2011; 26:395–407.
- 19 Arslanian-Engoren C, Patel A, Fang J, *et al.* Symptoms of men and women presenting with acute coronary syndromes. *Am J Cardiol* 2006; 98:1177–1181.

- 20 DeVon HA, Ryan CJ, Ochs AL, *et al.* Symptoms across the continuum of acute coronary syndromes: differences between women and men. *Am J Crit Care* 2008; 17:14–24.
- 21 Canto JG, Rogers WJ, Goldberg RJ, *et al.* Association of age and sex with myocardial infarction symptom presentation and in-hospital mortality. *JAMA* 2012; 307:813–822.
- 22 Tanabe P, Gimbel R, Yarnold PR, *et al.* Reliability and validity of scores on The Emergency Severity Index version 3. *Acad Emerg Med* 2004; 11:59–65.
- 23 Storm-Versloot MN, Ubbink DT, Kappelhof J, *et al.* Comparison of an informally structured triage system, the emergency severity index, and the Manchester triage system to distinguish patient priority in the emergency department. *Acad Emerg Med* 2011; 18:822–829.
- 24 Atack L, McLean D, LeBlanc L, Luke R. Preparing ED nurses to use the canadian triage and acuity scale with web-based learning. *J Emerg Nurs* 2004; 30:273–274.
- 25 Atack L, Rankin J, Then K. Effectiveness of a 6 week on-line course in the Canadian Triage and Acuity Scale for emergency nurses. *J Emerg Nursing* 2005; 31:436–441.
- 26 Considine J, Le Vasseur SA, Villanueva E. The Australasian Triage Scale: examining emergency department nurses' performance using computer and paper scenarios. *Ann Emerg Med* 2004; 44:516–523.
- 27 Wuerz RC, Travers D, Gilboy N, Eitel DR, Rosenau A, Yazhari R. Implementation and refinement of the emergency severity index. *Acad Emerg Med* 2001; 8:170–176.
- 28 Tanabe P, Travers D, Gilboy N, *et al.* Refining Emergency Severity Index triage criteria. *Acad Emerg Med* 2005; 12:497–501.
- 29 Boeije T, Frederikse MP, van der Heijden FH, Menci F. The Emergency Severity Index (version 3): a good predictor of admission, length of stay and mortality in a European emergency department. *Eur J Emerg Med* 2006;13:62.