

Lichens – masters of extraordinary symbiosis with potent pharmaceuticals

Waill A. Elkhateeb, Ghoson M. Daba

Department of Chemistry of Natural and Microbial Products, Pharmaceutical Industries Researches Division, National Research Center, Dokki, Giza, Egypt

Correspondence to Waill A. Elkhateeb, PhD, Department of Chemistry of Natural and Microbial Products, Pharmaceutical Industries Researches Division, National Research Center, El Buhouth Street, Dokki, Giza 12311, Egypt. Tel: (+20)1013241936; fax: (+20) 233 387 758; e-mail: waillahmed@yahoo.com

Received: 1 March 2020

Revised: 17 March 2020

Accepted: 2 June 2020

Published: 24 August 2020

Egyptian Pharmaceutical Journal 2020, 19:197–201

Lichens are an excellent example of a symbiotic relationship between members belonging to two unrelated separate kingdoms (fungi and algae), which results in the collection of secondary metabolites. These metabolites can be fungal originated, algal originated, or unique compounds not produced by either fungi or algae individually. Although involved since centuries in traditional folk medicine, lichens have attracted extra attention of scientists owing to the emergence of new diseases, which has required screening for novel compounds capable of curing or supporting currently used compounds. This review highlights the nature, importance, nutritional and pharmaceutical uses, and applications of these enigmatic dual organisms.

Keywords:

lichens, lichens substances, nutrition, pharmaceutical drugs, symbiosis

Egypt Pharmaceut J 19:197–201

© 2020 Egyptian Pharmaceutical Journal
1687-4315

Introduction

Lichens are a remarkable example of one of the most important balanced relations in the ecosystem, which is symbiosis. In the case of lichens, this relation defines an association between algae and fungi [1]. This relationship has resulted in the development of a unique organism that is different from its parental individual fungal partner and algal one. Symbiosis fulfils mutual benefit to each member in this relation. One of the simplest exchanged benefits is when the fungal partner derives food from the green algae, whereas the algae partner gets its mineral nutrition and moisture by the help of the fungus [2]. According to the form of the lichens growth, three types were identified, which are crustose, foliose, and fruticose [3,4]. The crustose form is the most simple, and it is noticed on soil, bark, rocks, or wood, whereas foliose and fruticose forms are more complicated, as they show erection or branching [2]. Lichens are capable of growing under different conditions but show sensitivity to many factors such as light, humidity, altitude, and to a greater extent, air quality [5,6]. Uses and applications of lichens are involved in industrial, medicinal, and nutritional fields, and examples are shown in Fig. 1. In this review, the contributions of lichens in traditional medicine have been clarified, and the important metabolites with biological activities originating from lichens are reported. Lastly, some uses and applications of lichens have been described and discussed.

Lichens in ancient medicine


Literatures describing folk medicinal uses of lichens in Europe were started in 1921 [7], whereas the first

global review discussing lichen uses was published in 1997 [8]. Numerous genera have been used from decades in North America, such as *Usnea*, *Letharia*, *Xanthoparmelia*, *Ramalina*, *Lobaria* and *Peltigera*, *Cladonia* and *Cladina*, and *Umbilicaria*. On the contrary, *Evernia* and *Pseudevernia*, *Cetraria*, *Ramalina*, *Lobaria* and *Peltigera*, *Cladonia*, and *Cladina* were commonly used in Europe. Lichens described in traditional medicine of Asian countries included *Usnea*, *Parmotrema* and *Hypotrachyna*, *Cladonia* and *Cladina*, *Thamnoia*, *Ramalina*, *Lobaria* and *Peltigera*, *Umbilicaria*, and *Lethariella*. North African countries also used the genera *Evernia* and *Pseudevernia*, *Xanthoparmelia*, and *Usnea* [9]. As noticed, the genus *Usnea* was commonly used worldwide owing to its medical importance [10]. The genus *Usnea* was identified, and five species in this genus were described in the 18th century under the name lichen, and then they were moved to the genus *Usnea* in 1780 [11]. Nowadays, species under the genus *Usnea* have reached ~350 species [12].

Each lichen genus was used specifically for treating certain disease. *Usnea* sp. is known for its demulcent activities and used to treat mild oral and pharyngeal inflammation. Moreover, *Usnea filipendula* Stirt was previously used for curing cuts and wounds in the former Soviet Union [13]. For stomach diseases, Northern America used *Letharia* [14], whereas

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Figure 1



Some pharmaceutical, nutritional, and industrial applications and uses of lichens.

spleen enlargement was cured in Arabian medicine by using *Alectoria usneoides*. Within Spanish folk medicine, the use of lichens in various medical ailments is also documented [14]. Curing and medicinal activities of lichens mainly rely on lichen substances secreted by that specific lichen [15–17].

Lichen substances

Metabolites produced by lichens are the result of responding to various external conditions as a kind of protection against different physical and biological influences. Lichens' metabolites can be classified into two main groups, primary and secondary metabolites [17]. Intracellular metabolites required for the growth and maintenance of cellular function are known as primary metabolites. Such metabolites can be produced by the fungal partner or the algae partner of the lichen, or unique metabolites produced nonspecifically. Amino acids, proteins, polyols, polysaccharides, vitamins, and carotenoids are considered as primary metabolites [18]. Primary metabolites are soluble in water; hence, they can be easily extracted by hot water. The amount of nitrogenous compounds within lichens ranges between 1.6 and 11.4% dry weight of the thallus, whereas carotenoids exist in the range 1.5–24 mg/g dry weight of the thallus. Examples for those carotenoids are β -carotene epoxide, α -cryptoxanthin, lutein, astaxanthin, and mutatoxanthin. Many vitamins are produced by lichens such as ascorbic acid, pantothenic acid, folic acid, nicotinic acid, riboflavin, biotin, thiamine, and α -tocopherol [18]. In contrast to primary metabolites, secondary metabolites are not

involved in either growth or reproduction of lichens; therefore, secondary metabolites do not participate in the basic molecular skeleton of the microorganism [17,19,20]. Till now, over one thousand metabolites have been originated and identified from lichens [21]. The majority of these metabolites are unique to lichens, whereas a minority exist in other fungi or higher plants. Moreover, the majority of secondary metabolites are poorly soluble in water; hence, their extraction requires using organic solvents [22]. One characteristic feature of lichens' secondary metabolites is their extreme stability [23]. However, some works have described variability in concentrations of metabolites such as usnic acid and atranorin [24]. The production of secondary compounds is genetically controlled and, in some cases, is dependent on geography and morphology of the species or genus [25]. Histologically, lichens' secondary metabolites are commonly existing in the medulla or to a lesser extent in the cortex. The most usual cortical-located metabolites are atranorin and usnic acid. Moreover, anthraquinones, xanthenes, and pulvinic acid derivatives were also reported to be located in the cortex. The expression of lichen metabolites differs among the layers of lichen thallus, and typical cortical metabolites can be distinguished from compounds usually found only in the medulla. The majority of the cortical compounds function as a light filter [26].

Generally, metabolites production, their variation, and concentration are usually taxon specific and depend on many factors, whereas some are synthesized only in permissive physiological stages. Hence, the production of metabolites in axenic cultures can differ significantly from that in nature [18]. For instance, fungal partner (mycobionts) grown without their algal partner (photobionts) produces expected secondary metabolites under certain conditions [27–29] but can also produce different metabolites from those found in symbiosis [30]. In general, different culture conditions are required by every lichen mycobiont to produce specific secondary metabolites. These conditions include specific temperatures, light levels, nutrient medium, pH, the presence of certain sugars or polyols, and stress [29]. According to their chemical structures and biosynthetic origins, lichen secondary metabolites are originated from three chemical pathways: shikimate, acetyl-malonate, and mevalonate [21]. Definitely every metabolite produced by lichens exerts a specific function that assists the lichen itself in one way or another. For example, some phenolic compounds such as melanins are

protectors against excessive radiation through ultraviolet B exposure through being accumulated in the thallus [31]. Such compounds exert potent antioxidant activity [32]. Furthermore, many phenolic compounds protect the lichen from being eaten by herbivorous animals [33], whereas other secondary metabolites display antimicrobial activities that prevent microbial attacks that may lead to degradation of the thallus [34]. Many lichen-synthesized secondary metabolites are responsible for keeping the equilibrium in the symbiotic relation, whereas others may dissolve rocks for improved attachment of the lichen [35,36].

Lichens' nutritional benefits

Owing to their mild toxicity and difficulty in digestion, people have traditionally conducted different preparation methods to make lichens edible, such as removing secondary compounds and/or hydrolyzing the lichen polysaccharides [32]. This can be accomplished by boiling or steaming which would help to hydrolyze the lichen polysaccharides into digestible forms. Many carbohydrates (polyols) exist in lichens such as adonitol (ribitol), mesoerythritol, glycerol, myo-inositol, D-mannitol, siphulitol, and volemitol; moreover, monosaccharides such as arabinose, D-fructose, D-galactose, D-glucose, D-tagatose, D-xylose had been previously reported [37]. Moreover, oligosaccharides and polysaccharides such as D-mannitol, peltigeroside, sucrose, trehalose, umbilicin, isolichenin, lichenin, and pustulan are also listed. Organic acids such as citric acid, glyceric acid, malic acid, oxalic acid, phosphoglyceric acid, and succinic acid are reported from many lichens [38]. On the contrary, some lichen polysaccharides can be extracted in significant yield, such as α -glucans, β -glucans, and galactomannans, which are generally expected to be originated from the fungal partner [37]. The nitrogenous compounds exist in many forms in lichens such as the amines, amino acids, and oligopeptides; examples include ammonia, choline, choline sulfate ester, ethanolamine, methylamine, trimethylamine, alanine, α -aminobutyric acid, arginine, asparagine, aspartic acid, betaine, cystine, phenylalanine, leucine, sarcosine, glutamic acid, glutamine, tryptophan, glycine, isoleucine, tyrosine, lysine, methionine, proline, serine, threonine, valine, and picroroccellin [38]. Carotenoids, chlorophylls, and phycobilins were reported in lichens, and they function as receptors of light energy. Carotenoids have another function as a protective agent that prevents the degradation of chlorophyll by molecular oxygen. The total carotenoid content varies from 23.25 to 123.5 g/g dry weight of the lichen [16]. Furthermore, some

sulfur-containing compounds have been reported from lichens such as dimethyl sulfone [38]. Many vitamins and growth factors were also extracted from lichens such as riboflavin, ascorbic acid, niacin, biotin, folic acid, pantothenic acid, thiamine, and vitamin B12 [37].

Lichens as a pollution indicator

To determine environmental contamination quantitatively, some organisms are used as tools in this process. Such organisms are known as biomonitors [39–43]. Owing to the high sensitivity of lichens to different environmental factors, lichens are strong candidates to have a role as bioindicators and/or biomonitors in two methods: (a) mapping all species existing in a specific area (method A), and (b) through individual sampling of lichen species and measuring pollutants accumulating in the thallus, or by transplanting lichens from an uncontaminated area to a contaminated one, and then morphological changes in the lichens thallus were measured (method B) [44–46]. Air pollution owing to industrial pollutants emitted from vehicles and different sources has an influence on diversity and distribution of lichens, which can be used for the Index of Atmospheric Purity. Mapping all species present in a specific area (method A) is a reliable promising method to monitor air pollution [47]. Examples of the most sensitive lichen genera are *Evernia*, *Candelariella*, *Parmelia*, and *Ochrolechia* [48].

Lichens' biological activities and medicinal uses

Natural products are a promising therapeutic alternative to the currently used antimicrobial treatment [49,50]. Lichen-derived products, which are known as lichen substances, and their antibiotic properties are attracting serious attention nowadays, because up to 50% of all discovered lichens have been reported to exert antibiotic properties [8,51]. The study by Burkholder *et al.* [52] was the first pioneered research describing lichens as antibacterial agents. Some of the most frequently reported lichen-derived metabolites with strong antimicrobial activities are usnic acid, vulpinic acid, and evernic acid, which can successfully inhibit growth of many gram-positive bacteria such as *Staphylococcus aureus*, *Bacillus megaterium*, and *Bacillus subtilis*. Nevertheless, such compounds had no effect on the tested gram-negative bacteria (*Escherichia coli* or *Pseudomonas aeruginosa*) [51,53]. The extracts of lichen species *Parmelia sulcata* were reported to contain salazinic acid, which has exhibited antibacterial properties

against *Aeromonas hydrophila*, *Bacillus cereus*, *Bacillus subtilis*, *Listeria monocytogenes*, *Proteus vulgaris*, *Yersinia enterocolitica*, *S. aureus*, *Streptococcus faecalis*, *Candida albicans*, and *Candida glabrata* [54]. Diethyl ether, acetone, and ethanol extracts of *Cetraria aculeate* contained protolichsterinic acid with promising antibacterial activity against nine bacteria belonging to gram-positive and gram-negative groups [55]. Most of the antibacterial activities were tested specifically on *Bacillus*, *Pseudomonas*, *E. coli*, *S. aureus*, *Klebsiella*, *Candida*, *Salmonella*, and *Yersinia* [56–59]. Among the *Bacillus* species, *Bacillus subtilis* was the most sensitive bacterium to lichen metabolites such as atranorin, protolichsterinic acid, salazinic acid, usnic acid, norstictic acid, protoacetraric acid, fumaroprotoacetraric acid, atranol, lecanoric acid, stictic acid, divericatic acids, and zeorin [60–68].

Conclusion

Understanding the importance of lichens, which are existing everywhere and have prestigious contributions in traditional folk medicine all over the world, is currently of great interest. Moreover, a remarkable list of applications are based on lichens, starting from their use as nutrients and animal feed, moving to their use as a source of dyes, their use in many countries as indicators for air quality and changes in climate, as lichens respond to the smallest change in climate or pollution in a way much faster than most other biological options, and finally ending with the pharmaceutical uses of their substances (metabolites). Further studies on lichens and their metabolites are highly required to get optimum benefits from these valuable dual organism.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Shukla V, Joshi GP, Rawat MS. Lichens as a potential natural source of bioactive compounds: a review. *Phytochem Rev* 2010; 9:303–314.
- Llano GA. Economic uses of lichens. *Econ Bot* 1948; 2:15.
- Clair SB, Clair LL, Weber DJ, Mangelson NF, Eggett DL. Element accumulation patterns in foliose and fruticose lichens from rock and bark substrates in Arizona. *Bryologist* 2002; 105:415–421.
- Monge-Najera J. Relative humidity, temperature, substrate type, and height of terrestrial lichens in a tropical paramo. *Rev Biol Trop* 2019; 67:1.
- Nayaka S. Methods, techniques in collection, preservation and identification of lichens. In: Rana TS, Nair KN, Upreti DK. (editors), *Plant taxonomy and biosystematics: classical and modern methods*. India: New India Publishing Agency; 2014.101–105.
- Ranković B. *Lichen secondary metabolites*. Cham: Springer International Publishing; 2015.
- Smith AL. *Lichens*. London: Cambridge University Press; 1921.
- Sharnoff SD. *Lichens and people*. 1997. Available at: <http://www.lichen.com/people.html>. [Accessed on 5 July 2011].
- Crawford SD. Lichens used in traditional medicine. In *Lichen secondary metabolites 31-97*. Cham: Springer; 2019.
- Crawford S. *Ethnolichenology of Bryoria fremontii: wisdom of elders, population ecology, and nutritional chemistry [MSc thesis]*. Victoria: University of Victoria, Canada 2007.
- Weber GH, Wiggers FH. *Primitiae Florae Holsaticae*. Litteris Mich. Frider. Bartschii Acad. Typogr, Germany. 1780.
- Thell A, Crespo A, Divakar PK, Kärnefelt I, Leavitt SD, Lumbsch HT, Seaward MR. A review of the lichen family Parmeliaceae – history, phylogeny and current taxonomy. *Nord J Bot* 2012; 30:641–664.
- Chevallier A. *The encyclopedia of medicinal plants*. London: Dorling Kindersley; 1996.
- Malhotra S, Subban R, Singh AP. Lichens – role in traditional medicine and drug discovery. *Internet J Alternat Med* 2008; 5:1–5.
- Reddy SM. *University botany I: (algae, fungi, bryophyta and pteridophyta) (Vol. 1)*. New Delhi, India: New Age International; 2001.
- Podterob AP. Chemical composition of lichens and their medical plications. *Pharma Chem J* 2008; 42:582–588.
- Ranković B, Kosanić M. Lichens as a potential source of bioactive secondary metabolites. In *Lichen Secondary Metabolites*. Cham: Springer; 2019. 1–26.
- Hawksworth D. Lichen secondary metabolites: bioactive properties and pharmaceutical potential. In: Ranković B editor. *The Lichenologist*. Cham, Switzerland: Springer International Publishing; 2015. 47:277–278.
- Galun M, Shomer-Ilan A. Secondary metabolic products. In: Galun M (editor) *CRC handbook of lichenology*. Boca Raton, FL: CRC 1988. III:3–8.
- Solhaug K, Lind M, Nybakken L, Gauslaa Y. Possible functional roles of cortical depsides and medullary depsidones in the foliose lichen *Hypogymnia physodes*. *Flora-Morphol Distrib Funct Ecol Plants* 2009; 204:40–48.
- Molnár K, Farkas E. Current results on biological activities of lichen secondary metabolites: a review. *Z Naturforsch C* 2010; 65:157–173.
- Bačkorová M, Jendželovský R, Kello M, Bačkor M, Mikeš J, Fedoročko P. Lichen secondary metabolites are responsible for induction of apoptosis in HT-29 and A2780 human cancer cell lines. *Toxicol In Vitro* 2012; 26:462–468.
- Rundel PW. The ecological role of secondary lichen substances. *Biochem Syst Ecol* 1978; 6:157–170.
- Ravinskaya AP, Vainshtein EA. Effect of some ecological factors on the content of lichen substances. *Ekologiya* 1975; 6:82.
- Zhou QM, Guo SY, Huang MR, Wei JC. A study of the genetic variability of *Rhizoplaca chrysoleuca* using DNA sequences and secondary metabolic substances. *Mycologia* 2006; 98:57–67.
- Marques J. A framework for assessing the vulnerability of schist surfaces to lichen-induced weathering in the Upper Douro region (NE Portugal). *Directores: Rubim Almeida y Graciela Paz*. Universidad: Universidade de Porto. Fecha de lectura; 2013.
- Mattsson JE. Lichen proteins, secondary products and morphology – a review of protein studies in lichens with special emphasis on taxonomy (Proceedings of the Symposia on Bryology and Lichenology at the 15 International Botanical Congress)-(Lichen Substances). *J Hattori Bo Lab* 1994; 76:235–248.
- Fazio AT, Adler MT, Bertoni MD, Sepúlveda CS, Damonte EB, Maier MS. Lichen secondary metabolites from the cultured lichen mycobionts of *Teloschistes chrysophthalmus* and *Ramalina celastri* and their antiviral activities. *Z Naturforsch C* 2007; 62:543–549.
- Hager A, Brunauer G, Türk R, Stocker-Wörgötter E. Production and bioactivity of common lichen metabolites as exemplified by *Heterodea muelleri* (Hampe) Nyl. *J Chem Ecol* 2008; 34:113–120.
- Brunauer G, Hager A, Grube M, Türk R, Stocker-Wörgötter E. Alterations in secondary metabolism of aposymbiotically grown mycobionts of *Xanthoria elegans* and cultured resynthesis stages. *Plant Physiol Biochem* 2007; 45:146–151.
- Gauslaa Y, Solhaug K. Fungal melanins as sun screen for symbiotic green algae in the lichen *Lobaria pulmonaria*. *Oecologia* 2001; 126:462–471.
- Muller K. Pharmaceutically relevant metabolites from lichens. *Appl Microbiol Biotechnol* 2001; 56:9–16.
- Mitrović T, Stamenković S, Cvetković V, Nikolić M, Tošić S, Stojičić D. Lichens as source of versatile bioactive compounds. *Biol Nyssana* 2011; 2:1–6.
- Emmerich R, Giez I, Lange OL, Proksch P. Toxicity and antifeedant activity of lichen compounds against the polyphagous herbivorous insect *Spodoptera littoralis*. *Phytochemistry* 1993; 33:1389–1394.

- 35 Seaward MR. Major impacts made by lichens in biodeterioration processes. *Int Biodeterior Biodegradable* 1997; 40:269–273.
- 36 Huneck S. The significance of lichens and their metabolites. *Naturwissenschaften* 1999; 86:559–570.
- 37 Akbulut G, Yildiz A. An overview to lichens: the nutrient composition of some species. *Kafkas Üni Fen Bilimleri Enstitüsü Dergisi* 2010; 3:2:79–86.
- 38 Culberson CF, Elix JA. Lichen substances. In: Dey PM, Harborne JB (editors) *Methods in plant biochemistry: plant Phenolics*. London: Academic 1989. 509–535.
- 39 Nimis P, Castello M, Perotti M. Lichens as biomonitors of sulphur dioxide pollution in La Spezia (Northern Italy). *Lichenologist* 1990; 22:333–344.
- 40 Gries C. Lichens as indicators of air pollution. *Lichen Biol* 1996; 1:1–29.
- 41 Hamada N, Miyawaki H. Lichens as bioindicators of air pollution. *Jap J Ecol* 1998; 48:49–60.
- 42 Nali C, Balducci E, Frati L, Paoli L, Loppi S, Lorenzini G. Integrated biomonitoring of air quality with plants and lichens: a case study on ambient ozone from central Italy. *Chemosphere* 2007; 67:2169–2176.
- 43 Jovan S. Lichen bioindication of biodiversity, air quality, and climate: baseline results from monitoring in Washington, Oregon, and California, Portland, U.S. Dept. of Agriculture 2008.
- 44 Richardson D. Lichens and man. In: *Frontiers in Mycology*. D. L. Hawkworth, ed. 1991; 187–210
- 45 Nash TH. (Ed.). *Lichenbiology*. Cambridge, United Kingdom: Cambridge University Press; 1996.
- 46 Conti M, Cecchetti G. Biological monitoring: lichens as bioindicators of air pollution assessment—a review. *Environm Poll* 2001; 114:471–492.
- 47 Firdous SS, Naz S, Shaheen H, Dar ME. Lichens as bioindicators of air pollution from vehicular emissions in district Poonch, Azad Jammu and Kashmir, Pakistan. *Pak J Bot* 2017; 49:1801–1810.
- 48 Stamenković S, Cvijan M, Arandjelović M. Lichens as bioindicators of air quality in Dimitrovgrad (South-Eastern Serbia). *Archives of Biological Sciences (Serbia)*. *Arch Biol Sci Belgrade* 2010; 62:643–648.
- 49 Ali MS, Azhar I, Amtul Z, Ahmad VU, Usmanhane K. Antimicrobial screening of some Caesalpiniaceae. *Fitoterapia* 1999; 70:299–304.
- 50 Nimri LF, Meqdam MM, Alkofahi A. Antibacterial activity of Jordanian medicinal plants. *Pharma Biol* 1999; 37:196–201.
- 51 Lawrey JD. Biological role of lichen substances. *Bryologist* 1986; 89:111–122.
- 52 Burkholder PR, Evans AW, McVeigh I, Thornton HK. Antibiotic activity of lichens. *Proc Natl Acad Sci U S A* 1944; 30:250–255.
- 53 Ingolfsdottir K, Gudmundsdottir GF, Ogmundsdottir HM, Paulus K, Haraldsdottir S, Kristinsson H, Bauer R. Effects of tenuiorin and methyl orsellinate from the lichen *Peltigera leucophlebia* on 5-/15-lipoxygenases and proliferation of malignant cell lines in vitro. *Phytomedicine* 2002; 9:654–658.
- 54 Candan M, Yilmaz M, Tay T, Erdem M, Türk AO. Antimicrobial activity of extracts of the lichen *Parmelia sulcata* and its salazinic acid constituent. *Z Naturforsch C* 2007; 62:619–621.
- 55 Türka AO, Yilmaz M, Kivanç M, Türk H. The antimicrobial activity of extracts of the lichen *Cetraria aculeata* and its protolichesterinic acid constituent. *Z Naturforsch C* 2003; 58:850–854.
- 56 Ranković B, Mišić M. The antimicrobial activity of the lichen substances of the lichens *Cladonia furcata*, *Ochrolechia androgyna*, *Parmelia caperata* and *Parmelia conspersa*. *Biotechnol Biotechnol Equip* 2008; 22:1013–1016.
- 57 Karthikaidevi G, Thirumaran G, Manivannan K, Anantharaman P, Kathiresan K, Balasubaramanian T. Screening of the antibacterial properties of lichen *Roccella belangeriana* (awasthi) from Pichavaram mangrove (*Rhizophora* sp.). *Adv Biomed Res* 2009; 3:127–131.
- 58 Karagöz A, Dođruoz N, Zeybek Z, Aslan A. Antibacterial activity of some lichen extracts. *J Med Plants Res* 2009; 3:1034–1039.
- 59 Martins M, Lima M, Silva FP, Azevedo-Ximenes E, Silva N, Pereira EC. *Cladia aggregata* (lichen) from Brazilian northeast: chemical characterization and antimicrobial activity. *Braz Arch Biol Technol* 2010; 53:115–122.
- 60 Manojlovic NT, Vasiljevic PJ, Gritsanapan W, Supabphol R, Manojlovic I. Phytochemical and antioxidant studies of *Laurera benguelensis* growing in Thailand. *Biol Res* 2010; 43:169–176.
- 61 Ranković B, Kosanić M, Sukdolac S. Antimicrobial activity of some lichens and their components. In: Anninos P, Rossi M, Pham TD, Falugi C, Bussing A, Koukhou M, editors. *Recent advances in clinical medicine*. Cambridge, UK: World Scientific and Engineering Academy and Society Press 2010; 279–286
- 62 Santiago KA, Borricano JNC, Canal JN, Marcelo DMA, Perez MCP, dela Cruz TEE. Antibacterial activities of fruticose lichens collected from selected sites in Luzon Island, Philippines. *Phil Sci Lett* 2010; 3:18–29.
- 63 Swathi D, Suchitha Y, Prashith Kekuda TR, Venugopal TM, Vinayaka KS, Mallikarjun N, Raghavendra HL. Antimicrobial, antihelmintic and insecticidal activity of a macrolichen *Everniastrum cirrhatum* (FR.). *Hale Int J Drug Dev Res* 2010; 2:780–789.
- 64 Zambare VP, Zambare AV, Christopher LP. Antioxidant and antibacterial activity of extracts from lichen *Xanthoparmelia somloensis*, native to the black hills, South Dakota, USA. *Int J Med Sci Technol* 2010; 3:46–51.
- 65 Elkhateeb WA, Daba GM. Lichens, an alternative drugs for modern diseases. *Int J Res Pharma Biosci* 2019; 6:5–9.
- 66 Elkhateeb WA, Daba GM, El-Dein AN, Sheir DH, Fayad W, Shaheen MN, et al. Insights into the in vitro hypocholesterolemic, antioxidant, anti-rotavirus, and anti-colon cancer, activities of the methanolic extracts of a Japanese lichen, *Candelariella vitellina*, and a Japanese mushroom, *Ganoderma applanatum*. *Egypt Pharma J* 2020; 19:67–73.
- 67 El-Garawani IM, Elkhateeb WA, Zaghlool GM, Almeer RS, Ahmed EF, Rateb ME, Moneim AE. *A. Candelariella vitellina* extract triggers in vitro and in vivo cell death through induction of apoptosis: a novel anticancer agent. *Food Chem Toxicol* 2019; 127:110–119.
- 68 Kuhnlein HV, Turner NJ. *Traditional plant foods of Canadian indigenous peoples: nutrition, botany and use*. Abingdon, England: Tylor & Francis; 1991.