



## Microwave Assisted Synthesis of Some Azo Disperse Dyes with Antibacterial Activities. Part 1

Morsy A. El-Asasery<sup>1\*</sup>, Abdelhaleem M. Hussein<sup>2</sup>, Abu-Bakr A. M. El-Adasy<sup>2</sup>, Mohamed O. Saleh<sup>2</sup>, Magda M. Kamel<sup>1</sup>

<sup>1</sup>Dyeing, Printing and Auxiliaries Department, Textile Research Division, National Research Centre, Cairo 12622, Egypt.

<sup>2</sup>Department of Chemistry, Faculty of Science, Al Azhar University, Assiut 71524, Egypt.



**T**HIS study aimed to synthesis a series of disperse dyes based on pyrazolopyrimidinones. Enaminones were used to prepare these dyes by reaction in acidic condition by using microwave irradiations. Their structures were established by using Elemental analysis, FTIR, Mass, UV and NMR spectroscopy. The antibacterial activities of these disperse dyes were also evaluated.

### Introduction

The vitality of microwave photons is so low when it contrasted and the connection bond energies and, in this way, microwaves don't influence in a straightway on the molecular structure of a compound and they don't adjust the electronic setup of particles. At the point when materials are irradiated in microwave, they are influenced in an unexpected way. Microwave heating is an elective strategy to conventional heating for quick, compelling and uniform heating [1-4].

Azo disperse dyes are considered as organic colorants that consist of a conjugated chromospheres azo group in association with at least one aromatic or heteroaromatic ring system which have excellent technical properties. In continuation of our interest in synthesis of condensed pyrazolopyrimidinones new dyestuffs [5-9], the current study deals with efficient preparation of condensed disperse dyes by using microwave irradiation as an energy source and studying their antibacterial activities of the synthesized disperse dyes against Gram negative and Gram positive bacteria were investigated.

### Material and Methods

All melting points were determined on an electrothermal digital melting point apparatus and are uncorrected. The infrared spectra were recorded on IR-470 infrared spectrophotometer, Shimadzu; and Pye Unicam SP3-100 spectrophotometer using KBr pellet technique at Assiut University. <sup>1</sup>H-NMR spectra were measured on a Varian 300 MHz in deuterated dimethylsulphoxide (DMSO-d<sub>6</sub>) using tetramethylsilane (TMS) as internal reference and the chemical shifts are expressed in ppm at Cairo University. Mass spectra were performed on HP model MS-5988 at Cairo University. Microanalyses for C, H, N and halogen were performed on a Vario El Elemental analyzer at Cairo University.

*General Procedure for the Synthesis of azo Disperse Dyes 3a-c, 5a-d and 7a,b.*

#### Method A

A mixture of arylhydrazonopyrazolones (4.1, 0.02 M) and enaminones (3.5 g, 0.02 M) was dissolved in glacial acetic acid 5 mL and refluxed for 3 h. the formed solid was collected and crystallized from DMF/water to give dyes **3a-e**.

#### Method B

A mixture of arylhydrazonopyrazolones

\*Corresponding author e-mail: elapaserym@yahoo.com

Received 7/12/2018; Accepted 19/12/2018

DOI: 10.21608/ejchem.2018.6547.1552

©2019 National Information and Documentation Center (NIDOC)

(0.02 M) and enaminones (g, 0.02 M) dissolved in glacial acetic acid 5 mL and irradiated in a microwave oven at power 420 watt for 5 min.

*3-(2-Methoxy-phenylazo)-7-phenyl-pyrazolo[1,5-a]pyrimidin-2-one (3a).*

Yield of method (A) (74.2%), Yield of method (B) (87.8%), m.p. 260-262 °C. IR (KBr):  $\nu$  = 3493  $\text{cm}^{-1}$  (NH), 3035  $\text{cm}^{-1}$  (CH-arom.), 2930  $\text{cm}^{-1}$  (CH-aliph.), 1632  $\text{cm}^{-1}$  (C=O of amid group), 1578  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 425 nm. MS:  $m/z$  = 345 ( $\text{M}^+$ , 65%), 238 (95%), 225 (46%), 182 (100%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 3.96 (s, 3H,  $\text{OCH}_3$ ), 7.47 (d, 2H,  $J$  = 4.8 MHz, arom-H), 7.61 (d, 2H,  $J$  = 5.10 MHz, arom-H), 7.67-7.74 (m, 5H, arom-H), 8.05 (d, 2H,  $J$  = 7.8 MHz, arom-H), 8.73 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. for  $\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}_2$  (345): C, 66.10; H, 4.35; N, 20.30. Found: C 65.91; H 4.24; N 20.19.

*3-(4-Methoxyphenylazo)-7-phenyl-pyrazolo[1,5-a]pyrimidin-2-one (3b).*

Yield of method (A) (84.3%), Yield of method (B) (92.8%), m.p. 222-224 °C. IR (KBr):  $\nu$  = 3493  $\text{cm}^{-1}$  (NH), 3035  $\text{cm}^{-1}$  (CH-arom.), 2930  $\text{cm}^{-1}$  (CH-aliph.), 1632  $\text{cm}^{-1}$  (C=O of amid group), 1578  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 418 nm. MS:  $m/z$  = 345 ( $\text{M}^+$ , 100%), 238 (52%), 210 (25%), 182 (57%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 3.84 (s, 3H,  $\text{OCH}_3$ ), 7.46 (d, 2H,  $J$  = 4.8 MHz, arom-H), 7.54-7.64 (m, 5H, arom-H), 7.79 (d, 2H,  $J$  = 8.7 MHz, arom-H), 8.06 (d, 2H,  $J$  = 8.4, arom-H), 8.73 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. for  $\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}_2$  (345): C, 66.10; H, 4.35; N, 20.30. Found: C 65.90; H 4.30; N 20.24.

*3-(2-Chlorophenylazo)-7-phenyl-pyrazolo[1,5-a]pyrimidin-2-one (3c)*

Yield of method (A) (71 %), Yield of method (B) (90.1%), m.p. 280-282 °C. IR (KBr):  $\nu$  = 3443  $\text{cm}^{-1}$  (NH), 3061  $\text{cm}^{-1}$  (CH-arom.), 1633  $\text{cm}^{-1}$  (C=O of amid group), 1601  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 435 nm. MS:  $m/z$  = 348.5 ( $\text{M}^+$ , 52%), 314 (25%), 238 (100%), 210 (25%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 7.67 (m, 5H, arom-H), 7.91 (d, 2H,  $J$  = 8.70 MHz, arom-H), 8.07 (d, 2H,  $J$  = 4.8 MHz, arom-H), 8.33 (d, 2H,  $J$  = 7.8 MHz, arom-H), 7.78 (d, 1H,  $J$  = 4.2 MHz, NH). Anal. Calcd. for  $\text{C}_{18}\text{H}_{12}\text{N}_5\text{OCl}$  (349.5): C, 61.80; H, 3.43; N, 20.03. Found: C 62.00; H 3.71; N 20.07.

*3-Phenylazo-7-p-tolyl-pyrazolo[1,5-a]pyrimidin-2-one (5a).*

Yield of method (A) (65.7%), Yield of method (B) (86.4%), m.p. 200-202 °C. IR (KBr):  $\nu$  = 3493  $\text{cm}^{-1}$  (NH), 3035  $\text{cm}^{-1}$  (CH-arom.), 2930  $\text{cm}^{-1}$  (CH-aliph.), 1632  $\text{cm}^{-1}$  (C=O of amid group), 1578  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 426 nm. MS:  $m/z$  = 329 ( $\text{M}^+$ , 60%), 252 (59%), 224 (36%), 129 (54%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 2.35 (s, 3H,  $\text{CH}_3$ ), 7.29 (d, 2H,  $J$  = 8.1 MHz, arom-H), 7.45 (d, 2H,  $J$  = 4.8 MHz, arom-H), 7.56-7.68 (m, 5H, arom-H), 8.05 (d, 2H,  $J$  = 8.4 MHz, arom-H), 8.70 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. For Chemical Formula:  $\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}$  (329): C, 69.29; H, 4.59; N, 21.26. Found: C 68.99; H 4.54; N 21.56.

Yield of method (A) (68.6%), Yield of method (B) (89.5%), m.p. 264-266 °C. IR (KBr):  $\nu$  = 3446  $\text{cm}^{-1}$  (NH), 3050  $\text{cm}^{-1}$  (CH-arom.), 2917  $\text{cm}^{-1}$  (CH-aliph.), 1629  $\text{cm}^{-1}$  (C=O of amid group), 1589  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 446 nm. MS:  $m/z$  = 359 ( $\text{M}^+$ , 72%), 252 (100%), 224 (30%), 196 (85%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 2.49 (s, 3H,  $\text{CH}_3$ ), 3.96 (s, 3H,  $\text{OCH}_3$ ), 7.47 (d, 2H,  $J$  = 4.8 MHz, arom-H), 7.61 (d, 2H,  $J$  = 5.10 MHz, arom-H), 7.67-7.74 (m, 5H, arom-H), 8.05 (d, 2H,  $J$  = 7.8 MHz, arom-H), 8.73 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. for  $\text{C}_{20}\text{H}_{17}\text{N}_5\text{O}_2$  (359): C, 66.84; H, 4.74; N, 19.49. Found: C 66.56; H 4.45; N 19.70.

*3-(2-Methoxyphenylazo)-7-p-tolyl-pyrazolo[1,5-a]pyrimidin-2-one (5b).*

Yield of method (A) (64.3%), Yield of method (B) (88.4%), m.p. 215-217 °C. IR (KBr):  $\nu$  = 3424  $\text{cm}^{-1}$  (NH), 3023  $\text{cm}^{-1}$  (CH-arom.), 2960  $\text{cm}^{-1}$  (CH-aliph.), 1632  $\text{cm}^{-1}$  (C=O of amid group), 1589  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 415 nm. MS:  $m/z$  = 359 ( $\text{M}^+$ , 24%), 343 (100%), 252 (95%), 242 (45%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 2.42 (s, 3H,  $\text{CH}_3$ ), 3.84 (s, 3H,  $\text{OCH}_3$ ), 7.07 (d, 2H,  $J$  = 9 MHz, arom-H), 7.42 (d, 2H,  $J$  = 4.5 MHz, arom-H), 7.79-7.86 (m, 4H, arom-H), 8.03 (d, 2H,  $J$  = 8.7 MHz, arom-H), 8.71 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. for  $\text{C}_{20}\text{H}_{17}\text{N}_5\text{O}_2$  (359): C, 66.84; H, 4.77; N, 19.49. Found: C 70.26; H 4.86; N 19.55.

*3-(4-Methoxyphenylazo)-7-p-tolyl-pyrazolo[1,5-a]pyrimidin-2-one (5c).*

Yield of method (A) (58.6%), Yield of method (B) (87.4%), m.p. 246-248 °C. IR (KBr):  $\nu$  = 3446  $\text{cm}^{-1}$  (NH), 3027  $\text{cm}^{-1}$  (CH-arom.), 2965  $\text{cm}^{-1}$  (CH-aliph.), 1629  $\text{cm}^{-1}$  (C=O of amid group), 1586  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 431 nm. MS:  $m/z$  = 363.5 ( $\text{M}^+$ , 60%), 328 (30%), 252 (100%), 224 (34%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 3.35 (s, 3H,  $\text{CH}_3$ ),

*3-(2-Chlorophenylazo)-7-p-tolyl-pyrazolo[1,5-a]pyrimidin-2-one (5d).*

Yield of method (A) (65.7%), Yield of method (B) (86.4%), m.p. 200-202 °C. IR (KBr):  $\nu$  = 3493  $\text{cm}^{-1}$  (NH), 3035  $\text{cm}^{-1}$  (CH-arom.), 2930  $\text{cm}^{-1}$  (CH-aliph.), 1632  $\text{cm}^{-1}$  (C=O of amid group), 1578  $\text{cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}}$  = 426 nm. MS:  $m/z$  = 329 ( $\text{M}^+$ , 60%), 252 (59%), 224 (36%), 129 (54%).  $^1\text{H-NMR}$  (DMSO- $\text{d}_6$ ) 2.35 (s, 3H,  $\text{CH}_3$ ), 7.29 (d, 2H,  $J$  = 8.1 MHz, arom-H), 7.45 (d, 2H,  $J$  = 4.8 MHz, arom-H), 7.56-7.68 (m, 5H, arom-H), 8.05 (d, 2H,  $J$  = 8.4 MHz, arom-H), 8.70 (d, 1H,  $J$  = 4.8 MHz, NH). Anal. Calcd. For Chemical Formula:  $\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}$  (329): C, 69.29; H, 4.59; N, 21.26. Found: C 68.99; H 4.54; N 21.56.

7.41-7.48 (m, 4H, arom-H), 7.54 (d, 2H,  $J=8.7$  MHz, arom-H), 7.78 (d, 2H,  $J=8.7$  MHz, arom-H), 7.99 (d, 2H,  $J=7.8$  MHz, arom-H), 8.70 (d, 1H,  $J=4.8$  MHz, NH). Anal. Calcd. For  $C_{19}H_{14}N_5OCl$  (363.5): C, 62.72; H, 3.86; N, 19.26. Found: C 62.57; H 3.63; N 19.39.

*7-(4-Bromophenyl)-3-(2-methoxyphenylazo)-pyrazolo[1,5-a]pyrimidin-2-one (7a).*

Yield of method (A) (78%), Yield of method (B) (89.8%), m.p. 292-294 °C. IR (KBr):  $\nu = 3446\text{cm}^{-1}$  (NH),  $3050\text{ cm}^{-1}$  (CH-arom.),  $2917\text{ cm}^{-1}$  (CH-alph),  $1629\text{ cm}^{-1}$  (C=O of amid group),  $1589\text{ cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}} = 444\text{ nm}$ . MS:  $m/z = 424$  ( $M^+$ , 31%), 316 (30%), 262 (62%), 235 (14%).  $^1\text{H-NMR}$  (DMSO- $d_6$ ) 3.84 (s, 3H,  $\text{OCH}_3$ ), 7.49 (d, 2H,  $J=4.5$  MHz, arom-H), 7.54 (d, 2H,  $J=8.4$  MHz, arom-H), 7.79-7.85 (m, 4H, arom-H), 8.02 (d, 2H,  $J=8.7$  MHz, arom-H), 8.73 (d, 1H,  $J=4.5$  MHz, NH). Anal. Calcd. for  $C_{19}H_{14}N_5O_2\text{Br}$  (423): C, 53.80; H, 3.30; N, 16.51. Found: C 53.55; H 3.60; N 16.85.

*7-(4-Bromophenyl)-3-(2-chlorophenylazo)-pyrazolo[1,5-a]pyrimidin-2-one (7b).*

Yield of method (A) (65.7%), Yield of method (B) (86.6%), m.p. > 300 °C. IR (KBr):  $\nu = 3442\text{cm}^{-1}$  (NH),  $3063\text{ cm}^{-1}$  (CH-arom.),  $1632\text{cm}^{-1}$  (C=O of amid group),  $1602\text{ cm}^{-1}$  (C=C). UV (DMSO)  $\lambda_{\text{max}} = 446\text{ nm}$ . MS:  $m/z = 472.5$  ( $M^+$ , 51%), 292 (34%), 316 (100%), 262 (75%).  $^1\text{H-NMR}$  (DMSO- $d_6$ ) 7.5 (d, 2H,  $J=4.8$  MHz, arom-H), 7.67-7.75 (m, 4H, arom-H), 7.83 (d, 2H,  $J=8.7$  MHz, arom-H), 8.02 (d, 2H,  $J=8.7$  MHz, arom-H), 8.73 (d, 1H,  $J=4.8$  MHz, NH). Anal. Calcd. For  $C_{18}H_{11}N_5OClBr$  (426.5): C, 50.41; H, 2.57; N, 16.34. Found: C 49.77; H 2.02; N 16.14.

*Antibacterial Activities Test*

All prepared disperse dyes were screened in vitro for their antibacterial activities against *Pseudomonas aeruginosa* as Gram negative bacteria and *Bacillus cereus*, *Micrococcus luteus* and *Staphylococcus aureus* as Gram positive bacteria by the agar diffusion technique [10]. 1 mg/mL solution in dimethylformamide (DMF) was used. The bacteria are maintained on nutrient agar. DMF showed no inhibition zones. The agar media were inoculated with different microorganism's culture tested after 24 hours of inoculated at 37 °C for bacteria. The diameter of inhibition zone (mm) was measured. The data obtained are summarized in Table 1 (These tests conducted at Assiut University).

## Results and Discussion

### *Synthesis and Characteristics*

Our study was based on the preparation of dyes based on microwave technology. In our previous research[1], when using microwave radiation in the preparation of dyes or dye intermediates, we have shown that this technique provides saving of both energy and reaction time, as well as giving better yield. From this point of view aryl-5-Amino-4-(phenyl-hydrazono)-2,4-dihydro-pyrazol-3-one 1a-c react with 3-dimethylamino-1-phenyl-propenone 2 to give the corresponding disperse dyes 3a-c whose structures were elucidated by using elemental analysis, mass spectral data,  $^1\text{H-NMR}$  spectroscopy as well as the IR, and UV (Scheme 1).


By the same way aryl-5-Amino-4-(phenyl-hydrazono)-2,4-dihydro-pyrazol-3-one 1a-d react with 3-dimethylamino-1-p-tolyl-propenone 4 by using microwave irradiation to yield the disperse dyes 5a-d whose identities were proved using elemental analysis, mass spectral data,  $^1\text{H-NMR}$  spectroscopy as well as the IR, and UV (Scheme 2).


Summarily, by using microwave irradiation aryl-5-Amino-4-(phenylhydrazono)-2,4-dihydro-pyrazol-3-one 1a,b react with 1-(4-bromophenyl)-3-dimethylamino-propenone 6 to yield the disperse dyes 7a,b whose structures were conducted using elemental analysis, mass spectral data,  $^1\text{H-NMR}$  spectroscopy as well as the IR, and UV (Scheme 3).


The obtained results revealed that the product yields of microwave irradiation were better than conventional heating and also save time from 3 hrs to 5 min.

### *Antibacterial Activity*

The obtained data that listed in Table 1 showed that the prepared dyes have yielded satisfactory and promising results and can therefore be used for various medical purposes whereas disperse dyes 3a-c, 5a-d and 7a,b showed positive antimicrobial activities against *Pseudomonas aeruginosa*, *Bacillus cereus*, *Micrococcus luteus* and *Staphylococcus aureus*. In contrast dyes 5a, 5d and 7b showed no activities against *Bacillus cereus*. The same phenomena repeated

Scheme 1. Preparation of disperse dyes **3a-c**.

Scheme 2. Preparation of disperse dyes **5a-d**.



Scheme 3. Preparation of disperse dyes 7a,b.

TABLE 1. Antibacterial results of the synthetic disperse dyes.

Dye No	G <sup>-</sup> ( inhibition zone in mm )		G <sup>+</sup> ( inhibition zone in mm )	
	<i>Pseudomonas aeruginosa</i> (cont.42)	<i>Bacillus cereus</i> (cont.18)	<i>Micrococcus luteus</i> (cont.24)	<i>Staphylococcus aureus</i> (cont.34)
3a	10	10	10	15
3b	12	13	12	12
3c	12	14	10	12
5a	12	0	10	10
5b	10	15	0	15
5c	10	15	12	12
5d	15	0	0	10
7a	10	10	0	15
7b	15	0	10	10

with dyes 5b, 5d and 7a showed no activities against of *Micrococcus luteus*, respectively.

### Conclusions

A series of disperse dyes were prepared by using both conventional and microwave heating. The obtained data indicated that the product yields of microwave irradiation were better than conventional heating. Finally, the antimicrobial activities of these dyes against Gram negative and Gram positive bacteria were evaluated.

### References

1. El-Asasery, M.A. Synthesis of some azo disperse dyes by the use of focused microwave heating. *Pol. J. Appl. Chem.* **50**, 75-81 (2006).
2. Al-Qalaf, F.; Almohammad, K.; El-Asasery, M.A.; Mahmoud, H. Synthesis of some biologically active monoazo disperse dyes derived from nicotinic acid derivatives under microwave irradiation for dyeing polyester fabrics. *Eur. J. Chem.* **4**, 211-215 (2013).
3. Al-Etaibi, A.M.; El-Asasery, M.A.; Mahmoud, H.M.; Al-Awadi, N.A. One-pot synthesis of disperse dyes under microwave irradiation: Dyebath reuse in dyeing of polyester fabrics. *Molecules* **17**, 4266-4280 (2012).
4. Al-Etaibi, A.M.; El-Asasery, M.A.; Ibrahim, M.R.; Al-Awadi, N.A. A facile synthesis of new monoazo disperse dyes derived from 4-hydroxyphenylazopyrazole-5-amines: evaluation of microwave assisted dyeing behavior. *Molecules* **17**, 13891-13909 (2012).
5. Tsai, P.C.; Wang, I.J. A facile synthesis of some new pyrazolo[1,5-a]pyrimidine heterocyclic disazo dyes and an evaluation of their solvatochromic behaviour. *Dyes Pigments* **74**, 578-584 (2007).
6. Ho, Y.W. Synthesis of some new azo pyrazolo[1,5-a]pyrimidine-thieno[2,3-b]pyridine derivatives and their application as disperse dyes. *Dyes Pigments* **64**, 223-230 (2005).
7. Rangnekar, D. W. and Puro, S. S. Synthesis of arylazopyrazolo [1,5-a] pyrimidines and their application to synthetic fibres. *Journal of Chemical Technology and Biotechnology*, **49**(3), 275-283 (1990).
8. Al-Etaibi, A.M.; Al-Awadi, N.A.; El-Asasery, M.A.; Ibrahim, M.R. Synthesis of some novel pyrazolo[1,5-a]pyrimidine derivatives and their application as disperse dyes. *Molecules*, **16**, 5182-5193 (2011).
9. Elapasery, A. M., Shakra, S.; Abbas, D.; Gaffer, H.; Allam, A.; Synthesis of some azo disperse dyes based on pyridone moiety and their application on polyester fabrics, *Egyptian Journal of Chemistry*, **60**(6). 97-102 (2006)
10. Isaacson, D.M., Kirschbaum, J. Assays of antimicrobial substances In *Manual of Industrial Microbiology and Biotechnology*; Demain, A.L., Solomon, N.A., Eds; ASM: Washington, DC, USA, (1986).

### تشبيد بعض الصبغات الآزوية المنتشرة بمساعدة الميكروويف مع أنشطتها المضادة للبكتيريا الجزء الأول

مرسى الاباصيرى<sup>1</sup>, عبد الحليم حسين<sup>2</sup>, أبو بكر العداسى<sup>3</sup>, محمد عثمان<sup>4</sup> و ماجدة كامل<sup>5</sup>  
<sup>1</sup>قسم الصباغة و الطباعة و المواد الوسيطة - شعبة بحوث الصناعات النسيجية - المركز القومى للبحوث - الدقى  
 - مصر.  
<sup>2</sup>قسم الكيمياء - كلية العلوم - جامعة الأزهر - اسيوط - مصر.

تم تشبيد فريد للصبغات الآزوية المنتشرة المعتمدة على مشتقات البيرازولوبيريمادينات باستخدام اشعة الميكروويف كطريقة خضراء و آمنة للبيئة و مقارنتها بطرق التحضير العادية و وجد أن استخدام هذه الطريقة موفرة للطاقة و وقت التفاعل كما انها تعطى ناتج تفاعل أعلى من الطرق العادية و تمت ايضا دراسة أنشطة هذه الصبغات المنتشرة ضد البكتيريا و وجد ان لها نشاط جيد.