

Effect of some botanical materials on certain biological aspects of the house fly, *Musca domestica* L.

Nabawy A. I. Elkattan, Khalafalla S. Ahmed, Saadya M. Elbermawy and Rabab M. Abdel-Gawad

Biological and Geological Sciences Department, Faculty of Education, Ain Shams University.

Abstract

The effects of *Lantana camara* (leaves), *Pelargonium zonale* (leaves), *Cupressus macrocarpa* (leaves), *Cyperus rotundus* (whole plant) and *Acacia nilotica* (seeds) powders on some biological aspects of house fly, *M. domestica* L. were tested. The effects of three lethal concentrations LC₂₅, LC₅₀ and LC₇₅ on the larval duration, pupation percent, pupal weight, pupal duration, adult emergence percent, sex ratio, adult longevity, and fecundity were determined. The induced malformed larvae, pupae and adults were recorded and photographed. The powders of the five plants were found to have promising effects in controlling this insect.

Keywords: *Musca domestica*; *Lantana camara*; *Pelargonium zonale*; *Cupressus macrocarpa*; *Cyperus rotundus*; *Acacia nilotica*; Biological studies.

Introduction

The house fly, *Musca domestica* L., is a serious pest to livestock and a public health pest that acts as a transmitter of many human and animal diseases (Emerson *et al.*, 1999; Douglass and Jesse, 2002; Mian *et al.*, 2002).

House fly has been successfully controlled by the application of various insecticides, but reports of insecticide resistance in this insect have been amply found (Kaufman *et al.*, 2001; Shono and Scott, 2004). For this reason, alternative house fly control strategies, including the use of botanical insecticides have been studied (Wang-Jian *et al.*, 2005; Ghoneim *et al.*, 2007; Pavela, 2008; Sripongpun, 2008; Tarelli *et al.*, 2009).

Plants and plant products are recently considered alternatives to conventional insect-control agents as they constitute a rich source of bioactive chemicals, against number of species including specific target insects, and are often biodegradable to non-toxic products (Hashem and Youssef, 1991).

The successful use of plant products in the control of certain insect species depends on contained substances that inhibit the developmental process of those insects (Kristensen and Jespersen, 2003). From these points of view, the aim of this research was to study the effect of some plant materials on the house fly population and the possibility of using these materials as larvicides for controlling the insect by treating insect's breeding places.

Material and methods

Insect rearing

Musca domestica L. colony was obtained from the Medical Insect Research Center, Dokki, Giza. The adults were allowed free access to sugar and cotton pads soaked in milk powder dissolved in water (10% w/v). Larvae were reared according to the method described by Pavela (2008) and Huang *et al.* (2008) on a mixture of sterilized bran (38 g), milk powder (2 g) and water (60 ml), and maintained at 27±2°C and 70±5% relative humidity (RH).

Tested plants

The sublethal concentrations (LC₂₅, LC₅₀ and LC₇₅) of leave's powder of *L. camara*, *P. zonale* and *C. macrocarpa*; whole plant powder of *C. rotundus*, and seed's powder of *A. nilotica* were determined in previous work (Elbermawy *et al.*, 2011).

Biological studies

The experiments were carried out on the 2nd instar larvae (3- days old). The larval media were treated with LC₂₅, LC₅₀ and LC₇₅ of each tested plant. The treated media was divided in 250 ml beakers each received 50g of media. Normal larvae were transferred from rearing media to each beaker (25 larvae). Control experiments were done as above but without any treatment. This procedure was repeated 4 times. All tests were carried out at laboratory conditions mentioned above.

Larvae were examined daily to estimate larval duration which was calculated as the intervals between the commencement of 1st instar larvae and that of pupation. It was calculated for each larva and then the mean value was taken. Mortality was recorded daily until pupation.

The resultant pupae were counted and weighed to determine the percent of pupation and the mean pupal weight. Observations were carried out daily to record pupal duration. The reduction in pupal weight and adult emergence was calculated according to Khazanie (1979). Percentage of total pupae developed to adults was estimated according to Sripongpun (2008) and the emergence of successfully metamorphosed adults was estimated in percentage according to Jimenez Peydro *et al.* (1995).

The emerged males and females adults were transferred daily to oviposition cages, containing sugar and cotton pads served for feeding and oviposition, the cotton pads were renewed daily. Mean longevity for each sex was calculated according to Fletcher *et al.* (1990). The total number of eggs was recorded and the number of eggs laid per female (fecundity) was calculated. Percent fecundity was

determined according to Crystal (1964). The oviposition deterrent index was calculated according to Lundgren (1975). The eggs were moved to Petri dishes containing filter paper moisten by water. Control and treated eggs were incubated under the same laboratory conditions. One day later, the emerged larvae were counted and the percent of egg hatch was determined. The sterility was calculated according to Toppozada *et al.* (1966). Any morphogenetic abnormalities that might occur in all developmental stages were recorded and photographed.

Results

Larval duration:

Results in Table 1 revealed that the larval duration of the control larvae of *M. domestica* was 6.31±0.98 days. A significant prolongation in the larval duration of the treated larvae was observed in larvae treated with LC₂₅ of *C. macrocarpa*. Also, a highly significant prolongation in the larval duration was observed in larvae treated with LC₂₅, LC₅₀ and LC₇₅ of *L. camara*, *P. zonale* and *C. rotundus* and LC₅₀ of *C. macrocarpa*. On the contrary, LC₅₀ of *A. nilotica* caused a highly significant reduction in the average larval period compared to controls. On the other hand, there was insignificant effect on the larval duration after treatment with LC₂₅ of *A. nilotica* and LC₇₅ of *C. macrocarpa* as compared with controls.

Pupation percent, pupal weight and pupal duration:

Results shown in Table 2 revealed that a significant reduction of pupation percent was induced by using LC₂₅ of *P. zonale* and LC₅₀ of *L. camara*. Also, a highly significant decrease in the pupation percent was observed in treatments with LC₇₅ of *L. camara*, LC₅₀ and LC₇₅ of *P. zonale*, all LC's of *C. rotundus* and *C. macrocarpa* and *A. nilotica*. The percent pupation was decreased as the concentration of plant powder increased.

Larvae of *M. domestica* raised on tested plant materials diets recorded a highly significant lower pupal average weight

and the average pupal weights dropped with increased concentration and the effect of plant materials is rated as follows: *P. zonale* > *C. rotundus* > *A. nilotica* > *C. macrocarpa* > *L. camara*. Also, a highly

significant prolongation in the pupal duration was observed in all tested concentration of all plant materials (Table 2).

Table 1: Effect of the tested plant materials on the larval duration of *M. domestica* treated as 2nd larval instar, at 27 °C.

Treatment	Larval duration (days)			Change %	t-Test	
	Min.	Max.	Mean ± SD		P-value	Significance level ⁽¹⁾
Control	5	8	6.31±0.98	-	-	-
<i>L. camara</i>	LC ₂₅	5	10	7.41±1.30	17.37	0.000
	LC ₅₀	6	10	7.47±1.48	18.44	0.000
	LC ₇₅	6	9	7.09±1.02	12.43	0.000
<i>P. zonale</i>	LC ₂₅	7	10	8.41±0.79	33.34	0.000
	LC ₅₀	8	10	9.44±0.62	49.69	0.000
	LC ₇₅	8	10	9.22±0.55	46.20	0.000
<i>C. rotundus</i>	LC ₂₅	6	10	7.45±1.14	18.09	0.000
	LC ₅₀	6	9	7.11±0.87	12.65	0.000
	LC ₇₅	6	8	7.05±0.62	11.80	0.000
<i>C. macrocarpa</i>	LC ₂₅	5	8	6.67±0.81	5.66	0.016
	LC ₅₀	6	8	6.78±0.59	7.46	0.001
	LC ₇₅	6	8	6.57±0.73	4.11	0.099
<i>A. nilotica</i>	LC ₂₅	6	8	6.10±0.38	-3.32	0.074
	LC ₅₀	5	7	5.07±0.30	-19.69	0.000

(1) Significance level: n.s. (insignificant), * (significant), ** (highly significant) as compared with control.

Effect of some....

Table 2: Effect of the tested plant materials on pupation percent, pupal weight and pupal duration of *M. domestica* treated as 2nd larval instar, at 27 °C.

Treatment		% Pupation	% Inhibition in pupation	Pupal weight (mg) (Mean± SD)	% Reduction in pupal weight	Pupal duration (days) (Mean±SD)	% Change in pupal duration
Control		97±3.83	0.00	19.66±2.34		4.51±1.14	
<i>L. camara</i>	LC ₂₅	79±16.77 ^{ns}	15.56	18.53±5.70 **	5.74	7.08±1.66 **	57.23
	LC ₅₀	74±12.44 *	20.71	18.19±4.64 **	7.48	7.50±1.65 **	66.48
	LC ₇₅	64±5.66 **	31.02	17.30±3.84 **	12.02	7.08±1.53 **	57.19
<i>P. zonale</i>	LC ₂₅	80±9.80 *	14.53	9.98±3.65 **	49.26	6.49±1.30 **	44.13
	LC ₅₀	63±9.45 **	32.05	9.51±2.83 **	51.64	6.93±1.67 **	53.73
	LC ₇₅	49±8.25 **	46.48	7.51±2.00 **	61.80	6.40±1.68 **	42.06
<i>C. rotundus</i>	LC ₂₅	71±6.83 **	23.80	10.90±2.49 **	44.55	5.87±1.20 **	30.37
	LC ₅₀	65±3.83 **	29.99	9.91±2.75 **	49.60	6.12±1.25 **	35.86
	LC ₇₅	37±2.00 **	58.86	8.38±1.74 **	57.38	6.61±1.72 **	46.79
<i>C. macrocarpa</i>	LC ₂₅	66±6.93 **	28.96	16.59±3.38 **	15.61	5.91±1.21 **	31.16
	LC ₅₀	59±6.83 **	36.18	14.61±3.27 **	25.69	6.13±1.33 **	36.11
	LC ₇₅	51±12.38 **	44.42	11.82±3.31 **	39.86	5.94±1.45 **	31.95
<i>A. nilotica</i>	LC ₂₅	80±6.53 **	14.53	12.96±3.51 **	34.07	5.34±1.00 **	18.59
	LC ₅₀	75±5.03 **	19.68	10.30±2.69 **	47.60	5.95±1.33 **	32.07

(1) Significance level: n.s. (insignificant), * (significant), ** (highly significant) as compared with control.

Percent adult emergence and adult longevity:

Results shown in Table 3, revealed a reduction in the percent of total pupae developed to adults. All the tested plant powders induced reduction in the percent of adult emerged from treated larvae. Changes in the sex ratios of emerged adults tended towards favoring males. The longevity of adults in both male and female flies was highly significantly decreased in all treatments comparing with the control.

Reproductive potential:

The treatment of *M. domestica* larvae with LC₂₅ of *L. camara* caused a significant decrease in the number of eggs deposited per resulting female. Also, all LC's of *P. zonale*, *C. rotundus*, *A. nilotica* and *C. macrocarpa*, LC₅₀ and LC₇₅ of *L. camara* caused a highly significant decrease in fecundity of adult females. The tested plant powders showed a highly significant decrease in the egg hatching percent (Table 4).

Table 3: Effect of the tested plant materials on the adult emergence percent, sex ratio and adult longevity of *M. domestica* treated as 2nd larval instar, at 27 °C.

Treatment	% of total pupae developed to adults	% Adult emergence	% Inhibition in adult emergence	Sex ratio		Adults longevity (days) (Mean±SD)	
				♂	♀	♂	♀
Control	100.00	97	0.00	49.48	50.52	16.52±1.15	18.10±1.31
<i>L. camara</i>	LC ₂₅	86.08	68	29.90	50.00	50.00	11.44±0.93 **
	LC ₅₀	62.16	46	52.58	54.35	45.65	8.32±1.65 **
	LC ₇₅	40.63	26	73.20	57.69	42.31	7.13±0.92 **
<i>P. zonale</i>	LC ₂₅	83.75	68	30.93	59.70	40.30	9.23±1.62 **
	LC ₅₀	85.71	50	44.33	46.30	53.70	8.12±0.73 **
	LC ₇₅	53.06	27	73.20	57.69	42.31	7.87±0.35 **
<i>C. rotundus</i>	LC ₂₅	95.77	68	29.90	54.41	45.59	6.67±0.84 **
	LC ₅₀	76.92	50	48.45	52.00	48.00	6.46±1.33 **
	LC ₇₅	72.97	27	72.16	51.85	48.15	5.07±0.92 **
<i>C. macrocarpa</i>	LC ₂₅	100.00	66	31.96	48.48	51.52	11.06±1.63 **
	LC ₅₀	86.44	51	47.42	58.82	41.18	10.63±1.10 **
	LC ₇₅	54.90	28	71.13	60.71	39.29	7.29±0.99 **
<i>A. nilotica</i>	LC ₂₅	87.50	70	27.84	65.71	34.29	7.61±1.34 **
	LC ₅₀	70.67	53	45.36	66.04	33.96	6.37±0.81 **
							8.06±0.87 **

(1) Significance level: n.s. (insignificant), * (significant), ** (highly significant) as compared with control.

Effect of some....

Table 4: Effect of the tested plant materials on Fecundity, Fecundity percent, % ODI, Hatchability and Sterility percent of *M. domestica* treated as 2nd larval instar, at 27 °C.


Treatment		Fecundity (no. eggs/female)	% Fecundity	% ODI	Egg hatchability (% egg hatching)	% Sterility
Control		191.35±18.55	100.00	0.00	95.69±0.47	0.00
<i>L. camara</i>	LC ₂₅	164.53±18.28*	85.99	7.53	89.64±2.14**	19.44
	LC ₅₀	88.81±11.96**	46.41	36.60	89.29±3.93**	56.69
	LC ₇₅	50.04±6.45**	26.15	58.54	75.77±5.89**	79.29
<i>P. zonale</i>	LC ₂₅	133.36±14.56**	69.70	17.86	84.69±4.78**	38.31
	LC ₅₀	131.13±19.09**	68.53	18.67	82.36±3.33**	41.01
	LC ₇₅	63.58±7.36**	33.22	50.12	67.26±5.0**	76.64
<i>C. rotundus</i>	LC ₂₅	52.37±6.49**	27.37	57.02	84.23±5.09**	75.91
	LC ₅₀	50.44±6.60**	26.36	58.28	81.46±2.41**	77.56
	LC ₇₅	25.03±3.10**	13.08	76.87	67.89±5.72**	90.72
<i>C. macrocarpa</i>	LC ₂₅	92.89±16.18**	48.54	34.64	90.06±2.74**	54.31
	LC ₅₀	50.76±4.52**	26.53	58.07	79.21±5.09**	78.04
	LC ₇₅	46.22±15.96**	24.15	61.09	65.78±2.82**	83.40
<i>A. nilotica</i>	LC ₂₅	73.22±3.25**	38.26	44.65	76.95±7.26**	69.23
	LC ₅₀	29.76±7.48**	15.56	73.08	76.81±8.68**	87.51

(1) Significance level: n.s. (insignificant), * (significant), ** (highly significant) as compared with control.

Morphogenetic effects:

In the present study, the application of all LC's of *L. camara*, *P. zonale*, *C. rotundus*, *C. macrocarpa* and *A. nilotica* against *M. domestica* induced different morphological abnormalities. Considerable number of larvae, pupae and adults showed obvious malformations after the treatment of 2nd instar larvae with plant powders. Malformations include

complete darkened larvae, curved larvae, irregular-shaped larvae, swelling larvae, larvae with patches of cuticle melanization, larval-pupal intermediates, compressed and shrinkage pupae, dry and darkened pupae, C-shaped pupa, peanut shaped pupa, and small sized pupae. Many adults could not emerge completely and remained concealed in the puparia. Other adults with defective wings, and deformed abdomen were also observed (Plates 1 - 3).



Effect of some....




Plate 2: Normal and treated pupae of *M. domestica*: a, normal pupa; b, *L. camara* treated; c, *P. zonale* treated; d-f, *C. rotundus* treated; g-i, *C. macrocarpa* treated and j-l, *A. nilotica* treated.




Plate 3: Normal and adults resulted from larvae treated with plant materials in *M. domestica*:
 a, normal adults; b-f, *L. camara* treated; g, *P. zonale* treated; h-m, *C. rotundus* treated; n-p, *C. macrocarpa* treated and q-t, *A. nilotica* treated.

Discussion

In the present study, prolongation of the larval duration with tested plants was similar to that reported in *M. domestica* by **Gad-Allah (1991)** using *Melia azedarach* and *Venca rosea*, **Ande (2001)** using *Peganum harmala*, *Acalypha indica* and *Calotropis gigantic*, **Assar (2002 and 2003)** using *Lupinus termis*, *Calotropis procera* and *Atriplex inflate* and **Bakr et al. (2003)** using *Artemisia monosperma*, *Conyza dioscoridis*, *Clerodendron inerme*, *Clocasia antiquorum*. Likewise, white and black mustard lengthened the duration of 2nd larval instars of *M. domestica* (**Abdel Kadder, 2005**). Also, shortened larval period after *A. nilotica* treatment was in accordance with **Shaalan et al. (2005)** in *Aedes aegypti* larvae treated with *Callitris glaucophylla*. They stated that larvae observed to pupate faster as their environment increased in toxicity. This is clearly a self preservation mechanism since the pupal form is less susceptible to the environment.

The percent pupation was decreased as the concentration of plant powder increased. Similar observation was also reported, reduction of percentage of pupation by 91.57%, after treatment of 3rd larval instar of *Synthesiomyia nudiseta* with LC₅₀ of *C. macrocarpa* oil (**Khalaf et al., 2009**). Similar effects of some botanical plant extracts have been reported on *M. domestica* by **Abou El Ela et al. (1995)**; **Ande (2001)**; **Assar (2002 and 2003)** and **Bakr et al. (2003)**.

The decrease of pupal weight in the present study may be attributed to the decrease in total water content or decreased intensity of protein biosynthesis (**Abdel Aal, 1996**). Also, it may be due to the lack of proper sclerotization of the newly formed puparium, or evaporation of body fluids leading to decreased pupal weight. The effect of the tested plant powders on the mean pupal weight of pupae treated as larvae agrees with the results obtained on *M. domestica* by **Kilani et al. (1991)**; **Ande (2001)**; **Assar (2003)** and **Bakr et al. (2003)**.

Prolongation in the pupal duration was observed in all tested concentration of all plant materials. Similar observation was also reported on *M. domestica* by **Assar (2003)** using *A. inflate*, and **Bakr et al. (2003)** using *Artemisia monosperma*, *Conyza dioscoridis*, *Eichhornia crassipes*, *Clerodendron inerme*, *Clocasia antiquorum*, and *Farestia aegyptia*. On the contrary, other studies reported that other plants reduced pupal duration **Bakr et al. (2003)** using *Zygophyllum coccineum* on *M. domestica* and **Khater and Shalaby (2008)** using *Cyperus esculentus* on *C. pipiens*

The decrease in the percentage of adult emergence of *M. domestica* due to treatment with the tested plant materials was similar to the data reported previously by these plants on other dipteran species. The total mean number of males and females of blowfly, *Chrysomya chloropyga* emerging from larvae feeding diet containing 5% of *L. camara* powder, were significantly less than those of the control (**Muse et al., 2003**). High reduction in adult emergence was achieved by larval treatment with *C. macrocarpa* and *A. officinarum* volatile oils against *Synthesiomyia nudiseta* (**Khalaf et al., 2009**).

Disturbance in sex ratio observed after treatment with botanical materials towards more males than females was similar to the data obtained by **Robert and Olson (1989)** they found a change in the sex ratio towards more males in *C. quinquefasciatus* after sub-lethal exposure propoxur and resmethrin. This is not always the case, since **Shaalan et al. (2005)** found a change in the sex ratio towards more females in *Aedes aegypti* after treatment with LC₂₅ of *Callitris glaucophylla*. The shortened adult longevity was also shown in *M. domestica* treated with plant extracts tested by **Gad-Allah (1991)** and **Shoukry (1997)**. On the contrary, the longevity of adult *M. domestica* was not affected by *A. inflate* (**Assar, 2003**) and jojoba oil (**Amer et al., 2004**).

The accumulation of the plant powders in different developmental stages of *M. domestica* might be expected to decrease the longevity of adult flies, as reported in *S. littoralis* after the treatment with *Abrus precatorius* extract (Dimetry and Abdallah, 1991).

The results obtained by the current study indicated that treatment of *M. domestica* larvae with all tested plant powders with all concentrations caused decrease in egg production. Some explanations were introduced by different authors revealing the possible reasons for the reduction of insect fecundity and as a result increasing sterility following the treatment with botanicals insecticides: (i) the weakened physical stage of the treated insects (Tripathi *et al.*, 2003); (ii) mild suppressing effect exerted by the oil on the insect's mating-decisive factor influencing the subsequent number of eggs laid by the insect (Engelmann, 1970); (iii) partial sterilization of females and/or males, or the inability of the sperms to be transferred to the females during copulation (Ismail, 1980); (iv) reduction in the number of normal sperms produced by male insect (El-Meniawi *et al.*, 1999); (v) a blockage in ovarian activity, as the tested botanical products may interfere with oogenesis which, in turn, results in a complete and irreversible sterility of insect female flies (Di Ilio *et al.*, 1999; Khan *et al.*, 2007) and (vi) a delay or reduction of ova giving some opportunities not for retention but for possible egg re-sorption within ovaries. Also, that delay could be due, in part, to a lower metabolic rate (Taher and Cutkomp, 1983; Lucantoni *et al.*, 2006).

Moreover, some extracts from *C. rotundus* prevented the sexual maturity of *S. gregaria* (Bakr *et al.*, 2008). Also, Saxena *et al.* (1992) found that extract of *L. camara* induced oviposition deterrent effect. The extract also had conspicuous activity against the eggs of pulse beetle, *Callosobruchus chinensis* deposited on treated seeds, leading to a pronounced reduction in progeny. As discussed by Weathersbee III and Tang (2002), the

disruption of reproductive capability could lead to substantial population decline over time. Furthermore, Dhar *et al.* (1996) revealed that exposure to neem extract suppressed rather than inhibited oviposition in mosquitoes. Disturbance in sex ratio observed after treatment with *A. nilotica* (ratio males : females was 2: 1) may be the reason for the low number of eggs deposited by females emerged from treated larvae as compared with females emerged from untreated larvae.

Reduction in the egg hatching percent by plant materials was similar to findings reported by many authors using different plant oils and extracts against *M. domestica*, in which the decrease of egg production accompanied with increasing sterility; among these are: *Matricaria chamomilla* and *Clerodendron inerme* (Shoukry, 1997) *Melia azedarach* extract (Radwan, 2000), extracts from leaves and flowers of *Datura innoxia* (Al-Zubaidi *et al.*, 2002) and *A. inflate* (Assar, 2003).

The morphological aberrations induced by plant powders were concentration dependant, in almost cases, the higher concentration the more morphogenetic aberrations. Adamski *et al.* (2005) observed that the degree of malformation was directly proportional to the concentration of pesticides. Our results made also clear co-relation with the recent findings reported from Khalaf *et al.* (2009) where the essential oil of *C. macrocarpa* had been reported to produced clear morphological abnormalities in *S. nudiseta*. Some deformed larvae were pigmented and larval-pupal intermediate, the resultant some individuals showed C-shaped pupae, elongated pupae and balloon shaped pupae, most of the pupae failed to reach adults, however, some emerged adult have various degrees of morphological abnormalities. Topical application of the ethanolic extract from *C. rotundus* onto the penultimate instar nymphs of *S. gregaria* resulted in the formation of defected adults (Bakr *et al.*, 2008). Similar abnormalities were reported by Hashem and Youssef (1991), they

observed dark intersegmental pigments on the 3rd larvae of *M. domestica* and fully formed pupa but with a constricted puparium after treatment the 1st instar larvae with methanolic extraction of leaves and flowers of *M. azedarach*. **Bakr et al.** (2003) found larval pupal intermediate as a result of treatment of *M. domestica* larvae with *A. monosperma*, *C. inerme* and *C. antiquorum*. **El-Domiaty et al.** (2003) found shrinkage of the pupae and folding of the wing of adults as a result of treatment of 3rd instar larvae of *M. domestica* with *P. nigra* volatile oil. **Sripongpun (2008)** observed small sized pupae (1 mm wide x 3 mm long) after treated *M. domestica* larvae with the extract of Chinese star anise fruits, while the size of the control ones was 2 mm wide x 5 mm long. In addition, the number of small pupae developed to adults was less than that of normal one.

Sometimes highly melanized pupae (pupa with darkened puparium) were noticed as a result of treatment. These abnormalities are similar to the effect of IGR's against *M. domestica* as pointed out by **Khalil et al. (2010)**. This indicates that plant powders have also IGR effect.

As a result of treatments, cuticle melanization in patches were observable in *M. domestica* larvae. This phenomenon was observed previously with **Shoukry (1996)** who studied the histopathological effects of Chamomile and Jasmine oils on the house fly larvae. Ultrastructure of muscles of the treated larvae showed that those compounds induced disorganization of light and dark bands of the muscles. This may be the possible explanation for the melanized patches of cuticle or may be due to the inhibition in melanin synthesis (**Gelbič and Němec, 2001**). **El Hadek (2002)** stated that the malformation in prepupal stage, as it appeared as larval-pupal intermediate, may be due to the treated larvae were unable or failed to free themselves from their old cuticle.

In the present study, some of the pupae failed to reach adults. This result was observed previously (**Jahan et al., 1990**), as a number of adults failed to come out from the puparium. Similar results were

obtained by **Naqvi et al. (2007)** using N-9 (extract from neem tree) against *M. domestica*. **Ande (2001)** stated that diet of housefly containing these plant materials no doubt contains desirable primary or secondary principles which may have developed from the interactions of the components of the diet. These principles elicit biological activities in respect of larval/pupal transformation and pupal eclosion hindrances.

Emerging of adults with malformed wings may be attributed to the failure of the wings to expand and flatten after adult emergence (**Saxena et al., 1981**). **Aly et al. (2010)** attributed the adult malformation of *S. gregaria* to the intervening of *F. bruguieri* extracts with the hormonally controlled program of morphogenesis. This may be due to the modification of the ecdysteroid titer, which in turn leads to changes in lysosomal enzyme activity causing overt morphological abnormalities (**Josephrajkumar et al., 1999**).

Conclusion

The results of the present biological studies suggest that the application of plant materials prevented normal development of the different developmental stages of *M. domestica*. Thus, *L. camara*, *P. zonale*, *C. rotundus*, *C. macrocarpa* and *A. nilotica* were nearly comparable with the insect growth regulators (IGR's) in its effects. All are able to reduce larval and pupal weights, adult emergence and the number of laid eggs, shorten adult life span and resulted in larval-pupal intermediates (**Shaurub et al., 1998** and **Naqvi et al., 2007**). According to the current data, *L. camara*, *P. zonale*, *C. rotundus*, *C. macrocarpa* and *A. nilotica* powders are harmful to *M. domestica*, not only reducing longevity of adults but also decreasing their reproductive potential. In conclusion, *L. camara*, *P. zonale*, *C. rotundus*, *C. macrocarpa* and *A. nilotica* show effective IGR-like activities and exhibit great promise in suppressing populations of *M. domestica*.

References

Abdel Kadder A H (2005). Toxicological and histopathological effects of some botanical extracts on the housefly *Musca domestica* L. (Muscidae: Diptera). Ph.D. Thesis, Fac. Sci., Zool. Dept., El- Minia Univ. Egypt.

Abdel-Aal A E (1996). Biological, histological and physiological effects of some insect growth regulators on the greasy cutworm, *Agrotis ipsilon* (Lepidoptera: Noctuidae). M.Sc. Thesis, Fac. of Sci., Cairo Univ. Egypt.

Abou El Ela R G, Helmy N M, El Monairy O M and Salah H (1995). Biological activity of an extract from *Hyoscyamus muticus* on *Musca domestica* (Diptera: Muscidae). Bull. Ent. Soc. Egypt. Econ. Ser., 22: 27- 35.

Adamski Z, Niewadzi M and Ziemnicki K (2005). Inheritance of chorionic malformations and insecticide resistance by *Spodoptera exigua*. J. Appl. Entomol., 129: 526-533.

Aly S A, El Ebarie A S and Hamadah K S (2010). Effects of the wild plant, *Fagonia bruguieri* on the adult performance and phase transition of *schistocerca gregaria* (orthoptera: acrididae). Egypt. Acad. J. Biolog. Sci., 3 (2): 133-147.

Al-Zubaidi F S, Al Rubaie H M and Al Okaily L N (2002). The effects of terpenoid compounds extracted from *Datura innoxia* Mill on the biological performance of *Musca domestica* L. (Diptera: Muscidae). Proc. of the 2nd Inter. Conf. on Biological Sciences, Fac. Sci. Tanta Univ.

Amer M S, Ghoneim K S, Al Dali A G, Bream A S and Hamadah K H (2004). Assessment of the bioactivity of Margosan-o and Jojoba against the house fly *Musca domestica* (Diptera: Muscidae). Al-Azhar Bull.Sci., 15(2): 9-24.

Ande A T (2001). Biological activities of some plant materials against the housefly – *Musca domestica*. Nigerian Soci. for Experimental Biol. J., 1(4): 293-296.

Assar A A (2002). Biological and histopathological effects of the water pant extract of oshaar and termis on the larvae of the house fly, *Musca domestica vicina* Macq. (Diptera: Muscidae). J. Egypt. Ger. Soc. Zool., 38 (E): 33-51.

Assar A A (2003). Biological effects of the water extract of qataf *Atriplex inflate* (Chenopodiaceae) on the house fly, *Musca domestica vicina* Macq. (Diptera: Muscidae). J. Egypt. Ger. Soc. Zool., 41 (E): 15-28.

Bakr M E, Nassef N E, Assar A A, El-Sobky M M and Shams El Din S A (2003). Biological and morphological effects of water extracts of some plants on the house fly, *Musca domestica vicina* Macq. (Diptera: Muscidae). J. Egypt. Ger. Soc. Zool., 41 (E), 29-47.

Bakr R F A, Hussein M A, Hamouda L S, Hassan H A, Elsokary Z F (2008). Effect of some insecticidal agents on some biological aspects and protein patterns of desert locust, *Schistocerca gregaria* (Forskal). Egypt. Acad. Soc. Environ. Develop., 9(2): 29-42.

Crystal M M (1964). Insect fertility: inhibition by folic acid derivatives. Science, New Series, 144 (3616): 308 – 309.

Dhar R, Dawar H, Garg S, Basir S F and Talwar G P (1996). Effect of volatiles from neem and other natural products on gonotrophic cycle and oviposition of *Anopheles stephensi* and *An. Culicifacies* (Diptera: Culicidae). J. Med. Entomol., 33:195–201.

Di Ilio V, Cristofaro M, Marchini D, Nobili P and Dallai R (1999). Effects of a neem compound on the fecundity and longevity of *Ceratitis capitata* (Diptera: Tephritidae). J. Econ. Entomol., 92 (1): 76 – 82.

Dimetry N Z and Abdallah E F (1991). Feeding deterrent effects of various extracts of the bead tree, *Abrus precatorius* seeds towards *Spodoptera littoralis* (Boisd.). Bull. Ent. Soc. Egypt. Econ. Ser., 17: 85 – 92.

Douglass E S and Jesse C (2002). Integrated pest management for fly control in Maine dairy farms. Texas Agricultural Extension Service, 4. 6. pp.

Elbermaway S M, Elkattan N A I, Ahmed K S and Abdel Gawad R M (2011). Screening for bioactive plant materials against house fly, *Musca domestica* L. (Diptera: Muscidae). Functional Plant Science and Biotechnology, 5(1): 45-51.

El Domiaty M M, El Shafae M M, Abdel Aal M M and Rashad E M (2003). Chemical composition and insecticidal activity of *Populus nigra* buds; growing in Egypt. J. Egypt. Acad. Soc. Environ. Develop., 3(2): 21-40.

El Hadek M K M (2002). Phytochemical and toxicological effect of plant extractives against *Spodoptera littoralis* (Biosd.) and *Agrotis ipsilon* (Hufnagel). M.Sc. Thesis, Econ. Entomol. & Pestic. Dept., Fac. Agric., Cairo Univ.

El Meniawi F A, Hashem M, El Mesieri S M and Rawash I A (1999). Physiological effects of neem seed extracts on the reproductive activities of the cotton leafworm, *Spodoptera littoralis* Boisd. Alex. J. Agric. Res., 44(1): 79 – 96.

Emerson P M, Lindsay S W, Walraven G E L, Faal H, Bogh C, Lowe K and Bailey R L (1999). Effect of fly control on trachoma and diarrhoea. Lancet 353: 1401-1403.

Engelmann F (1970): The physiology of insect reproduction. Pergamon Press, New York.

Fletcher M G, Axtell R C and Stinner R E (1990): Longevity and fecundity of *Musca domestica* (Diptera: Muscidae) as a function of temperature. *J. Med. Entomol.*, 27(5): 922-926.

Gad Allah S Z (1991): The effect of some plant extracts on *Musca domestica* vicina. M. Sc. Thesis Fac. of sci., Cairo Univ. Egypt.

Gelbič I and Němec V (2001): Developmental changes caused by metyrapone and azadirachtin in *Spodoptera littoralis* (Boisd.) (Lep., Noctuidae) and *Galleria mellonella* (L.) (Lep., Pyralidae). *J. Appl. Entomol.*, 125: 417-422.

Ghoneim K S, Amer M S, Bream A S, Al Dali A G and Hamadah K S (2007): Effectiveness of Margosan-O and Jojoba on some reproductive aspects of the house fly, *Musca domestica* (Diptera: Muscidae). *Inter. J. Agric. Biol. Friends Sci. Publishers*, Faisalabad, Pakistan, 9 (2); 338-341.

Hashem H O and Youssef N S (1991): Developmental changes induced by methanolic extracts of leaves and fruits of *Melia azedarach* L. on the house fly, *Musca domestica vicina* Macq. *J. Egypt Ger. Soc. Zool.*, 3 (E): 35-52.

Huang Q, Liu M, Feng J and Liu Y (2008): Effect of dietary benzoxadiazole on larval development, cuticle enzyme and antioxidant defense system in housefly (*Musca domestica* L.). *Pestl. Biochem. and Physiol.*, 90: 119-125.

Ismail I E (1980): Physiological studies on the effect of juvenile hormone analogues upon the cotton leafworm *Spodoptera littoralis* Boisd (Lep., Noctuidae). Ph.D. Thesis, Cairo, Univ. Egypt.

Jahan M, Ahmad I and Naqvi S N H (1990): Toxic teratogenic effects of Juliflorine and Margosan-OTM on the *Musca domestica* L., larvae. *Proc. Pak. Congr. Zool.* 10: 293-299.

Jimenez Peydro R, Gimeno Martos C, Lopez Ferror J, Serrano Delgado C and Moreno Mari J (1995): Effects of the insect growth regulator, cyromazine, on the fecundity, fertility and offspring development of Mediterranean fruit fly, *Ceratitis capitata* Wied (Diptera: Tephritidae). *J. Appl. Entomol.*, 119: 435-438.

Josephrajkumar A, Subrahmanyam B and Srinivasan (1999): Plumbagin and azadirachtin deplete haemolymph ecdysteroid levels and alter the activity profiles of two lysosomal enzymes in the fat body of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Eur. J. Entomol.*, 96: 347-353.

Kaufman P E, Scott J G and Rutz D A (2001): Monitoring insecticide resistance in house flies (Diptera: Muscidae) from New York dairies. *Pest Management Science* 57: 514-521.

Khalaf A A, Hussein K T and Shoukry K K (2009): Biocidal activity of two botanical volatile oils against the larvae of *Synthesiomyia nudiseta* (Wulp) (Diptera: Muscidae). *Egypt. Acad. J. Biolog. Sci.*, 2 (1): 89-101.

Khalil M S, Assar A A, Abo El Mahasen M M and Mahmoud S H (2010): Morphological effects of some insect growth regulators on *Musca domestica*, (Diptera, Muscidae). *Egypt. Acad. J. biolog. Sci.*, 2 (2): 29- 36.

Khan M, Hossain M A and Islam M S (2007): Effects of neem leaf dust and a commercial formulation of a neem compound on the longevity, fecundity and ovarian development of the melon fly, *Bactrocera cucurbitae* (Coquillett) and the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pak. J. Biol. Sci.*, 10: 3656-3661.

Khater H F and Shalaby A A (2008): Potential of biologically active plant oils to control mosquito larvae (*Culex pipiens*, Diptera: Culicidae) from an Egyptian locality. *Rev. Inst. Med. trop. S. Paulo*, 50(2):107-112.

Khazanie R (1979): Elementary Statistics. Good Year Publishing Co., California, U.S.A.

Kilani I M, El Dib M A and El Fishawi A A (1991): The effect of aqueous neem, *Azadirachta indica* A.Juss, seed kernel extract on the house fly, *Musca domestica* L. *Egypt J. Biological Pest Control*, 1(1):121-128.

Kristensen M and Jespersen J B (2003): Larvicide resistance in *Musca domestica* (Diptera: Muscidae) populations in Denmark and establishment of resistant laboratory Strains. *J. Econ. Entomol.*, 96(4): 1300-1306.

Lucantoni L, Giusti F, Cristofaro M, Pasqualini L, Esposito F, Lupetti P and Habluetzel A (2006): Effects of a neem extract on blood feeding, oviposition and oocyte ultrastructure in *Anopheles stephensi* Liston (Diptera: Culicidae). *Tissue Cell* 38: 361-371.

Lundgren L (1975): Natural plant chemicals acting as oviposition deterrent on cabbage butter flies, *Pieris rapae* (L.) and *P. napi* (L.). *Zool. Ser.*, 4: 250-258.

Mian L S, Maag H and Tacal J V (2002): Isolation of *Salmonella* from muscoid flies at commercial animal establishments in San Bernardino County, California. *J. Vector Ecol.* 27: 82-85.

Muse W A, Lajide L and Adediri C O (2003): Effect of some nigerian plants on survival, oviposition, and emergence of adult blowfly, *Chrysomya chloropyga* (Wied.)

(Diptera: Calliphoridae). *J. Asia-Pacific Entomol.*, 6 (1): 69-75.

Naqvi S N H, Tabassum R, Khan M F, Yasmin N, Nurulain S M and Burney A A (2007). Toxic, residual, and teratomorphic effect of a neem extract (N-9) in comparison to Coopex 25 WP (Permethrin + Bioallethrin) against *Musca domestica* L. (Holland strain). *Turk. J. Zool.*, 31:127-130.

Pavela R (2008). Insecticidal properties of several essential oils on the house fly (*Musca domestica* L.). *Phytotherapy Research* 22 (2): 274-278.

Radwan M E H (2000). Effect of crude methanolic extracts of leaves and fruits of chinaberry tree *Melia azedarach* L. on the adult female housefly *Musca domestica vicina* Macq. M.Sc. Thesis, Fac. Sci. Alex. Univ.

Robert L L and Olson J K (1989). Effects of sublethal dosages of insecticides on *Culex quinquefasciatus*. *J. Am. Mosq. Contr. Assoc.* 5: 239-246.

Saxena R C, Dixit O P and Harshan V (1992). Insecticidal action of *Lantana camara* against *Callosobruchus chinensis* (Coleoptera: Bruchidae). *J. of Stored Prod. Res.*, 28 (4): 279-281.

Saxena R C, Waldbauer G P, Liquido N J and Puma B C (1981). Effect of neem oil on the leaf-folder, *Cnaphlocrosis medinalis*. Proc. 1st Intn. Neem Conf. Rottach-Egern (1980): 189-203.

Shaalan E A S, Canyon D V, Younes M W F, Abdel Wahab H and Mansour A H (2005). Effects of sub-lethal concentrations of synthetic insecticides and *Callitris glaucophylla* extracts on the development of *Aedes aegypti*. *J. Vector Ecol.*, 30(2): 295-298.

Shaurub E H, Ahmed Z A and El Naggar S E M (1998). Impacts of pyriproxyfen and extracts of *Schinus terebinthifolius* Raddi on development, reproduction and reproductive organs in *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *J. Egypt. Ger. Soc. Zool.*, 27 (E): 57 – 82.

Shono L Z and Scott J G (2004). Indoxacarb resistance in the house fly, *Musca domestica*. *Pestic. Biochem. Physiol.*, 80: 106–112.

Shoukry I F I (1996). Morphogenic and histopathological effects induced by two natural volatile oils in the house fly, *Musca domestica* L. (Diptera: Muscidae). Proc. 1st Conf. on the role of science in the development of Egypt. soc. and environ. Benha branch, Zag. Univ. 21 – 23 October, 1-12.

Shoukry I F I (1997). Delayed effects of the volatile oils of *Matricaria chamomilla* and *Clerodendron inerme* on fecundity, fertility and ovarian amino acids of the female *Musca domestica* L. *Bull. Ent. Soc. Egypt, Econ. Ser.*; 24: 52-60.

Sripongpun G (2008). Contact toxicity of the crude extract of Chinese star anise fruits to house fly larvae and their development. *Songklanakarin J. Sci. Technol.*, 30 (5): 667-672.

Taher M and Cutkomp K. (1983). Effect of sublethal doses of DDT and three other insecticides on *Tribolium confusum*. *J. Stored Prod. Res.*, 19:43-50.

Tarelli G, Zerba E N and Alzogaray R A (2009). Toxicity to vapor exposure and topical application of essential oils and monoterpenes on *Musca domestica* (Diptera: Muscidae). *J. Econ. Entomol.*, 102 (3): 1383-1388.

Toppozada A, Abd Allah S and El Defrawi M E (1966). Chemosterilization of larvae and adult of the Egyptian cotton leafworm, *Spodoptera littoralis* by apholate-metapa and tepea. *J. Econ. Entomol.*; 59 : 1125-1128.

Tripathi A K, Prajapati V, Khanuja S P S and Kumar S (2003). Effect of d-limonene on three stored-product beetles. *J. Econ. Entomol.*; 96(3): 990-995.

Wang-Jian, Li-ya and Lei Chaoliang (2005). The repellency and fumigant activity of *Artemisia vulgaris* essential oil to *Musca domestica vicina*. *Chinese. Bull. Entomol.*, 42(1): 51-53.

Weathersbee III A A and Tang Y Q (2002). Effect of neem extract on feeding, growth, survival, and reproduction of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). *J. Econ. Entomol.*, 95: 661-667.

تأثير عدد من المواد النباتية على بعض النواحي البيولوجية للذبابة المنزلية، مسماً دوميسكا ل.

نبوى عبد الرحمن القطنان، خلف الله صابر أحمد، سعدية محمد البرماوى، رباب مجدى عبد الجاد
قسم العلوم البيولوجية والجيولوجية - كلية التربية - جامعة عين شمس - القاهرة

تم اختبار تأثير مطحون أوراق نباتات اللانتانا والبلارجونيوم والسرور الليمونى ومطحون نبات السعد ومطحون بنور السنط على النواحي البيولوجية للذبابة المنزلية مسماً دوميسكا. وقد تمت دراسة تأثير ثلاثة تركيزات مميتة (LC₂₅, LC₅₀, LC₇₅) على طول عمر اليرقات، نسبة التعذر، وزن العذارى، طول عمر العذارى، نسبة خروج الحشرات البالغة، نسبة الإناث للذكور، طول عمر الطور البالغ، نسبة الخصوبة. كما تمت دراسة وتصوير التشوهات الناتجة فى كل من اليرقات والعذارى والذبابة البالغة بعد المعاملة. وقد أثبتت الدراسة أن مطحون النباتات الخمسة له تأثير واضح على كل النواحي البيولوجية مما يضيف طريقة جديدة آمنة إلى طرق مقاومة هذه الحشرة.