

# Bacterial pathogens and health problems of *Acipenser gueldenstaedtii* and *Acipenser baerii* sturgeons reared in the eastern Black Sea region of Turkey

Kayış, Ş.<sup>1\*</sup>; Er, A.<sup>1</sup>; Kangel, P.<sup>2</sup> and Kurtoğlu, İ. Z.<sup>1</sup>

<sup>1</sup>Department of Aquaculture, Faculty of Fisheries Sciences, Recep Tayyip Erdoğan University, 53100 Rize, Turkey; <sup>2</sup>Post-Graduate, Republic of Turkey Ministry of Food and Agriculture and Livestock of Aquaculture Ordu, Turkey

\*Correspondence: Ş. Kayış, Department of Aquaculture, Faculty of Fisheries Sciences, Recep Tayyip Erdoğan University, 53100 Rize, Turkey. E-mail: aquasevki@msn.com

(Received 22 Jun 2016; revised version 1 Sept 2016; accepted 24 Sept 2016)

## Summary

Farming sturgeons is an economically important practice in a number of Asian and European countries. However, since it is not widely implemented in Turkey, fertilized eggs necessary for research and industrial activities are imported from Germany. Due to the interest of several fish farms in culturing sturgeon in Turkey and the lack of relevant data, this study investigated bacteria related health problems of two different sturgeon species, the diamond sturgeon (*Acipenser gueldenstaedtii*) and the Siberian sturgeon (*Acipenser baerii*). The fungal, parasitic and bacterial pathogens found in these fish were investigated until the fish reached about 3 kg of weight (3+ years). A number of bacterial disease pathogens (*Acinetobacter radioresistens*, some *Aeromonas* and *Pseudomonas* species and *Bacillus mycoides*) and parasite *Trichodina* sp. and fungus *Saprolegnia* sp. were identified in the fish. Both phenotypic and molecular characterizations of the isolated bacteria were performed. Furthermore, swim bladder and spinal problems, cannibalism, tumor growth and mechanical injuries on the external surface of the fish were observed during the study period.

**Key words:** Bacteria, Health problems, Sturgeons, Turkey

## Introduction

Sturgeons are valuable because of their meat and caviar. However, they take longer to mature and are more difficult to culture compared to other fish species (Chebanov and Galich, 2013). The production of feed for this species may also be considered as another limiting factor in its culture. It is believed that sturgeons are comparatively more resistant to fish diseases; nevertheless, many studies have shown that their diseases are idiopathic and involve different pathogens.

The most common bacterial pathogen associated with sturgeons (*Huso huso* and *Acipenser gueldenstaedtii*) held in a marine environment is *Vibrio anguillarum* (Austin and Austin, 2007). The authors of this study believe that *V. anguillarum* infections increase fish intestinal and peritoneal bleeding. Other reports state that *V. alginolyticus* and *Pasteurella* sp. cause mortality in *Acipenser baerii* (Costinar *et al.*, 2010). Apart from the mobile *Aeromonas* sp., *Aeromonas salmonicida* has been reported to cause a disease that manifests itself as furuncles on the skin of *Acipenser oxyrinchus oxyrinchus* (Mohler, 2003). Yersiniosis, or enteric redmouth disease, is frequently seen in aquaculture units and caused by *Yersinia ruckeri*, which has been isolated from *A. baerii* in rare cases (Noga, 2010). *Streptococcus dysgalactiae*, a Gram-positive bacterium, has been isolated from *Acipenser schrenckii* in China. Hemorrhages, abdominal swelling and ascites have also been reported in cultured sturgeons (Yang and Li, 2009). *Flavobacterium johnsonae* was isolated from cultured sturgeons

(*Acipenser gueldenstaedtii*) (Bauer *et al.*, 2002) in Russia, and *Aeromonas hydrophila* and *Pseudomonas fluorescence* were isolated from adult *A. gueldenstaedtii*. These bacteria are known to cause serious conditions such as hemorrhage, exophthalmus and ulcers, ultimately leading to the death of the cultured fish (Prearo *et al.*, 2009; Noga, 2010).

Studies on parasites in sturgeons show that *Trichodina reticulate*, a ciliated parasite, and *Diplostomum spathaceum*, a digenean parasite, are present in reared *Acipenser persicus* larvae (Bazari *et al.*, 2010). The authors of this report mention that these parasites were isolated from fish specially reared and released in their habitat. They also found out that these parasites were isolated from various sturgeon species in their natural habitat in Russia (Bauer *et al.*, 2002).

Fungal diseases of sturgeons listed in a handbook on white sturgeon aquaculture (*Acipenser transmontanus*) include *Saprolegnia* sp. as infectious agents, especially in fingerling and spawning adults (Conte, 1988). Jalilpoor *et al.* (2006) also isolated *Penicillium* spp., *Fusarium* spp., *Mucor* spp. and *Saprolegnia* sp. from the eggs of *A. persicus*, and found that the mortality rate was highest (7-22%) during *Saprolegnia* sp. infestation. Three distinct viral diseases, white sturgeon iridovirus (WSIV), white sturgeon herpesvirus-1,2 (WSHV-1, 2) and shovelnose sturgeon iridovirus (SSIV) (Yu-Ping and Di, 2005), are known to occur in sturgeons.

Sturgeon aquaculture is underdeveloped in Turkey, therefore, limited information is available on their pathogen profile, especially when compared to rainbow

trout (*Oncorhynchus mykiss*), European sea bass (*Dicentrarchus labrax*) and seabream (*Sparus aurata*) aquaculture. The pathogens isolated from sturgeons in Turkey are, in fact, isolated from eggs imported from other countries for scientific studies, and individual fish raised from these imported eggs. So far, only bacterial pathogens have been isolated from sturgeons in Turkey and include *F. johnsonae* (Karataş *et al.*, 2010), *A. hydrophila*, and *Flavobacterium hydatis*, all isolated from *A. gueldenstaedtii* (Timur *et al.*, 2010). Another study investigated deformations during larval development and early feeding stages of the Siberian sturgeon (*Acipenser baerii*) (Kurtoglu *et al.*, 2015).

Despite the live natural stock of sturgeons of the Black Sea coastal line, sturgeon culture in Turkey is underdeveloped and scientific data is insufficient. However, in the last few decades, several studies on sturgeon culture have been conducted and aquaculture enterprises have shown more interest in farming this fish. The present study focused on the bacteriological, fungal and parasitological pathogens associated with two different species of sturgeons, namely, the diamond sturgeon (*Acipenser gueldenstaedtii*) and the Siberian sturgeon (*Acipenser baerii*), reared in the North Eastern part of Turkey.

## Materials and Methods

Fertilized eggs of the diamond sturgeon (*Acipenser gueldenstaedtii*) and the Siberian sturgeon (*Acipenser baerii*) were imported from Germany in 2011 as part of the Recep Tayyip Erdoğan University (RTEU) aquaculture research activity. These eggs were incubated in the RTEU Aquaculture Research and Application Center and were periodically inspected. Five fish were sampled from each species monthly for health problems until they reached an average weight of 3 kg  $\pm$  0.25. The study was conducted between July 2011 and May 2014 and all health problems observed during this period were carefully recorded. Fish were fed with appropriate commercial trout feed for different sizes. Quality parameters of the water used in the present study are presented in Table 1.

**Table 1:** Water quality parameters

Parameters	Spring water for incubation	Rearing water
Temperature (°C)	11	4-17
pH	7.2	6.5-7.5
Dissolved oxygen (mg/L)	11.4	7-11.1
Ammonia (un-ionized) (mg/L)	>0.01	>0.01
Nitrite (mg/L)	>0.01	>0.01
Alkalinity (mg/L as CaCO <sub>3</sub> )	13	50
Total hardness (mg/L as CaCO <sub>3</sub> )	25	35

## Fungal and parasitic examinations

All fungal infections observed in the eggs were examined microscopically, at a low magnification of  $\times 4$ . Samples taken to identify parasitic infections were prepared using scrapings from gills and the skin and

stool samples. These samples were examined microscopically at a magnification of  $\times 40$  (Timur and Timur, 2003) and the parasites were identified based on morphological criteria (Lom and Dykova, 1992). The parasites isolated from the fish were fixed and stained by ethyl alcohol (10%) and crystal violet used in Gram-stain (one drop/10 cc).

## Bacteriological examinations

In order to eliminate non-target and contaminant bacteria, the outer surfaces of normal fish and diseased fish affected by gas problems, external injuries and exophthalmia were cleared for the presence of pathogenic bacteria and cleaned with 70% alcohol. After cleaning, liver, kidney and spleen samples of the fish (5 g-3 kg) were transferred to Tryptic Soy Agar (TSA) under aseptic conditions. The bacterial colonies obtained from these samples were found to be pure colonies and were preserved in media containing 15% glycerol at  $-70^{\circ}\text{C}$  for future use (Lasee, 1995).

Before subjecting the isolates to phenotypic tests, certain pre-tests, such as cytochrome oxidase, Gram-staining, catalase, and motility tests, were conducted. The Analytical Profile Index (API) test (API 20E and API 20NE) was also performed on all bacteria. Glutamate Starch Phenol Red (GSP) selective agar was used to distinguish between the *Aeromonas* and *Pseudomonas* species. Blood agar hemolysis was used to identify the *Bacillus* species, and the bacteria were classified as  $\beta$ -hemolytics or non-hemolytics. Rod-shaped Gram-stained bacteria that showed the presence of an endospore were subjected to motion tests as necessary. The susceptibility to penicillin was also taken into consideration to characterize the bacterial isolates.

For the purpose of identifying bacterial species accurately, DNA obtained from the bacteria was subjected to a universal PCR reaction and the PCR product was sequenced. Bacterial DNA was isolated as follows. Twenty-four-hour fresh bacterial cultures in Tryptic Soy Broth (TSB) medium were centrifuged in 0.5 ml tubes at  $4000 \times g$  for 5 min, the supernatant was then discarded, and 100  $\mu\text{L}$  distilled water was added to the pellet. This mixture was boiled to  $100^{\circ}\text{C}$  for 10 min and centrifuged at  $20000 \times g$  for 2 min. The supernatant was then stored at  $-20^{\circ}\text{C}$ . A PCR reaction for the 16S rRNA gene common to all eubacteria used universal primers (27 F 5' AGA GTT TGA TCC TGG CTC AG-3', 1492 R 5' GTT TAC CTT GTT ACG ACT T-3') to obtained of the PCR product (1465-bp). Nucleotide sequences were aligned using the BioEdit Sequence Alignment Editor (North Carolina State University, Raleigh, North Carolina) and these sequences were identified using the BLAST program (<http://www.ncbi.nlm.nih.gov>) (Altnok and Grizzle, 2001).

## Results

### Fungal and parasitic detections

From the beginning, the most distinct health problem was found to be suberication of the dead eggs of both

sturgeon species. The fungus isolated from the dead eggs was identified as *Saprolegnia* sp. From the hatching stage to the larval stage, the survival rate of the sturgeons was found to be 40% on day 90. *Trichodina* sp. was the only parasitic protozoan species found in the gills of the two species (prevalence, 0.5%) (Fig. 1). During the study period, no tumors were observed in the juveniles.




Fig. 1: *Trichodina* sp. (scale bar, 10  $\mu$ m)

### The problems of the feeding stage

As the artemia cysts were not completely removed after Artemia (*Artemia salina*) feeding, they were consumed by the larvae. Thus the leading health problem at this stage was death (30% mortality rate) caused by the deposition of *A. salina* cysts in the intestine. During artificial bait adaptation, a low rate of cannibalism (1%) was especially observed in *A. gueldenstaedtii* larvae that had completed the egg stage.

### Swim bladder problem

After three months, 5% of the fish were found to be affected by gas problems. Inflation of the swim bladder occurred in both species of sturgeon larvae (*A. baerii* and *A. gueldenstaedtii*) using the artificial bait (Figs. 2A, B). The fish also swam upside-down on the surface, exhibited problems in bait taking (Figs. 2C, D), and

displayed permanent spinal curvature due to the diurnal gas problem. A bacterial examination of these sturgeons showed the presence of *A. hydrophila* and *Bacillus mycoides* in the pneumatic duct which connects the air bladder and the pharynx.

### Bacterial detections

A total of 30 different bacterial strains were isolated from the external lesions and the internal organs of the sturgeons, and biochemical tests carried out after bacterial purification identified these bacteria as belonging to seven different bacterial species. Information on these bacterial species is given in Table 2. The species not subjected to the API test, Gram-positive rods (Fig. 3D) that formed a fringe shaped colony in TSA medium (Figs. 3A, B), demonstrated  $\beta$  hemolysis (Fig. 3C) and resistance to penicillin, were non-motile and were all classified as *B. mycoides* (Fig. 3C). The presence of many bacterial species implied a mixed infection of these fish, and hemorrhages were observed on the mouth of all bacteria-infected fish (Fig. 4A), apart from the presence of expanded internal organs, and unilateral exophthalmos (Fig. 4D).

### Other problems

Tumors were observed occasionally at the end of the ventral fins of the Russian sturgeons (*A. gueldenstaedtii*) that were older than two years and had reached a weight of 3 kg (Figs. 5C, D). Such tumors were not observed in *A. baerii* individuals. Moreover, mechanical injuries on the mouth were observed, especially in *A. baerii* individuals (Figs. 5A, B).

### Discussion

Fungal infestation of fish eggs is a common problem in all cultured fish species, and many studies have tried to overcome this problem by treating the eggs with disinfectants and improving water quality (Timur and Timur, 2003; Lasee, 1995; Noga, 2010). *Saprolegnia* sp. infestations during the incubation period are frequent in rainbow trout commonly reared in Turkey, and formaldehyde applications are used to treat these infections. An intense infestation of *Saprolegnia* sp. species also causes heavy loss in sturgeon eggs, and the reported death rate of *Acipenser persicus* eggs due to


Table 2: Bacterial species and their isolation area on fish

Bacteria	API CODES	Mole	n	Fish species	Tissues of fish
<i>Acinetobacter radioresistens</i>	API 20NE 0000032 (96.7%)	NA	3	<i>Acipenser gueldenstaedtii</i>	Skin, liver
<i>Aeromonas hydrophila</i>	API 20NE 7047127 (99.7%)	94%	7	<i>Acipenser baerii</i> <i>Acipenser gueldenstaedtii</i>	Mouth, skin, liver, pneumatic duct
<i>Aeromonas sobria</i>	API 20NE 7136755 (98.4%)	97%	2	<i>Acipenser baerii</i>	Liver
<i>Bacillus mycoides</i>	NA	NA	5	<i>Acipenser baerii</i>	Esophagus
<i>Citrobacter freundii</i>	API 20E 1604573 (95%)	95%	2	<i>Acipenser baerii</i> <i>Acipenser gueldenstaedtii</i>	Liver, kidney
<i>Pseudomonas</i> sp.	API 20NE 3467747 (94%)	95%	8	<i>Acipenser gueldenstaedtii</i>	Mouth, liver, kidney
<i>Pseudomonas fluorescens</i>	API 20NE 0057555 (99.6%)	96%	4	<i>Acipenser baerii</i>	Skin
<i>Pseudomonas putida</i>	API 20NE 0142457 (99.8%)	96%	1	<i>Acipenser baerii</i>	Liver
<i>Serratia</i> sp.	API 20E 5305363 (91%)	93%	3	<i>Acipenser gueldenstaedtii</i>	Liver

n: Number of bacteria, and NA: Not application




**Fig. 2:** Inflation of swim bladder, fill with gas and spinal deformities in *Acipenser baerii* larvae. Gas problems in swim bladders of different size of fish (A, B), spinal deformities (C, D) (scale bars A: 3 cm, and B: 7 cm)




**Fig. 3:** *Bacillus mycoides* isolated from the esophagus of *Acipenser baeri*. The colonies of *Bacillus mycoides* on TSA medium (A, B), β hemolysis on blood agar (C), and Gram-stain of the bacteria (D)





**Fig. 4:** External deformations on the fish caused by the bacteria. Hemorrhages on around the mouth (A), and the base of dorsal and caudal fins (B, C), unilateral exophthalmus (black arrow) on the fish *Acipenser gueldenstaedtii*, spiraculum (white arrow) (D) (scale bars, A: 2 cm, B: 2 cm, C: 5 cm, and D: 5 cm)



**Fig. 5:** Mechanical injuries and tumor on *Acipenser gueldenstaedtii*, mechanical injuries on the mouth of the fish (A, B), clear tumors on the base of the anal fins of the fish (C, D) (scale bars, A: 5 cm, and C: 4 cm)

*Saprolegnia* sp. is between 7% and 22% (Jalilpoor *et al.*, 2006). Post-infection measures and ensuring good microbiological quality of the water and its sufficient ventilation are very important. Careful removal of the infected eggs is another technique effective in decreasing loss. Nanosil, Chloramine-T and hydrogen peroxide can also be used to prevent fungal infestation of sturgeon eggs (Ghazvini and Rudsari, 2012). Available data on cannibalism in sturgeons predominantly pertain to the giant sturgeon (*Huso huso*); nevertheless, reports of cannibalism in other sturgeon species also exist. Cannibalism has been observed in *A. gueldenstaedtii* larvae at post-incubation days 18 and 36 (Memiş *et al.*, 2009). In the present study, we observed a low rate of cannibalism which occurred only in *A. gueldenstaedtii* larvae. To avoid cannibalism, it is important to pay attention to bait sequence and live and artificial baits. For various reasons, the swim bladder disease is very common in fish; the supersaturation of gas, bacterial infections, and sudden changes in water temperature and pressure have all been reported as causes of air bladder diseases (Austin and Austin, 2007). In this study, the air bladder gas disease was observed in both species of sturgeon due to the supersaturation of gas after adaptation to artificial bait. This can be due to the fact that sturgeons are physostome fish and their air bladder is connected to their esophagus. A large quantity of bacteria was isolated from the esophageal samples, and *A. hydrophila* was found to be the predominant species. Bacteria such as *Yersinia ruckeri*, *A. hydrophila* and *Vibrio* spp., have an important role among pathogens that cause fish diseases (Kayis *et al.*, 2009). Literature on bacterial pathogens of sturgeons is limited, probably because the aquaculture of sturgeons is not as common as that of other species and also because sturgeons are quite resistant to diseases after they reach a certain size. Bacterial pathogens isolated from sturgeons in recent years include *Flavobacterium johnsoniae*, *A. hydrophila* (*Acipenser gueldenstaedtii*) (Timur *et al.*, 2010) and *Flavobacterium columnare* (*Acipenser oxyrinchus desotoi*) (Altınok *et al.*, 2001). In our study, we isolated *A. hydrophila*, *Acinetobacter radioresistens*, *Aeromonas sobria*, *A. salmonicida*, *B. mycoides*, *Citrobacter freundii*, *Pseudomonas fluorescens*, *Pseudomonas putida* and *Serratia* sp. species from the sturgeons; this knowledgebase is important as it will help create a bacterial profile of sturgeons reared in Turkey. The existence of parasitic pathogens in sturgeons has been frequently reported, especially in those cultured in earth pools. *T. reticulata*, *D. spathaceum*, and helminths are some of the parasites that have been isolated from sturgeons (Bazari *et al.*, 2010). In our study, *Trichodina* sp. was the only parasitic species isolated from sturgeons. It is possible that the use of fiberglass tanks rather than soil pools, provided more favorable conditions for sturgeon culture, especially during the incubation period. Nevertheless, reports of parasites isolated from sturgeons are available and specifically differentiate between fish reared in soil pools and those cultured in their natural environment.

In the present study, we reported on the health problems of sturgeons from the incubation period until they reached a body weight of 3 kg. The following conclusions and recommendations are based on our observations. As aquaculture of sturgeons in Turkey can be initially developed only by importing eggs, more detailed studies are necessary to investigate health problems during production stages. Sturgeons need to be treated with appropriate disinfectants during the incubation period, dead eggs should be removed immediately and the water should be properly ventilated. Baiting should be done frequently in the larval stage to avoid cannibalism. To prevent gas disease and hemorrhage-related deaths due to bacterial pathogens, microbial contamination and water quality should be carefully monitored. Using fiberglass and concrete ponds instead of earth pools may help reduce or even avoid parasite infestations.

## Acknowledgement

The study was funded by Recep Tayyip Erdoğan University Research Project Fund (Project number; 2011.103.02.3).

## References

- Altınok, I and Grizzle, JM (2001). Effects of low salinities on *Flavobacterium columnare* infection of euryhaline and freshwater stenohaline fish. *J. Fish Dis.*, 24: 361-367.
- Austin, B and Austin, DA (2007). *Bacterial fish pathogens: diseases of farmed and wild fish*. 4th Edn., Chichester, UK, Praxis Publ. Ltd., PP: 357-411.
- Bauer, ON; Pugachev, ON and Voronin, VN (2002). Study of parasites and diseases of sturgeons in Russia: a review. *J. Appl. Ichthyol.*, 18: 420-429.
- Bazari Moghaddam, S; Mokhayer, B; Shenavar Masouleh, AR; Jalilpoor, J; Masoumzadeh, M and Alizadeh, M (2010). Study on parasitic infestation on Persian sturgeon (*Acipenser persicus*) larvae and fingerlings in the Shahid Dr. Beheshti Hatchery. *JIFRO.*, 9: 342-351.
- Chebanov, MS and Galich, EV (2013). Sturgeon hatchery manual. FAO Fisheries and Aquaculture Technical Paper 558. PP: 1-17.
- Conte, FS (1988). Hatchery manual for the white sturgeon (*Acipenser transmontanus* Richardson): with application to other North American Acipenseridae. University of California (System). Cooperative Extension, UCANR Publications.
- Costinar, L; Herman, V; Pascu, C; Marcu, AD; Marcu, A and Faur, B (2010). Isolation and characterization of *Vibrio alginolyticus* and *Pasteurella* spp. from Siberian sturgeon *Acipenser baerii*. *Lucrari Stiintifice Medicina Veterinara*. 43: 125-128.
- Ghazvini, A; Rudsari, HV; Takami, GA; Masuleh, ARS and Ashourpour, A (2012). Disinfection efficiency of three anti-fungal agents (nanosil, chloramine-T and hydrogen peroxide) on Persian sturgeon (*Acipenser persicus*, Borodin 1897) larvae. *Int. J. Biol.*, 4: 138-145.
- Jalilpoor, J; Masouleh, AS and Masoumzadeh, M (2006). Fungal flora in *Acipenser persicus* eggs with particular emphasis on *Saprolegnia* sp. (Oomycetes) and mortality during mass incubation at the Shahid Beheshti hatchery. *J.*

- Appl. Ichthyol., 22: 265-268.
- Karataş, S; Ercan, D; Stenium, TM; Turgay, E; Memiş, D and Candan, A** (2010). First isolation of a *Flavobacterium johnsoniae* like bacteria from Cultured Russian sturgeon in Turkey. J. Anim. Vet. Adv., 9: 1943-1946.
- Kayis, S; Ozcelep, T; Capkin, E and Altinok, I** (2009). Protozoan and metazoan parasites of fish in the Turkey and their applied treatments. IJA., 61: 93-102.
- Kurtoğlu, İZ; Ak, K; Delihasan Sonay, F; Kayış, Ş; Balta, F; Köse, Ö and Şahin, T** (2015). Encountered deformations, during the larvae development and early feeding stages in Siberian sturgeon, *Acipenser baerii*. ECJSE., 2: 1-11.
- Lasee, BA** (1995). Introduction to fish health management, U.S. Fish and Wildlife Service. La Crosse Fish Health Center 555, Lester Avenue Onalaska, Wisconsin, 54650.
- Lom, J and Dykova, L** (1992). Protozoan parasites of fishes. Dev. Aquacult. Fish. Sci., 26: 250-301.
- Memiş, D; Ercan, E; Çelikkale, MS; Timur, M and Zarkua, Z** (2009). Growth and survival rate of Russian sturgeon (*Acipenser gueldenstaedtii*) larvae from fertilized eggs to artificial feeding. Turkish J. Fish. Aquatic Sci., 9: 47-52.
- Mohler, JV** (2003). Culture manual for the Atlantic sturgeon. A Region 5 U.S. Fish & Wildlife Service Publication 300 Westgate Center Drive Hadley, Massachusetts.
- Noga, EJ** (2010). *Fish disease – diagnosis and treatment*. 2nd Edn., New Jersey, Hoboken, Wiley-Blackwell. PP: 197-350.
- Prearo, M; Squadrone, S; Fioravanti, ML; Giorgi, I; Marturano, S; Gasparri, F; Abete, MC and Zannoni, RG** (2009). Italian farmed sturgeon: mortality events during 2004-2008, WAS 2009, Veracruz, Mexico. P: 681.
- Timur, G; Akaylı, T; Korun, J and Yardımcı, ERA** (2010). Study on bacterial haemorrhagic septicemia in farmed Young Russian sturgeon in Turkey (*Acipenser gueldenstaedtii*). Turk. J. Fish. Aquat. Sci., 25: 19-27.
- Timur, M and Timur, G** (2003). *Fish diseases*. Publications of Istanbul University, Publication No: 4426, Section of Aquatic Publication Number: 5. PP: 1-213.
- Yang, W and Li, A** (2009). Isolation and characterization of *Streptococcus dysgalactiae* from diseased *Acipenser schrenckii*. Aquaculture. 294: 14-17.
- Yu-Ping, H and Di, W** (2005). A review of sturgeon virosis. J. Forestry Res., 16: 79-82.