

# Long-Term Results of Various Operations for Lumbar Disc Herniation: Analysis of over 39,000 Patients

George J. Dohrmann Nassir Mansour

Section of Neurosurgery, University of Chicago Medical Center, Chicago, Ill., USA

### Key Words

Lumbar disc herniation · Lumbar disc surgery · Long-term follow-up · Lumbar disc operations · Operations for lumbar disc herniation · Results of lumbar disc surgery

### Abstract

**Objective:** To determine the long-term follow-up of the various operations for lumbar disc herniation in a large patient population. **Subjects and Methods:** Patients who had operations for lumbar disc herniation (microdiscectomy, endoscopic microdiscectomy and the 'classical operation', i.e. laminectomy/laminotomy with discectomy) were collected from the world literature. Patients who had follow-ups for at least 2 years were analyzed relative to the outcome. The outcome was graded by the patients themselves, and the operative groups were compared to one another. **Results:** 39,048 patients collected from the world literature had had lumbar disc operations for disc herniations. The mean follow-up period was 6.1 years, and 30,809 (78.9%) patients reported good/excellent results. Microdiscectomy was performed on 3,400 (8.7%) patients. The mean follow-up was 4.1 years with 2,866 (84.3%) good/excellent results, while 1,101 (3.6%) patients had endoscopic microdiscectomy. There, the mean follow-up was 2.9 years with 845 (79.5%) good/excellent results. The classical operation was performed on 34,547 (88.5%) patients with a mean follow-up

period of 6.3 years, and 27,050 (78.3%) patients had good/excellent results. These results mirror those with discectomy and the placement of prosthetic discs. **Conclusions:** The analysis of 39,048 patients with various operations for lumbar disc herniation revealed the same pattern of long-term results. Patients who had microdiscectomy, endoscopic microdiscectomy or the classical operation (laminectomy/laminotomy with discectomy) all had approximately 79% good/excellent results. None of the operative procedures gave a different outcome.

© 2015 S. Karger AG, Basel

### Introduction

Operations for lumbar disc herniation are numerous [1–56] and have been performed since the 1934 publication by Mixter and Barr [33]. Indeed, operations for lumbar disc herniation are the most frequently done neurosurgical procedures. However, unlike the long-term success rate of posterior operations for cervical disc herniation, which is 94%, the overall long-term success rate for operations for lumbar disc herniations is considerably lower. For this reason, various operations were done with the hope that the long-term success rate would improve.

Since the 'classical operation' (laminectomy/laminotomy with discectomy), other approaches have been used.

Of these, the most popular are 2 operative procedures: (a) microdiscectomy and (b) endoscopic microdiscectomy. The average size of the published series of operations (classical, microdiscectomy and endoscopic microdiscectomy) is only several hundred patients/series, and most series did not attempt assessing the long-term outcome. The purpose of the present study was to analyze the long-term outcome of operations for lumbar disc herniation and then, specifically, the outcomes for each of the following: microdiscectomy, endoscopic microdiscectomy and the classical operation. To nullify the occasional unusual result, the goal of the study was to analyze the largest number of such patients published to date.

## Subjects and Methods

All patients operated on for lumbar disc herniation with radiculopathy and followed for a minimum of 2 years postoperatively were collected from the world literature (using references cited in published studies, the studies themselves and the website PubMed, the search was complete). Good/excellent outcome was measured by the patients' own analyses because the best outcome measure is most simply what the patient thinks about the outcome [19, 24]. Hobbs et al. [19] noted that the patient's perception was the 'true measure of success'. Then, the operations with good/excellent outcomes were tabulated. The patients were divided into groups by the type of operative procedure, i.e. microdiscectomy, endoscopic microdiscectomy and the classical operation, and were analyzed by the time of follow-up and by which patients had good/excellent results by the patients' own assessment.

## Results

Thirty-nine thousand forty-eight patients collected from the world literature had had operations for lateral lumbar disc herniation with radiating pain and met the follow-up requirement of at least 2 years. Of the 39,048 operations, 95% of lumbar disc herniations were at the lowest 2 levels of the lumbar spine, and 49 and 46% were at L<sub>4-5</sub> and L<sub>5-S<sub>1</sub></sub>, respectively. Of the remaining 5% lumbar disc herniations, 0.15% were at L<sub>1-2</sub>, 0.65% were at L<sub>2-3</sub> and 4.2% were at L<sub>3-4</sub> (table 1). The mean follow-up period in this series was 6.1 years. Of all the patients, 30,809 (78.9%) had good/excellent outcomes (table 2). Microscopic discectomy was performed on 3,400 (18.7%) patients with a mean follow-up of 4.1 years. Good/excellent results occurred in 32,917 (84.3%) patients (table 3). The endoscopic microdiscectomy group consisted of 1,101 (3.6%) patients with a mean follow-up period of 2.9 years, and 845 (79.5%) patients had good/excellent results (table 4). Of the

**Table 1.** Level of herniated lumbar discs

L <sub>1-2</sub>	0.15%
L <sub>2-3</sub>	0.65%
L <sub>3-4</sub>	4.2%
L <sub>4-5</sub>	49%
L <sub>5-S<sub>1</sub></sub>	46%

39,048 patients, 34,547 (88.5%) had the classical operation (laminectomy/laminotomy with discectomy). The mean follow-up was 6.3 years. The patients had 78.3% good/excellent results (table 5).

## Discussion

In the 8 decades since the publication by Mixter and Barr [33], many studies of the surgical management of lumbar disc herniation with radiculopathy have been published showing the results of laminectomy/laminotomy with discectomy. Another operative approach was described over 4 decades later, i.e. microdiscectomy [7, 52, 55]. Later still, another surgical approach to lumbar disc herniation was developed with the advent of endoscopic microdiscectomy [17]. Many series were published about the above 3 surgical techniques. Significantly fewer publications dealt with the long-term results of these different operative approaches.

In studies focused on the long-term results of the surgical management of lumbar disc herniation, most series that were published averaged several hundred patients. In the 45 studies analyzed here, the mean number of patients was 382/series. This study analyzes 39,048 patients operated for lumbar disc herniation with radiculopathy and followed for over 6 years.

Approximately 79% of the outcomes, graded by the patients, were good/excellent. A series of this size is not affected by slight variations in technique with various surgeons and by variations in the patients' ages and gender.

Each of the operations for this problem was an attempt to improve the outcome by using different operative approaches and techniques; however, as is shown in the present analysis, there is no real difference in the long-term outcome with the above operations. Good/excellent outcomes were 79% overall and 84% for microdiscectomy, 80% for endoscopic microdiscectomy and 78% for the classical operation (laminectomy/laminotomy and discectomy). Another attempt at improving the outcome was the use of the prosthetic disc; however, in long-term

**Table 2.** Long-term results of operations for lumbar disc herniation

Authors [Ref.]	Patients, n	Mean follow-up, years	Good/excellent results, %
Asch et al. [2]	212	2	80
Atlas et al. [3]	217	10	69
Bakhsh [4]	39	10	79
Butterman [5]	100	2.5	92
Casal-Moro et al. [6]	120	5	95
Chang et al. [8]	26	3	Not given
Cooper and Feuer [10]	100	2	94
Davis [11]	984	10.8	89
Dewing et al. [12]	183	2.1	85
Dvorak [14]	371	10.5	72
Ebeling et al. [15]	485	2	73
Findlay et al. [16]	79	10	83
Gurdjian et al. [18]	623	9	74
Hsu et al. [20]	226	2	82
Jansson et al. [21]	22,261	6	78
Jensdottir et al. [22]	134	20.7	91
Kotilainen et al. [23]	237	2	92
Lewis et al. [26]	83	7.5	89
Liu et al. [27]	82	6.4	84
Loupasis et al. [28]	109	12.7	64
Mariconda et al. [29]	201	27.8	90
Marin [30]	600	10	83
Martinez Quinones et al. [32]	142	5	93
Moore et al. [34]	100	8.6	93
Naylor [35]	204	17.5	79
Nykqvist et al. [36]	197	12.9	81
Österman et al. [37]	28	2	93
Padua et al. [38]	120	12.1	77
Papavero and Caspar [39]	200	Not given	79
Pappas et al. [40]	654	1.5	96
Parker et al. [41]	111	3.1	68
Peul et al. [42]	125	2	81
Salenius and Laurent [43]	695	6	63
Schoegg et al. [44]	672	6.3	77
Schramm et al. [45]	3,238	4	80
Silverplats et al. [46]	140	7.3	70
Spangfort [47]	2,503	Not given	77
Tregonning et al. [48]	91	9.7	63
Vik et al. [49]	124	8.5	81
Weber [50]	56	4	86
Weinstein et al. [51]	245	4	84
Williams [52]	530	3	91
Woertgen et al. [53]	98	2.3	66
Wu et al. [54]	1,231	2.3	77
Yorimitsu et al. [56]	72	14.3	87
Total	39,048		
Mean		6.1	78.9

**Table 3.** Long-term results of operations for lumbar disc herniation: microdiscectomy

Authors [Ref.]	Patients, n	Mean follow-up, years	Good/excellent results, %
Asch et al. [2]	212	2	80
Cooper and Feuer [10]	100	2	94
Dewing et al. [12]	183	2.1	85
Ebeling et al. [15]	485	2	73
Findlay et al. [16]	79	10	83
Jensdottir et al. [22]	134	20.7	91
Kotilainen et al. [23]	237	2	92
Moore et al. [34]	100	8.6	93
Österman et al. [37]	28	2	93
Papavero and Caspar [39]	100	Not given	79
Pappas et al. [40]	353	1.5	96
Peul et al. [42]	125	2	81
Schoegg et al. [44]	672	6.3	77
Vik et al. [49]	62	8.5	81
Williams [52]	530	3	91
Total	3,400		
Mean		4.1	84.3

**Table 4.** Long-term results of operations for lumbar disc herniation: endoscopic microdiscectomy

Authors [Ref.]	Patients, n	Mean follow-up, years	Good/excellent results, %
Casal-Moro et al. [6]	120	5	95
Chang et al. [8]	26	3	Not given
Liu et al. [27]	82	6.4	84
Wu et al. [54]	873	2.3	77
Total	1,101		
Mean		2.9	79.5

**Table 5.** Long-term results of operations for lumbar disc herniation: laminectomy/laminotomy with discectomy

Authors [Ref.]	Patients, n	Mean follow-up, years	Good/excellent results, %
Atlas et al. [3]	217	10	69
Bakhsh [4]	39	10	79
Butterman [5]	100	2.5	92
Davis [11]	984	10.8	89
Dvorak [14]	371	10.5	72
Gurdjian et al. [18]	623	9	74
Hsu et al. [20]	226	2	82
Jansson et al. [21]	22,261	6	78
Lewis et al. [26]	83	7.5	89
Loupasis et al. [28]	109	12.7	64
Mariconda et al. [29]	201	27.8	90
Marin [30]	600	10	83
Martinez Quinones et al. [32]	142	5	93
Naylor [35]	204	17.5	79

**Table 5** (continued)

Authors [Ref.]	Patients, n	Mean follow-up, years	Good/excellent results, %
Nykqvist et al. [36]	197	12.9	81
Padua et al. [38]	120	12.1	77
Papavero and Caspar [39]	100	Not given	79
Pappas et al. [40]	301	1.5	96
Parker et al. [41]	111	3.1	68
Salenius and Laurent [43]	695	6	63
Schramm et al. [45]	3,238	4	80
Silverplats et al. [46]	140	7.3	70
Spangfort [47]	2,503	Not given	77
Tregonning et al. [48]	91	9.7	63
Vik et al. [49]	62	8.5	81
Weber [50]	56	4	86
Weinstein et al. [51]	245	4	84
Woertgen et al. [53]	98	2.3	66
Wu et al. [54]	358	2.3	77
Yorimitsu et al. [56]	72	14.3	87
Total	34,547		
Mean		6.3	78.3

studies (46 patients at 3.2 years of follow-up; 105 patients at 4.3 years of follow-up), the good/excellent results were 77 and 79%, respectively [9, 25]. All of the operations analyzed have good/excellent results of around 79%. Different approaches and different techniques do not appear to have made any real difference in the long-term outcome.

The results of posterior operations for lumbar disc herniation are not as good as the results of posterior operations for cervical disc herniation. An analysis of over 3,000 such posterior operations for cervical disc herniation with an 8.5-year mean follow-up revealed 94% good/excellent results [13]. Why is there this difference of 79% versus 94%? Surely the operative procedures were successful in both groups, but much more so in the cervical spine. The

reason for this difference needs further analysis and, perhaps, yet another approach, surgical or otherwise, to the problem of lumbar disc herniation with radiculopathy.

## Conclusion

Each of the operations for lumbar disc herniation (microdiscectomy, endoscopic microdiscectomy and laminectomy/laminotomy with discectomy) had approximately 79% good-to-excellent results. There was no difference in the long-term follow-up in any of the operative groups, including the use of a lumbar disc prosthesis ('artificial disc').

## References

- 1 Abramovitz JN, Neff SR: Lumbar disc surgery: results of the Prospective Lumbar Discectomy Study of the Joint Section on Disorders of the Spine and Peripheral Nerves of the American Association of Neurological Surgeons and the Congress of Neurological Surgeons. *Neurosurgery* 1991;29:301–308.
- 2 Asch HL, Lewis J, Moreland DB, et al: Prospective multiple outcomes study of outpatient lumbar microdiscectomy: should 75 to 80% success rates be the norm? *J Neurosurg* 2002;96(suppl 1):34–44.
- 3 Atlas SJ, Keller RB, Wu YA, et al: Long-term outcomes of surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: 10 year results from the Maine Lumbar Spine Study. *Spine (Phila Pa 1976)* 2005;30:927–935.
- 4 Bakhsh A: Long-term outcome of lumbar disc surgery: an experience from Pakistan. *J Neurosurg Spine* 2010;12:666–670.
- 5 Buttermann GR: Treatment of lumbar disc herniation: epidural steroid injection compared with discectomy. A prospective randomized study. *J Bone Joint Surg* 2004;86:670–679.
- 6 Casal-Moro R, Castro-Menendez M, Hernandez-Blanco M, et al: Long-term outcome after microendoscopic discectomy for lumbar disk herniation: a prospective clinical study with 5-year follow-up. *Neurosurgery* 2011;68:1568–1575.
- 7 Caspar W: A new surgical procedure for lumbar disc herniation causing less tissue damage through a microsurgical approach. *Adv Neurosurg* 1977;4:74–80.

8 Chang S-S, Fu T-S, Liang Y-C, et al: Results of microendoscopic discectomy performed in the 26 cases with a minimum 3 years follow-up. *Chang Gung Med J* 2009;32:89–96.

9 Cinotti G, David T, Postacchini F: Results of disc prosthesis after a minimum follow-up period of 2 years. *Spine (Phila Pa 1976)* 1996; 21:995–1000.

10 Cooper DF, Feuer H: Lumbar microdiscectomy. *J Indiana State Med Assoc* 1982;74: 674–675.

11 Davis RA: A long-term outcome analysis of 984 surgically treated herniated lumbar discs. *J Neurosurg* 1994;80:415–421.

12 Dewing CB, Provencher MT, Riffenburgh RH, et al: The outcomes of lumbar microdiscectomy in a young, active population. Correlation by herniation type and level. *Spine (Phila Pa 1976)* 2008;33:33–38.

13 Dohrmann GJ, Hsieh JC: Long-term results of anterior versus posterior operations for herniated cervical discs: analysis of 6,000 patients. *Med Princ Pract* 2014;23:70–73.

14 Dvorak J, Gauchat M-H, Valach L: The outcome of surgery for lumbar disc herniation. I. A 4–17 years' follow-up with emphasis on somatic aspects. *Spine (Phila Pa 1976)* 1988;13: 1418–1422.

15 Ebeling U, Reichenberg W, Reulen HJ: Results of microsurgical lumbar discectomy. Review on 485 patients. *Acta Neurochir (Wien)* 1986;81:45–52.

16 Findlay GF, Hall BI, Musa BS, et al: A 10-year follow-up of the outcome of lumbar microdiscectomy. *Spine (Phila Pa 1976)* 1998;23: 1168–1171.

17 Foley KT, Smith MM: Microendoscopic discectomy. *Techn Neurosurg* 1997;3:301–307.

18 Gurdjian ES, Webster JE, Ostrowski AZ, et al: Herniated lumbar intervertebral discs – an analysis of 1176 operated cases. *J Trauma* 1961;1:158–176.

19 Hobbs J, Bina R, Dohrmann G, et al: What's new in spine surgery. *Med Princ Pract* 2013; 22:101–102.

20 Hsu WK, McCarthy KJ, Savage JW, et al: The Professional Athlete Spine Initiative: outcomes after lumbar disc herniation in 342 elite professional athletes. *Spine J* 2011;11:180–186.

21 Jansson KA, Nemeth G, Granath F, et al: Surgery for herniation of a lumbar disc in Sweden between 1987 and 1999. An analysis of 25,576 operations. *J Bone Joint Surg Br* 2004;86-B:841–847.

22 Jendsdottir M, Gundmundsson K, Hannesson B, et al: 20 years follow-up after the first microsurgical lumbar discectomies in Iceland. *Acta Neurochir (Wien)* 2007;149:51–58.

23 Kotilainen E, Valtonen S, Carlson CA: Microsurgical treatment of lumbar disc herniation: follow-up of 237 patients. *Acta Neurochir (Wien)* 1993;120:143–149.

24 Lattig F, Grob D, Kleinstueck FS, et al: Ratings of global outcome at the first post-operative assessment after spinal surgery: how often do the surgeon and patient agree? *Eur Spine J* 2009;18(suppl 3):S386–S394.

25 Lemaire JP, Skalli W, Lavaste F, et al: Intervertebral disc prosthesis. Results and prospect for the year 2000. *Clin Orthop Relat Res* 1997; 337:64–76.

26 Lewis PJ, Weir BKA, Borad RW, et al: Long-term prospective study of lumbosacral discectomy. *J Neurosurg* 1987;67:49–53.

27 Liu W-G, Wu X-T, Guo J-H, et al: Long-term outcomes of patients with lumbar disc herniation treated with percutaneous discectomy: comparative study with microendoscopic discectomy. *Cardiovasc Intervent Radiol* 2010;33: 780–786.

28 Loupasis GA, Stamos K, Katonis PG, et al: Seven- to 20-year outcome of lumbar discectomy. *Spine (Phila Pa 1976)* 1999;24:2313–2317.

29 Mariconda M, Galasso O, Secondulfo V, et al: Minimum 25-year outcome and functional assessment of lumbar discectomy. *Spine (Phila Pa 1976)* 2006;31:2593–2599.

30 Marin GA: Lumbar disc protrusion. Evaluation and study of 600 discectomies with one to ten years followup. *Int Surg* 1974;59:154–155.

31 Maroon JC: Current concepts in minimally invasive discectomy. *Neurosurg* 2002;51: S137–S145.

32 Martinez Quinones JV, Aso J, Consolini F, et al: Long-term outcomes of lumbar microdiscectomy in a working class sample (in Spanish). *Neurocirugia* 2011;22:235–244.

33 Mixter WJ, Barr JS: Rupture of the intervertebral disc with involvement of the spinal canal. *N Engl J Med* 1934;211:210–215.

34 Moore AJ, Chilton JD, Uttley D: Long-term results of microlumbar discectomy. *Br J Neurosurg* 1994;8:319–326.

35 Naylor A: The late results of laminectomy for lumbar disc prolapse. A review after ten to twenty-five years. *J Bone Joint Surg Br* 1974; 56:17–29.

36 Nykvist F, Hurme M, Alaranta H, et al: Severe sciatica: a 13-year follow-up of 342 patients. *Eur Spine J* 1995;4:335–338.

37 Österman H, Seitsalo S, Karppinen J, et al: Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up. *Spine (Phila Pa 1976)* 2006;31:2409–2414.

38 Padua R, Padua S, Romanini E, et al: Ten- to 15-year outcome of surgery for lumbar disc herniation: radiographic instability and clinical findings. *Eur Spine J* 1999;8:70–74.

39 Papavero L, Caspar W: The lumbar microdiscectomy. *Acta Orthop Scand Suppl* 1993;251: 34–37.

40 Pappas CTE, Harrington T, Sonntag VKH: Outcome analysis in 654 surgically treated lumbar disc herniations. *Neurosurgery* 1992; 30:862–866.

41 Parker SL, Xu R, McGirt MJ, et al: Long-term back pain after a single level discectomy for radiculopathy: incidence and health care cost analysis. *J Neurosurg Spine* 2010;12:178–182.

42 Peul WC, van den Hout WB, Brand R, et al: Prolonged conservative care versus early surgery in patients with sciatica caused by lumbar disc herniation: two year results of a randomized controlled trial. *BMJ* 2008;336: 1355–1358.

43 Salenius P, Laurent LE: Results of operative treatment of lumbar disc herniation. A survey of 836 patients. *Acta Orthop Scand* 1977;48: 630–634.

44 Schoegl A, Reddy M, Matula C: Functional and economic outcome following microdiscectomy for lumbar disc herniation in 672 patients. *J Spinal Disord Tech* 2003;16:150–155.

45 Schramm J, Opel F, Umbach W, et al: Complications after lumbar operation on vertebral disks. Results of a statistical survey (in German). *Nervenarzt* 1978;49:26–33.

46 Silverplats K, Lind B, Zoega B, et al: Clinical factors of importance for outcome after lumbar disc herniation surgery: long-term follow-up. *Eur Spine J* 2010;19:1459–1467.

47 Spangfort EV: The lumbar disc herniation. A computer-aided analysis of 2,504 operations. *Acta Orthop Scand Suppl* 1972;142:1–95.

48 Tregonning GD, Transfeldt EE, McCulloch JA, et al: Chymopapain versus surgery for lumbar disc herniation. 10-year results of treatment. *J Bone Joint Surg Br* 1991;73:481–486.

49 Vik A, Zwart JA, Hulleberg G, et al: Eight year outcome after surgery for lumbar disc herniation: a comparison of reoperated and not reoperated patients. *Acta Neurochir (Wien)* 2001;143:607–611.

50 Weber H: Lumbar disc herniation. A controlled, prospective study with ten years of observation. *Spine (Phila Pa 1976)* 1983;8:131–140.

51 Weinstein JN, Lurie JD, Tosteson TD, et al: Surgical versus non-operative treatment for lumbar disc herniation: four-year results for the Spine Patient Outcomes Trial (SPORT). *Spine (Phila Pa 1976)* 2008;33:2789–2800.

52 Williams RW: Microlumbar discectomy. A conservative surgical approach to the virgin herniated lumbar disc. *Spine (Phila Pa 1976)* 1978;3:175–182.

53 Woertgen C, Rothoerl RD, Brem K, et al: Variability of outcome after lumbar disc surgery. *Spine (Phila Pa 1976)* 1999;24:807–811.

54 Wu X, Zhuang S, Mao Z, et al: Microendoscopic discectomy for lumbar disc herniation. Surgical technique and outcome in 873 consecutive cases. *Spine (Phila Pa 1976)* 2006;31: 2689–2694.

55 Yasargil MG: Microsurgical operation of herniated lumbar disc. *Adv Neurosurg* 1977;4: 81–84.

56 Yorimitsu E, Chiba K, Toyama Y, et al: Long-term outcomes of standard discectomy for lumbar disc herniation. A follow-up study of more than 10 years. *Spine (Phila Pa 1976)* 2001;26:652–657.