

Original Article**Effects of Ramadan fasting on acute upper gastrointestinal bleeding due to peptic ulcer***M. H. Emami**, *H. Rahimi*****Abstract**

BACKGROUND: As, acid negatively affects duodenal and gastric mucosal defense, we designed this study to investigate if fasting during Ramadan can promote peptic ulcer and its complication, acute upper gastrointestinal bleeding.

METHODS: All patients files who had admitted to hospital with acute upper gastrointestinal bleeding from 10th day of Ramadan till one month later, in 2002 to 2004 were reviewed. Patients were divided into two groups: fasting group who were fasting, at least 10 days before admission and non-fasting group.

RESULTS: A total of 236 patients were included in the study; fasting group involved 108 subjects and non-fasting group 128 ones. Both groups were matched regarding sex, predisposing factors and underlying diseases. Fasting group showed more frequent duodenal ulcer (38%) compared to non-fasting group (19.5%) ($P = 0.002$). The frequency of esophageal varices was significantly higher in non-fasting group. There was no correlation between fasting and other causes of gastrointestinal bleeding. In fasting group 38%, and in non-fasting group 18.9%, had previous dyspeptic symptoms ($P = 0.001$). Two groups were similar regarding prognostic factors.

CONCLUSION: It seems Ramadan fasting can increase acute upper gastrointestinal bleeding due to duodenal ulcer, but it does not make its prognosis poorer than that in non-fasting patients. Dyspeptic symptoms could be considered as alarming signs for patients who want to fast during Ramadan.

KEYWORDS: Ramadan, fasting, acute upper gastrointestinal bleeding, duodenal ulcer.

JRMS 2006; 11(3): 170-175

Muslims fast from dawn to sunset during Ramadan (the ninth lunar month in Hegira calendar) and it is obligatory for all healthy adults to fast¹. Over 50% of muslims fast every day or on some days of Ramadan and some of them can not fast due to medical problems forbidding fasting².

Three major causes of upper GI bleeding are peptic ulcer disease (PUD), gastric erosion and esophageal varices. PUD includes duodenal, gastric and stomal ulcers account for about 50% of upper GI bleeding episodes³. Acid is still an independent factor that, negatively, affects mucosal defense and plays an important

role in the pathogenesis of PUD, especially, duodenal ulcer⁴. Roles of gastric secretion, motility, and mucosal abnormalities as primary pathogenic elements in ulcer development have also been noted⁴. Factors that can cause bleeding or rebleeding from peptic ulcer include H. Pylori, NSAIDS and gastric acidity⁵⁻⁷.

During Ramadan, gastrin, pepsin and acid secretions increase and then, will return to pre-Ramadan level one month later⁸⁻¹⁰. It is indicated that the median pH is 2.3 prior to Ramadan, 1 at day 10 and day 24 of Ramadan and 1.6 one month later¹⁰. The gastric acid secretion is increased by 45% at day 10 of Ramadan

Poursina Hakim Institute, Department of Gastroenterology, Isfahan University of Medical Sciences, Isfahan, Iran.

*Assistant Professor of Internal Medicine and Gastroenterology, Isfahan University of Medical Sciences, Isfahan, Iran.

**Resident of Internal Medicine

Correspondence to: Dr Hojatollah Rahimi, Resident of Internal Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.
e-mail: h_rahimi@resident.mui.ac.ir

compared with its level prior to Ramadan and the diurnal increase is much more than the nocturnal one (122% and 25%, respectively)¹⁰. The gastric acid secretion is 23% higher than baseline even one month after Ramadan¹⁰. These modifications in gastric acid secretion are likely fundamental to the increase of dyspeptic symptoms observed during the Ramadan.

One study showed higher rate of peptic ulcer complications during Ramadan compared with the same period before and after it;¹¹ but fasting had no influence on the outcome of duodenal ulcers treated by lansoprazole, omeperazole or H₂-blockers; i.e. healing rate (90% to 95%) was similar in fasting and non-fasting groups¹²⁻¹⁴.

In another study, psychological changes especially changes in irritability were increased continuously during Ramadan and reached its peak at the end of the month especially in smokers¹⁵. More over, during Ramadan fasting, medication intake compliance decreases; therefore it may worsen or relapse underlying disease such as PUD in fasting patients. Also in rats, fasting less than 24 hours causes no ulceration in stomach, but starvation (for 5 days) increases the number of mucosal ulcerations and aggravated acid-induced gastric erosions¹⁶⁻¹⁹. Finally, Ramadan fasting increases bleeding and coagulation time and causes in vitro decrease in the platelet responses of different aggregating agents (ADP, adrenaline and collagen)²⁰.

On the other hand, during Ramadan fasting, smoking and drinking alcohol which are risk factors of PUD and Mallory-Weiss syndrome are prohibited. Although, the overall estimation was an increment of AUGIB due to PUD with Ramadan fasting, there might be some other psycho-socio-environmento-spiritual factors that could affect the final outcome.

Does Ramadan actually increase the rate of AUGIB or not? The question was investigated in this study.

Methods

In a historical cohort study, all patients who had admitted to Alzahra hospital, Isfahan, Iran with AUGIB during a 3-month period (Ramadan month, one month before and one month later) for 3 consecutive years (2002 to 2004) were enrolled in the study. Approximately all patients were admitted during autumn season except the month before Ramadan of 2004 which included the last 15 days of September too. Alzahra is the only referral hospital of Isfahan province during the past 8 years for the management of AUGIB and emergency endoscopy. All patients files with AUGIB from the 10th day of Ramadan month to one month after it from 2002 to 2004 were reviewed and filled out the required information including the prognostic factors in a data sheet, designed for this purpose. Prognostic factors that affect outcome of AUGIB include, the cause of bleeding (most important), severity of the initial bleeding that is assessed by the transfusion requirement, age of the patient, concomitant diseases, onset of bleeding during hospitalization, emergency surgery, medical and conservative therapies and therapeutic endoscopic techniques²¹⁻²⁶.

The gathered information included age, sex, recent use of NSAIDs, recent use of anticoagulants, past history of smoking, underlying diseases, frequency distribution of causes of AUGIB (table 3), previous dyspeptic symptoms (epigastric pain and fullness, heart burn, nausea, early satiety and bloating), prognostic factors and outcome of AUGIB (re-bleeding, need to surgery, volume of infused packed cells, number of days in hospital and mortality rate). Then, they were contacted and questioned about their fasting status from one month before AUGI and, the number of fasting days.

For analytical purposes, the patients were divided into fasting group who were fasting, at least 10 days before admission and non-fasting group. The data were analyzed by SPSS software, using t-test, Chi-square and Mann-Whitney tests. Data expressed as mean \pm SD,

unless stated otherwise. P values less than 0.05 were statistically considered significant.

Results

Of patients with AUGIB (245 patients) from the 10th day of Ramadan to one month after from 2002 to 2004, we could contact 238 patients; 108 patients had been fasting at least 10 days (46%) and 128 patients had not been fasting any days (54%). Two patients had been fasting less than 10 days and they were eliminated from the study.

In fasting group, mean days of fasting was 20. They were 16-96 years old and their mean age was 48 ± 19 years; 70 males (64.8%) and 38 females (35.2%). Non-fasting group were 17-91 years with the mean age of 57 ± 19 years; 96 males (75%) and 32 females (25%).

Patients in fasting group were younger than non-fasting group (P<0.05) (table 2). Both groups were comparable regarding male/female ratio, predisposing factors for GI bleeding (recent consumption of NSAIDs and anti-coagulants and smoking) and previous underlying diseases that may promote GI bleeding (peptic ulcer disease, cirrhosis, CRF, COPD and coagulopathy) (table 2).

There was not any significant difference between two groups in AUGIB whether in Ramadan or one month later but, the number

of non-fasting patients with AUGIB was significantly higher than fasting patients in the month before Ramadan (table 1). Frequency of duodenal ulcer in fasting group (38%) was significantly more than that in non-fasting group (19.5%), which was significant only for 20-60 years old patients (P<0.05), and not for others. Conversely, the frequency of esophageal varices in non-fasting group (11.7%) was more than that in fasting group (3.7%) (P<0.05), which was significant for patients more than 40 years old (P<0.05). The frequency of other causes of AUGIB but erosive gastritis, in fasting group, was less than that in non-fasting group but, this difference was not significant (table 3). Gastric ulcer and erosive gastritis more frequently occurred in patients more than 60 years old in non-fasting group (P<0.05).

Fasting group showed more dyspeptic symptoms (38%) compared with non-fasting group (18.9%); (P = 0.001 and odds ratio = 2.62).

Regarding the outcome of AUGIB in both groups, there was no significant difference between them from the points of rebleeding in hospital, need to surgery, volume of infused packed cells and number of days in hospital (NS). But mortality rate was higher in non-fasting group (12.5%/3.7%) (table 4).

Table 1. The frequency of acute upper gastrointestinal bleeding in Ramadan.

Month→ Year↓	One month before Ramadan NF/F(n)	Month of Ramadan NF/F(n)	One month after Ramadan NF/F(n)
2002	59/0	28 (18)	29 (11)
2003	55/1	42 (30)	37 (18)
2004	68/1	43 (22)	35 (9)
Sum	182/2	113 (70)	101 (38)

NF/F (n); number of non-fasting group/fasting group

Table 2. Frequency of predisposing factors and underlying diseases of acute upper gastrointestinal bleeding in fasting and non-fasting groups.

Factor	Fasting group	Non-fasting group	P-value
Mean age (year)	48	57	0.001
Sex (male%/female%)	64.8/35.2	75/25	0.088
Recent use of NSAIDs (%)	21.3	28.1	0.227
Recent use of anticoagulant Drugs (%)	2.8	4.7	0.445
Previous smoking (%)	22.2	33.6	0.54
Underlying diseases (%)	17	28	0.861

Table 3. Frequency of the causes of acute upper gastrointestinal bleeding in

fasting and non-fasting groups.				
	Fasting group	Non-fasting group	P-value	Odds ratio
Duodenal ulcer (%)	38%	19.5%	0.002	2.52
Erosive gastritis(%)	17.6%	16.4%	0.809	1.09
Gastric ulcer (%)	13%	18%	0.292	0.68
Esophageal varices (%)	3.7%	11.7%	0.024	0.29
Mallory-Weiss (%)	2.8%	3.1%	0.876	0.886
Other causes (%)	12%	18%	0.207	0.625
Obscure AUGIB (%)	13%	13.3%	0.943	0.972
All above causes but duodenal ulcer and and erosive gastritis	44.4%	64.1%	0.543	0.449

Table 4. Outcome of acute upper gastrointestinal bleeding in fasting and non-fasting groups.

	Fasting group	Non-fasting group	P-value
rebleeding in hospital (%)	12%	14.1%	0.646
need to surgery (%)	8.3%	7.8%	0.884
mean volume infused packed cells (ml)	964.6	965.9	0.991
mean number of days in hospital (days)	4.68	4.81	0.954
mortality (%)	3.7%	12.5%	0.016

Discussion

It is observed that AUGIB has seasonal fluctuation with low prevalence in winter and an increase in spring and autumn with two peaks in April and October²⁷. The seasonal prevalence parallels that of duodenal ulcer bleeding, which follows a similar fluctuation²⁷. Approximately all of our patients were studied at autumn but one month before Ramadan from 2002 to 2004 was nearly within October. GI-bleeding due to DU was increased during Ramadan in our study, and more frequent dyspeptic symptoms in fasting group could be considered as alarm signs for patients who want to fast during Ramadan.

On the other hand, the frequency of GI-bleeding because of almost all other causes was decreased; this decrement was significant only in bleeding due to esophageal varices. Less frequent esophageal varices in fasting group could be explained by decrement in portal pressure during fasting. Patients with portal hypertension have very high azygos blood flow in comparison with healthy controls²⁸. Eating significantly increases azygos blood flow both in controls (38%) and in cirrhotics (27%)^{29, 30}. Therefore, compared with fasting

conditions, regular food intake plays probably an important role in the maintenance of effective hepatic blood flow in cirrhotic patients³¹ It seems therefore that fasting can decrease episodes of AUGIB due to esophageal varices.

Fasting group were younger than non-fasting group. This is obvious that older age group do not fast because of other concomitant diseases and disabilities that they have or because they take drugs. As mentioned above, duodenal ulcer was more common in fasting group and it occurred in younger patients while other causes of GI bleeding were more common in older patients.

Both fasting and non-fasting groups were similar regarding prognostic factors but, lower mortality rate in fasting group might be due to their younger age compared with non-fasting group. Mortality rates because of upper GI hemorrhage vary from 3.5% to 7% in the United States^{23, 26, 32}, and 14% in the United Kingdom³³.

Conclusion

Ramadan fasting can increase AUGIB by aggravating duodenal ulcer, but it does not make its prognosis poorer than non-fasting patients. Dyspeptic symptoms could be considered as

alarming signs for patients who want to fast during Ramadan. Therefore, we recommend caution in advising fasting to those with previous history of DU and dyspepsia, except with concurrent prescription of proton pump inhibitors as prophylaxis. Further evaluation of the effects of prophylactic drugs such as H₂-blockers and proton pump inhibitors and

Helicobacter eradication in preventing AUGIB due to duodenal ulcer should be done.

Acknowledgment

We are grateful to Dr. Ziba Farajzadegan and Dr. Abdolmahdi Baghai for their statistical consultations, and Dr. Movahed Abtahi for his comments on writing the article.

References

1. Baghareh chapter. *The Koran*. 2000.
2. Zafer O, Tulay B. **Effects of fasting during Ramadan on compliance with medical treatment.** *Letters to the editor. ASM* 1998; 18[5], 479.
3. Silverstein FE, Gilbert DA, Tedesco FJ, Buenger NK, Persing J. **The national ASGE survey on upper gastrointestinal bleeding. I. Study design and baseline data.** *Gastrointest Endosc* 1981; 27:73-79.
4. John DV, William DC, James MS. **Pathogenesis of peptic ulcer disease.** In: Yamada T, editor. *Textbook of gastroenterology*. Philadelphia: Lippincott Williams & Wilkins, 2003: 1326-1328.
5. Dronfield MW, Langman MJ, Atkinson M, Balfour TW, Bell GD, Vellacott KD et al. **Outcome of endoscopy and barium radiography for acute upper gastrointestinal bleeding: controlled trial in 1037 patients.** *Br Med J (Clin Res Ed)* 1982; 284:545-548.
6. Green FW, Jr., Kaplan MM, Curtis LE, Levine PH. **Effect of acid and pepsin on blood coagulation and platelet aggregation. A possible contributor prolonged gastroduodenal mucosal hemorrhage.** *Gastroenterology* 1978; 74:38-43.
7. Khuroo MS, Yattoo GN, Javid G, Khan BA, Shah AA, Gulzar GM et al. **A comparison of omeprazole and placebo for bleeding peptic ulcer.** *N Engl J Med* 1997; 336:1054-1058.
8. Hakkou F, Tazi A, Iraqui L, Celice-Pingaud C, Vazier J. **[The observance of Ramadan and its repercussion on gastric secretion].** *Gastroenterol Clin Biol* 1994; 18:190-194.
9. Iraki L, Bogdan A, Hakkou F, Amrani N, Abkari A, Touitou Y. **Ramadan diet restrictions modify the circadian time structure in humans. A study on plasma gastrin, insulin, glucose, and calcium and on gastric pH.** *J Clin Endocrinol Metab* 1997; 82:1261-1273.
10. Iraki L, Abkari A, Vallot T, Amrani N, Khelifa RH, Jellouli K et al. **[Effect of Ramadan fasting on intragastric pH recorded during 24 hours in healthy subjects].** *Gastroenterol Clin Biol* 1997; 21:813-819.
11. Donderici O, Temizhan A, Kucukbas T, Eskioglu E. **Effect of Ramadan on peptic ulcer complications.** *Scand J Gastroenterol* 1994; 29:603-606.
12. Hosseini-asl K, Rafeian-kopaei M. **Can patients with active duodenal ulcer fast Ramadan?** *Am J Gastroenterol* 2002; 97:2471-2472.
13. Mallk GM, Mubarik M, Hussain T. **Acid peptic disease in relation to Ramadan fasting: a preliminary endoscopic evaluation.** *Am J Gastroenterol* 1995; 90:2076-2077.
14. Mehdi A, Ajmi S. **[Effect of the observance of diurnal fast of Ramadan on duodenal ulcer healing with lansoprazole. Results of a prospective controlled study].** *Gastroenterol Clin Biol* 1997; 21:820-822.
15. Kadri N, Tilane A, El Batal M, Taltit Y, Tahiri SM, Moussaoui D. **Irritability during the month of Ramadan.** *Psychosom Med* 2000; 62:280-285.
16. Auguste LJ, Angus L, Stein TA, Wise L. **Starvation and mucosal prostaglandin-E₂ in gastric stress ulceration.** *Crit Care Med* 1988; 16:610-611.
17. Hung CR, Neu SL. **Acid-induced gastric damage in rats is aggravated by starvation and prevented by several nutrients.** *J Nutr* 1997; 127:630-636.
18. Moron F, Nagy L, Mozsik G. **The gastric fundic mucosal energy systems and the HCl-induced lesions in rats in dependence on starvation.** *Acta Med Hung* 1984; 41:253-258.
19. Nakagawa K, Okada A, Kawashima Y. **Acute gastric mucosal lesions--a new experimental model and effect of parenteral nutrition.** *JPEN J Parenter Enteral Nutr* 1985; 9:571-582.
20. Aybak M, Turkoglu A, Sermet A, Denli O. **Effect of Ramadan fasting on platelet aggregation in healthy male subjects.** *Eur J Appl Physiol Occup Physiol* 1996; 73:552-556.

21. Corley DA, Stefan AM, Wolf M, Cook EF, Lee TH. **Early indicators of prognosis in upper gastrointestinal hemorrhage.** *Am J Gastroenterol* 1998; 93:336-340.
22. Graham DY, Smith JL. **The course of patients after variceal hemorrhage.** *Gastroenterology* 1981; 80:800-809.
23. Longstreth GF. **Epidemiology of hospitalization for acute upper gastrointestinal hemorrhage: a population-based study.** *Am J Gastroenterol* 1995; 90:206-210.
24. Lumsden K, MacLarnon JC, Dawson J. **Giant duodenal ulcer.** *Gut* 1970; 11:592-599.
25. Terdiman JP, Ostroff JW. **Risk of persistent or recurrent and intractable upper gastrointestinal bleeding in the era of therapeutic endoscopy.** *Am J Gastroenterol* 1997; 92:1805-1811.
26. Yavorski RT, Wong RK, Maydonovitch C, Battin LS, Furnia A, Amundson DE. **Analysis of 3,294 cases of upper gastrointestinal bleeding in military medical facilities.** *Am J Gastroenterol* 1995; 90:568-573.
27. Thomopoulos KC, Katsakoulis EC, Margaritis VG, Mimidis KP, Vagianos CE, Nikolopoulou VN. **Seasonality in the prevalence of acute upper gastrointestinal bleeding.** *J Clin Gastroenterol* 1997; 25:576-579.
28. Bosch J, Mastai R, Kravetz D, Bruix J, Rigau J, Rodes J. **Measurement of azygos venous blood flow in the evaluation of portal hypertension in patients with cirrhosis. Clinical and haemodynamic correlations in 100 patients.** *J Hepatol* 1985; 1:125-139.
29. Betal D, Hughes ML, Whitehouse GH, Roberts N. **Postprandial decrease in splenic volume demonstrated by magnetic resonance imaging and stereology.** *Clin Anat* 2000; 13:404-409.
30. Sugano S, Yamamoto K, Takamura N, Momiyama K, Watanabe M, Ishii K. **Azygos venous blood flow while fasting, postprandially, and after endoscopic variceal ligation, measured by magnetic resonance imaging.** *J Gastroenterol* 1999; 34:310-314.
31. Tochio H, Kudo M, Nishiura S, Okabe Y. **Intrahepatic spontaneous retrograde portal flow in patients with cirrhosis of the liver: reversal by food intake.** *AJR Am J Roentgenol* 2001; 177:1109-1112.
32. Cooper GS, Chak A, Harper DL, Pine M, Rosenthal GE. **Care of patients with upper gastrointestinal hemorrhage in academic medical centers: a community-based comparison.** *Gastroenterology* 1996; 111:385-390.
33. Rockall TA, Logan RF, Devlin HB, Northfield TC. **Risk assessment after acute upper gastrointestinal haemorrhage.** *Gut* 1996; 38:316-321.